UTILITY OF RS, GIS AND GPS IN AGRICULTURE AND ALLIED SCIENCES

Authors

Er. Deepak Chouhan

Dr. Rajveer Singh Yadav

Er. Jaypalsinh Chavda

Dr. Kunal Bhelave

Mr. Vaqar Malik

Publishing in 2024

Copyright ©2023 by Royal Book Publishing. All Rights Reserved

Published by
Royal Book Publishing
21/11, K M Nagar,
Ayodhiyapatinam, Salem.
Tamil Nadu-636103
Cell:+91 7598141199
Email: contact@royalbookpublishing.com
Website: www.royalbookpublishing.com

Published in India.

International Standard Book Number (ISBN) : 9788119821327

No part of this Edited book (E - book) may be reproduced in any form, by photostat, microflim, xerography, or any other means, or incorporated into any information retrievel system, electronic or mechanical, without the written permission of the publisher. For permission to photocopy or use material electronically from this work, please contact the Book Publisher or Book Editor through the e-mail.

Trademark notice: Any brand names and product names mentioned in this book are subjected to trademark, brand or patent protection and are trademarks or registered trademark of their respective holders. The use of brand names, product names, common names, trade names, product description, etc. even without a particular marking in this work is in no way to be constructed to mean that such names any be regarded as unrestricted in respect of trademark and brand protection legislation and could thus be used by anyone.

Royal book publishing also publishes it's books in a variety of Electronic formats. Some content that appears in print may not be available in Electronic formats. For more information visit our publication website www.royalbookpublishing.com

9 788119 821327

Preface

The field of agriculture is currently experiencing a technological revolution, with remote sensing, GIS, and GPS playing a pivotal role in this transformation. By utilizing these technologies to gather extensive data from various sources such as land, air, and space, remote sensing provides farmers and researchers with unprecedented insights into crop growth, soil moisture levels, and weather patterns. These valuable insights enable farmers to make more informed decisions, optimize their yields, and minimize their environmental impact. The integration of remote sensing, GIS, and GPS has been instrumental in the field of agriculture and crop management, allowing for local, regional, and even global scale estimation and analysis. This has been made possible through the simultaneous development of crop models and remote sensing techniques.

The book titled "Utility of RS, GIS and GPS in Agriculture and Allied Science" serves as a comprehensive guide to the current state of remote sensing, geographic information system and global position system in agriculture allied science fields. It thoroughly explores the latest scientific advancements and their practical applications in innovative solutions for farmers. The book covers a wide range of topics, including satellite farming, machine learning, big data, and the integration of remote sensing with other precision agriculture technologies. One of the most exciting aspects of remote sensing is its ability to support sustainable and efficient agricultural practices. The book dedicates sections to how remote sensing, GIS and GPS is being utilized to enhance crop yield, reduce water and fertilizer usage, detect and manage pests and diseases and natural resources management. It

also emphasizes the potential of remote sensing in supporting precision conservation and precision farming.

This book is an essential resource for researchers, practitioners, and policymakers alike. Whether you are a farmer seeking to improve your crop management techniques or a researcher working towards groundbreaking advancements in precision agriculture, this book serves as an indispensable guide to the current state of remote sensing in precision agriculture. We firmly believe that this book will prove to be an invaluable resource for anyone involved in the field.

Er. Deepak Chouhan
Dr. Rajveer Singh Yadav
Er. Jaypalsinh Chavda
Dr. Kunal Bhelave
Mr. Vaqar Malik

TABLE OF CONTENT

SL. NO	TITLE NAME	PAGE. NO
1	**Remote Sensing in Agriculture** Deepak Chouhan, Aniket Rajput	1-12
2	**Geographic Information System in Agriculture** Deepak Chouhan, Aniket Rajput	13-27
3	**Principles of Geographic Information Systems** Deepak Chouhan, Aniket Rajput	28-37
4	**Environmental Geoinformatics Monitoring and Management** Dr. K.M. Gojiya, Er. B.A. Karangiya, Prof. M.J. Gojiya	38-48
5	**Uses of GIS, GPS and VRA in Precision Agriculture** Tanveer Ahmad Ahngar, Ruby Jan, Zahoor Ahmad Dar	49-60
6	**Role of Geospatial Technology in Agriculture** Dr. K.M. Gojiya, Er. B.A. Karangiya, Dr. S. K. Chavda	61-71
7	**Applications of Remote sensing, GIS and GPS in Floriculture** G.N. Nagajyothi, C.T. Pratheeksha, T.V. Udaya	72-92
8	**Remote Sensing and Geographic Information System in Forestry** Nikita Kumari, Ajit Tippannavar	93-109
9	**Application of RS, GIS and GPS in Fruit Science**	110-132

	Anju Yadav, Dhaneshra Kumari Gurjar	
10	**Application of RS, GIS and GPS in Plant Pathology** Ravichandra, Sudarshan Varma, Channabasava	**133-157**
11	**Application of RS, GIS and GPS in Soil Resource Mapping** Sudarshan Varma, K. Vilakar, Ravichandra	**158-179**
12	**Optical Sensing Technology (OST) for Detection of Plants Diseases** Satendra Kumar Sharma, Nentia Chib, Deepak Kumar	180-208
13	**Image Classification, GIS Integration and Acreage Estimation for Land Management** Dhaval Kirankumar Dwivedi	209-222
14	**Application of RS, GIS and GPS in Agronomy** Tanveer Ahmad Ahngar, Ruby Jan, Zahida Rashid	223-238
15	**Geospatial Technologies for Crops and Soils** Seerat Jan , Tanveer A. Ahanger, Shahid Rasool	239-257
16	**Geoinformatics, Nanotechnology and Precision Farming** Insha Jan, Tanveer Ahmad Ahngar, Arzeena Shabir	258-270

CHAPTER – 1

Remote Sensing in Agriculture

Deepak Chouhan[1] , Aniket Rajput[2]

[1] Scientist (Agricultural Engineering), Krishi Vigyan Kendra, Shahdol (M. P.)

[2] M. Tech (SWE), College of Agricultural Engineering (M. P.)

Abstract

The application of remote sensing in quantifying the crop health status is trending. Sensors are served as early warning systems for countering climatic or biological aberrations before having negative impacts on crop yield. Remote sensing applications play a significant role in the agriculture sector for evaluating plant health, yield and crop loss (%) estimation, irrigation management, identification of crop stress, weed and pest detection, weather forecasting, gathering crop phenological information etc. Remote sensing inputs is used for forecasting of crop yields in conjunction with crop simulation models is getting popular day by day for its potential benefits. Remote sensing reduces the collection of amount of field data and improves the precision of the estimates. Remote sensing is used for crop stress caused by biotic and abiotic factors can be monitored and quantified. Monitoring of vegetation cover for acreage estimation, weed management through precision agriculture, mapping and monitoring drought conditions and maintenance of vegetation health, checking of nutrient and moisture status of the field, assessment of crop condition under stress-prone environment, measurement of crop evapotranspiration, gathering and transferring predictions of atmospheric dynamics through different observational satellites are the major agricultural applications of remote sensing technologies. Normalized difference vegetation index (NDVI), leaf area index (LAI), General Yield Unified Reference Index (GYURI), and

ISBN : 9788119821327

vegetation condition index (VCI) are some of the indices that have been used for mapping and monitoring drought and assessing vegetation health and productivity. Remote sensing with other advanced technologies like geographical information systems (GIS), Global position system (GPS) are playing a massive role in the assessment and management of several agricultural activities. For improving agricultural economy the state or district-level information systems based on available remote sensing information are required to be utilized efficiently.

Keywords:- Remote sensing, Agriculture, Normalized Difference Vegetation Index, Yield

Introduction :-

Remote sensing application is the art and science of gathering information about the objects or area of the real world at a distance without coming into direct physical contact with the object under study. Remote sensing technique is monitoring for the earth's resources that combines space technology with ground observations to provide greater precision and accuracy. The utilization of electromagnetic spectrum (visible, infrared, and microwaves) for measuring the earth properties is the principle behind remote sensing. In figure 1 show that the typical responses of the wavelength regions are different, so they are used for distinguishing the vegetation, bare soil, water and other features. The crop growth monitoring, water resources mapping and water status under field conditions, land use pattern and land cover changes, disease and insect infestation monitoring, harvesting date forecasting and yield prediction, precision farming, and weather forecasting all are possible though remote sensing. Remote sensing data can substantially aid in the monitoring the earth's surface features by providing timely, synoptic, cost-efficient, and repetitive information about the earth surface (Justice et al. 2002). Remote sensing inputs are combined with crop simulation

ISBN : 9788119821327

models are useful for crop yield forecasting, because ground-based and air-based platforms are time-consuming and have limited applications. The space-based satellite technologies are becoming increasingly important for gathering spatio-temporal metrological and crop condition information to supplement existing approaches.




Figure 1 :- Typical spectral reflectance curve for vegetation, dry bare soil and water

Remote sensing technology in contemporary agriculture, offering useful data to both farmers and researchers. The efficiency and effectiveness of agricultural techniques have substantially increased as a result of the deployment of remote sensing technology. The remote sensing technique gathers information about crops, soil, water, and other vital agricultural components using satellites, planes, and drones. Remote sensing data is used for produce images and maps that may be examined to spot potential issues and possibilities. Its making informed judgments about crop management, fertilization, irrigation, and pest control with this knowledge ultimately results in higher yields and lower costs. In precision agriculture the development is the technology to improve crop yield while minimizing waste and maximizing resource utilization, is another area where remote sensing is having a huge

ISBN : 9788119821327

impact. Bajaj et. al., 2023 also studied the application of RS, GIS and GPS in Agriculture in India.




Figure 2 :- Remote sensing modern Application in Agriculture.

Crop Inventory :-

Remote sensing can play an important role in agriculture by providing timely spectral information which can be used for assessing the bio-physical indicators of plant health. The physiological changes that occur in a plant as a result of stress may modify the spectrum reflectance/ emission properties, allowing remote sensing tools to identify stress (Menon, 2012). In the crop growing season crop monitoring is required at regular intervals to take suitable steps and to determine the likelihood of output loss due to any stressful event. The

ISBN : 9788119821327

different factors influence crop growth phases and development, including planting date, available soil moisture, air temperature, day length, and soil condition. The circumstances of the plants and their productivity are determined by these elements. In the corn crop yields, is example, might be harmed by very high temperatures during pollination. As a result, knowing the temperature at the time of corn pollination could aid forecasters in making more accurate predictions about maize yields (Nellis et al., 2009). Satellite data is used to monitoring drought in recent years, and the use of the Normalized Difference Vegetation Index (NDVI) and Vegetation Condition Index (VCI) for identifying agricultural drought in various regions with varying ecological conditions has been accepted globally (Anyamba et al., 2001; Ji and Peters, 2003).

Monitoring of vegetation cover :-

Remote sensing is play a vital role in the area of crop classification, crop acreage estimation and yield assessment, for this work aerial pictures and digital image processing techniques were used by several research investigations. Remote sensing reduces the amount of field data required and enhances the accuracy of estimates (Kingra et al., 2016). A significant improvement in crop characterization, discrimination, modeling, and mapping is known to be possible with hyperspectral data, compared with broadband multispectral remote sensing (Thenkabail et al., 2013). For crop health and productivity, some remote sensing approaches place a greater emphasis on physical factors of the crop system, such as nutrient stress and water availability. Normalized Difference Vegetation Index is the most commonly used index to assess the vegetation condition (Rouse et al., 1974, Calvao and Palmeirim, 2004, Wallace et al., 2004). The normalised difference vegetation index (NDVI), vegetation condition index (VCI), leaf area index (LAI), General Yield Unified Reference Index (GYURI), and Temperature Crop Index (TCI) are all examples of indices that have been

ISBN : 9788119821327

used for drought mapping and monitoring as well as vegetation health and productivity assessment (Doraiswamy et al., 2003).

Weed identification and management :-

Effectively weed management by precision weed management techniques and precision agriculture combined with remote sensing is promising technology today. For weed mapping ground surveying is very labour intensive and time-consuming, remote sensing is a good method for mapping weed information for a specific location. However, the potential applications in weed detection for site–specific weed management is image–based remote sensing (Johnson et al., 1997; Moran et al., 1997; Lamb et al., 1999). Remote sensing technology uses in standing crops for difference in spectral reflectance qualities between weeds and crops to identify weeds and aids in the compilation of weed maps in the field, allowing for the application of site-specific and need-based herbicides for weed management.

Nutrient and water status :-

Nutrient and water stress management are the two most important domains where remote sensing and GIS can be used to help with precision farming. The remote sensing and GIS is to detect nutritional challenges aids us in site-specific nutrient management, allowing us to lower cultivation costs and boost fertiliser usage efficiency for crops. Precision farming technologies can help in semi-arid and arid regions to make more efficient use of water. Drip irrigation, is example, can be used in conjunction with remotely sensed data such as canopy air temperature differential to boost water use efficiency by lowering runoff and percolation losses (Das and Singh, 1989). Remote sensing data is used for collect information on agricultural water demand, water use, soil moisture state, and related crop growth at various crop-growing stages. For example, Bandara (2003) evaluated the effectiveness of three big irrigation projects in Sri Lanka using NOAA satellite data. Estimates

ISBN : 9788119821327

of crop-water use based on remote sensing were compared to actual water availability to assess irrigation efficiency in this study.

Agro metrological Application :-

Agriculture is highly influenced by climatic and metrological phenomena. The metrological data are collected using various observation station point networks in space. The ability of conventional agrometrological techniques to utilize this data for forecasting yields and real-time agricultural monitoring is severely constrained. Several fundamental agrometrological characteristics can now be measured accurately and often thanks to satellite metrology (example surface albedo, surface temperature, evapotranspiration, solar radiation, rain fall). The main inputs for the agrometeorology model is included minimum and maximum temperatures, considerable rainfall occurring at biweekly intervals, and so on. The geostationary satellites is used for weather and climate remote sensing is recognized as the single most important development in the last 25 years for tracking the Earth's vegetation, weather, and climate. The two most crucial inputs for agricultural meteorology are data on vegetation and meteorological.

Water resource management :-

Recent decades seen that the availability of water resources in global and regional decrease, which calls for careful management of cutting-edge technology. Remote sensing is one of the efficient methods for evaluating and keeping track of water resources. In order to provide more precise estimates of the information needed for applications involving water resources, hyperspectral remote sensing is emerging as a more thorough method of examining spatial, spectral, and temporal variations. Active and passive microwave remote sensing provides a unique capability to obtain observations of soil moisture at global and regional scales that help satisfy the science and application needs for hydrology (Njoku and Entekhabi, 1996, Schugge et al., 2002).

ISBN : 9788119821327

Groundwater is one of the most important natural resources because it promotes economic growth, ecological diversity, and human health. Our ecosystems and the survival of future generations are in danger due to the excessive use of this essential resource.

Infestation of pests and diseases :-

Remote sensing has become an indispensable technique for monitoring and quantifying crop stress caused by biotic and abiotic variables. Remote sensing methodologies must be refined for identifying insect breeding grounds, formulating plans to limit their spread, and applying effective control measures. In order to relate differences in spectral responses to yellowing of leaves, chlorosis, and foliage reduction over a specific time period, the remote sensing approach has been used in insect defoliation assessment and monitoring and This approach makes the assumption that these differences can be correlated, classified, and interpreted (Franklin, 2001). The remote sensing applications has included detecting and mapping defoliation, characterization of pattern disturbances etc. and providing data to pest management decision support system (Lee et al., 2010). The Landasat 5 TM image can be used to accurately detect and quantify disease for site-specific Wheat Streak Mosaic disease management in the wheat crop (Mirik et al., 2013).

Land use and land cover :-

In the land use/land cover mapping involved identification of surface features at different scales and their hierarchical classification which is important in the study of global change. The human activities, including deforestation, biodiversity loss, and global warming is environmental issues to significantly impacted on land usage and land cover. The land use/ land cover maps are created from satellite imagery were compared pixel by pixel using a change detection approach. Mapping of land use and land cover has always been crucial to all

ISBN : 9788119821327

geographical research since it contains the fundamental data on features that are present on the surface, including their area, position, shape, and pattern.

Crop Yield forecasting :-

Crop yields have been forecasted using remote sensing, generally based on statistical-empirical connections between yield and vegetation indices (Thenkabail et al., 2002, Casa and Jones 2005). For critical national food policy planning crop production statistics prior to harvest. Crop production forecasting is an important aspect for Crop yield predictability. The crop yield is influenced by a number of factors, including crop type, field water and nutrient status, weed infestation, pest and disease infestation, and weather conditions. The crop condition and performance are indicated by the growth and decay of the spectral response curve.

Precision agriculture :-

A growing number of scientists, engineers, and large-scale crop growers are using remote sensing technology as part of precision farming (Liaghat and Balasundram, 2010). Precision farming's major goal is to minimise cost of cultivation, improve management, and increase resource efficiency by using data collected by sensors installed in farm machinery. Precision farming's most advanced component is variable rate technology (VRT), Sensors are fitted on moving farm machinery, which contain a computer that gives input recommendation maps and thereby controls the application of inputs based on GPS receiver data (NRC, 1997).

Conclusion :-

Remote sensing is useful in assessing various abiotic and biotic stresses in different crop and also very useful in detecting and management of various crop issues even at small farm holdings. To effectively utilize the information on crops for improvement of economy

ISBN : 9788119821327

there is a need to develop state or district level information system based on available information on various crops derived from remote sensing and GIS approaches. The governments can use remote sensing data in order to make important decisions about the policies they will adopt or how to tackle national issues regarding agriculture. A new and nontraditional remote sensing application involves the implanting of nano-chips in plant and seed tissue that can be used in near-real time to monitor crop. Clearly, these and other new approaches will reinforce the importance of remote sensing in future analysis of agricultural sciences.

References :-

1. Anyamba A, Tucker CJ, Eastman JR. NDVI anomaly patterns over Africa during the 1997/98 ENSO warm event. Int. J. Remote Sens. 2001;22:1847-1859.
2. Bajaj, S., Mishra , S., Chouhan , D., Bora , S. S., Parashar , S., Bargi, U. S., & Billore , V. (2023). A Comprehensive Review of Application of RS, GIS and GPS in Agriculture India. International Journal of Environment and Climate Change, 13(11), 2597–2608.
3. Bandara, KMPS. Monitoring irrigation performance in Sri Lanka with high frequency satellite measurements during the dry season. Agricultural Water Management. 2003; 58 (2): 159–170.
4. Calvao T. and Palmeirim, J.M. (2004). Mapping Mediterranean scrub with satellite imagery: biomass estimation and spectral behaviour. International Journal of Remote Sensing, 25: 3113–3126.
5. Das DK and Singh G. Estimation of evapotranspiration and scheduling irrigation using remote sensing techniques. Proc. Summer Inst. On agricultural remote sensing in monitoring crop growth and productivity, IARI, New Delhi, 1989, 13-17.
6. Doraiswamy, P. C., Sinclair, T. R., Hollinger, S., Akhmedov, B., Stern, A. and Prueger, J. (2005). Application of MODIS derived parameters for regional crop yield assessment. Remote Sens Environ., 97: 191-202.

ISBN : 9788119821327

7. Franklin S. Remote Sensing for Sustainable Forest Management. Lewis publisher, Boca Raton, Florida, 2001, 407.
8. Ji L, Peters AJ. Assessing vegetation response to drought in the northern Great Plains using vegetation and drought indices. Remote Sens Environ. 2003;87:85-98.
9. Johnson GA, Cardina J, Mortensen DA. Site–specific weed management: Current and future direction. In: The State of Site–Specific Management for Agriculture. Wiley Pubblisher; 1997. pp. 131-147.
10. Justice, C. O., Townshend, J. R. G., Vermata, E. F., Masuoka, E., Wolfe, R. E., Saleons, N., Ray, D. P. and Morisette, J. T. (2002). An overview of MODIS Land data processing and product status. Remote Sens Environ., 83: 3 15.
11. Lamb DW, Weedon MM, Rew LJ. Evaluating the accuracy of mapping weeds in seeding crops using air-borne digital imaging: Avena spp.in seeding triticale. Weed Research. 1999;39(6): 481-492.
12. Lee W, Alchanatis V, Yang C, Hirafuji M, Moshou D and Li C. Sensing technologies for precision specialty crop production. Computer and Electronic in Agriculture, (2010); 74: 2-33.
13. Liaghat S, Balasundram SK. A review: The role of remote sensing in precision agriculture. American Journal of Agricultural and Biological Sciences. 2010;5 (1):50-55.
14. Menon ARR. Remote sensing applications in agriculture and forestry. A paper from the proceedings of the Kerala environment congress, 2012, pp. 222-235.
15. Mirik, M., Ansley, J., Price, J. A., Workneh, F., Rush, C. M., Remote Monitoring of Wheat Streak Mosaic Progression Using Sub-Pixel Classification of Landsat 5 TM Imagery for Site Specific Disease Management in Winter Wheat. Advances in Remote Sensing., 2: 16-28.

ISBN : 9788119821327

16. Moran MS, Inoue Y, Barnes EM. Opportunities and limitations for imagebased remote sensing in precision crop management. Remote Sensing of Environment. 1997;61(3):319-246.
17. Nellis MD, Pricey KP, Rundquist D. Remote Sensing of Cropland Agriculture. The SAGE Handbook of Remote Sensing, 2009, (26).
18. Njoku, E. G. and Entekhabi, D., "Passive microwave remote sensing of soil moisture," Journal of Hydrology, vol. 184, no. 1-2, pp. 101–129, 1996.
19. NRC. Precision Agriculture in the 21st Century Geospatial and Information Techniques in Crop Management. Washington, DC: National Academy Press; 1997. p. 149.
20. Rouse JW, Haas RH, Schell JA, Deering DW. Monitoring vegetation systems in the Great Plains with ERTS. Third ERTS Symposium, NASA. 1973;351 (1):309-317.
21. Schmugge, T. J., Kustas, W. P., Ritchie, J. C., Jackson, T. J., and Rango, A., "Remote sensing in hydrology," Advances in Water Resources, vol. 25, no. 8-12, pp. 1367–1385, 2002.
22. Thenkabail PS, Mariotto I, Gumma MK, Middleton EM, Landis DR, Huemmrich KF. Selection of hyperspectral narrow bands (HNBs) and composition of hyperspectral two band vegetation indices (HVIs) for biophysical characterization and discrimination of crop types using field reflectance and hyperion/EO-1Data. IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing. 2013; 6 (2):427-439.
23. Wallace, J.F., Caccetta, P.A. and Kiiveri, H.T. (2004). Recent developments in analysis of spatial and temporal data for landscape qualities and monitoring. Australian Ecology, 29: 100–107.

ISBN : 9788119821327

CHAPTER – 2

Geographic Information System in Agriculture

Deepak Chouhan[1] , Aniket Rajput[2]

[1] Scientist (Agricultural Engineering), Krishi Vigyan Kendra, Shahdol (M. P.)

[2] M. Tech (SWE), College of Agricultural Engineering (M. P.)

Abstract

In recent centuries, the advent of technological innovations has greatly enhanced our ability to increase agricultural production in order to meet the growing demands of the global population. While the fourth digital revolution in agriculture has been made possible by advancements in digital technologies, we are also faced with various challenges such as limited arable land, depleting water resources, and the impact of climate change. These challenges emphasize the urgent need for unprecedented measures to ensure agricultural resilience and support the world population. One technology that has played a crucial role in monitoring crops and implementing effective management practices to enhance crop productivity is the Geographic Information System (GIS). Alongside other partner technologies like remote sensing, global positioning system, artificial intelligence, computational systems, and data analytics, GIS has proven to be instrumental in various aspects of agriculture, ranging from land-use planning to monitoring crop-soil-yield and post-harvest operations. By combining GIS with digital technologies and exploring new and emerging applications, precision farming and sustainable food production goals are being realized. The integration of these technologies enables farmers to make informed decisions and

ISBN : 9788119821327

implement targeted strategies, ultimately leading to improved agricultural practices and the achievement of sustainable food production.

Keywords :- Geographic Information System (GIS), Water resource management, Precision farming, Crop monitoring, Remote sensing**.**

Introduction :- The world population is projected to grow close to 10 billion by 2050, supply the sufficient food for those population we need to produce about 50% more food compared to 2013 production to meet the global demand (FAO 2017). In this goal we needs to be met while facing the challenges of climate change, the limited scope of arable land expansion, and dwindling water resources. The expected food production also needs to incorporate different agricultural practices for sustainable management of croplands to conserve water resources, preserve soil health, and encompass biodiversity (Gomiero, 2016). For considering these challenges and constraints in achieving food production its need to monitoring of crop growth and crop health and timely interventions to maintain or improve crop productivity while reducing wastage of inputs and resources, this goal achieved by advances in sensors, communication technologies, computational systems, and powerful data analytics. This technologies is efficient use of agricultural inputs and reduce environmental losses to increased and sustainable production for achieving food security. Achieving this goal through monitoring of soil and crop and providing data driven insights for targeted or site-specific management of crops productivity several tools and technologies use as Global Positioning System (GPS), geographic information system (GIS), Artificial Intelligence (AI), remote sensing

ISBN : 9788119821327

(RS), Internet of Things (IoT), and Big Data Analytics (Delgado et al., 2019).

Geographic Information System (GIS), is a key foundational technology, is defined as a powerful system comprising tools for the collection, storage, and retrieval of data at will, as well as analyzing, transforming, and displaying the spatial data for a specific purpose (Burrough & Mcdonnel, 1998, Delgado et al., 2019). GIS is provides the spatial context and information from different features data layer. It is a tool to manipulate spatial and non-spatial data and presents them through different map formats as intuitive and illustrative. GIS has been making an impact in diverse domains that include geography, environmental sciences, natural resources, forestry, agriculture, food, manufacturing, banking, and health services (Soomro, 2015). In the recent decades application of GIS tools is increase for applications in agriculture at local, regional, national, or global scales. These application along with remote sensing, GPS, and data analytics to understanding the crops and soil measures. Many studies have reported the use of GIS for diverse applications in different crops (Gebeyehu, 2019). The readers are strong appreciation for the role of GIS technology in agriculture, here we have reviewed the most widely used and emerging applications of this GIS technology, either by itself or in combination with other partner technologies, and how it making major impacts on agricultural productivity and supply chains.

Technologies with GIS :- The use of GIS and two other key geospatial technologies in combination is responsible for the power and impactful contributions of GIS in different domains. These three technologies play an important role in realizing the goals of applications and, therefore, is briefly described below.

ISBN : 9788119821327

Geographic information system (GIS) :- GIS is considered the brain of precision agriculture due to its role in supporting the collection, storage, retrieval, and analysis of trait and location data, and the utility of data-driven solutions, especially in site-specific management (Kumar, et al., 2017). Digital GIS maps differ from traditional maps in that they contain multiple layers of information, with each layer containing information about a specific attribute such as soil testing, rainfall, nutritional status, pest infestation, yield, or Provide a map. Additionally, GIS provides analytics. The ability to use statistical tools and geospatial analysis allows the extraction of interrelationships between attributes, and the resulting insights are valuable for decision-making regarding management practices.

Global positioning system (GPS):- The latitude, longitude, and elevation of a location are provided by this satellite-based positioning and navigation system, which makes it possible to determine positional information. Farmers and researchers can reliably identify fields, map field boundaries, water bodies, infested or problematic areas in the field, and understand the relationship to several other attributes both inside and outside the boundaries of a given field thanks to the location data gathered by GPS receivers. Precision agriculture is all about lowering input costs and increasing output by applying water, fertilisers, herbicides, and pesticides just where needed. This kind of high-fidelity field mapping makes this possible.

Remote sensing (RS) :- Crop monitoring and interventions to increase farm output have been transformed by remote sensing's many uses and technologies in agriculture (Sishodia, et al., 2020 & Bajaj, et al., 2023). RS is essential for achieving the objectives of precision agriculture, together with GPS, GIS, and other instruments. This combination is essential for

ISBN : 9788119821327

enabling a number of applications, such as soil mapping, crop growth monitoring, estimation of soil moisture and fertility, detection of biotic (drought and flood) and abiotic (pests and diseases), and yield estimation, that form the basis for site-specific management of fields.

GIS applications :- Considered the fourth revolution in agriculture, the advent of digital agriculture has completely changed the way farming is done, thanks to advances in geospatial technology, sensors, artificial intelligence, robotics, and other tools and technologies. Imagery and non-imagery data and spatial context are required to accurately identify problem areas within agricultural fields and to monitor and manage every step of the entire agricultural value chain. GIS, with its component tools and analysis modules, as well as data collected by partner technologies such as remote sensing and GPS, provides the intuition of information for data-driven decision-making to improve crop productivity. Provides accurate and clear visualization. GIS has long been used in agricultural applications, but advances in technology have rapidly increased its use in recent years. Some of the most common and new applications are listed and discussed below.

Water resource management :-

Adequate water supply is a fundamental requirement to meet the food production needs of a growing world population. As mentioned earlier, farmers are responsible for feeding approximately 10 billion people in 2050 and will need to increase food production by 50% compared to 2013 (FAO 2017). The availability of clean water is decreasing and relying solely on rain is not a profitable option for farmers (Consoli & Vanella, 2014). Geospatial technology supported by remote sensing has already proven itself in water resource management

ISBN : 9788119821327

(Elhag, 2016). Many researchers have supported the potential of GIS in groundwater management. Tripathi et al., 2003, integrate the MODFLOW groundwater model into the GIS information system for watershed prioritization. Sikdar et al., 2004, GIS and remote sensing are combined to delineate zones of groundwater potential. Determining the suitability of irrigation for a particular geographic location is one of the most common uses of GIS. The ISAREG irrigation planning model was integrated into GIS with the aim of producing efficient irrigation planning advice and identifying practices to account for water conservation and salinity management (Fortes, et al., 2005). These results indicate that the model for the selection of irrigation plans and water-saving measures in both the wet and dry seasons is successful. Although intensification of irrigation is beneficial for food production, it has been pointed out that soil salinization and waterlogging are the main drawbacks of intensification of irrigated agriculture, and the use of GIS and remote sensing techniques to monitor problem areas. The emphasis will be on usage to plan conservation and preventive measures.

Land suitability assessment and land use planning :-

Geographic Information System (GIS) serves as a valuable tool for evaluating land quality for various purposes. When it comes to land use planning, the use of a Multi-criteria decision-making (MCDM) approach utilizing GIS is widely favored by researchers. These researchers make use of various GIS features such as the distribution of soil types, soil texture mapping, groundwater levels, soil fertility, soil pollution, soil hydraulic conductivity, slope, soil texture, depth to water-table, and groundwater electrical conductivity. Additionally, they also consider climate conditions, topography, and satellite data to analyze the intricate

ISBN : 9788119821327

relationships, dependencies, and overall impact of these factors on achieving sustainable land use. Chen et al., 2010 evaluated weight sensitivity of MCDM model for land suitability evaluation for irrigated agriculture. They aimed to study the sensitivity of converting weights of the input capabilities on the model output. The outcomes suggested a sturdy influence of sensitivity and, therefore, they've advocated giving unique emphasis in this criterion. Zolekar and Bhagat, 2015, have used GIS-primarily based MCDM version with IRS P6 LISS-IV pix as enter for the evaluation of agricultural practices in hilly areas. The rank of influential criteria changed into determined by way of correlation analysis and tips from scientific literature. The mixed use of faraway sensing and GIS turned out to be beneficial for land suitability evaluation. The integrated application of fuzzy, AHP, and GIS helped to bypass the problems on account of the uncertainties, subjectivities, and hierarchy traits of the conventional land suitability evaluation technique. GIS is a powerful tool to delineate the take a look at place, manage geographic information, manner maps, and gift results in land suitability evaluation. Integration of Fuzzy set and AHP methods with GIS presents a unique and effective combination in applying for land suitability analysis.

Soil health and fertility management :-

Soil productivity is directly linked to soil fertility, as it determines the availability of nutrients and water for crops. Unfortunately, soil fertility has been declining due to various factors such as pollution, sealing, overgrazing, waterlogging, excessive use of agricultural chemicals, and erosion. To effectively manage soil health and fertility, it is crucial to assess its status and plan site-specific management practices or precision farming techniques (Velayutham & Vhattacharya, 2000).

ISBN : 9788119821327

Assessing soil fertility involves analyzing macronutrients (N, P, and K), micronutrients (Zn, Mn, and Fe), pH levels, soil organic carbon (SOC), water holding capacity, erosion status, and moisture content (Shen, et al., 2016, Yang, et al., 2019). Geospatial analysis techniques like spatial interpolation, Multi-Criteria Decision Analysis (MCDA), and Ordered Weighted Averaging (OWA) are commonly used to assess the spatiotemporal variability of soil health and fertility, providing valuable information for decision makers (Malczewski, et al., 2003, Malczewski, et al., 2005).

Evaluating soil erosion status is a vital aspect of assessing soil quality and understanding its spatial variation is crucial for effective agricultural planning (Kusre, et al., 2018). Fuzzy mathematics, combined with indicators such as soil organic matter (SOC), total N, total P, total K, available N, available P, available K, pH value, and cation exchange capacity, can be used in ArcGIS to determine soil fertility. This analysis revealed that mid- and low-yielding fields had low soil fertility, which was directly related to the soil profile configuration (Li & Zhang, 2011). The correlation between crop productivity and soil fertility is evident, and GIS-based soil maps and fertility status provide valuable information on crop suitability for specific fields. Leena et al., 2016, proposed a soil fertility management decision support system that utilizes GIS-enabled cloud technology. This system utilizes soil tests and crop responses to make fertilizer recommendations, helping farmers optimize their fertilizer usage and maximize yields. The spatial nutrient variation generated by this system serves as an excellent e-governance tool for government agencies.

ISBN : 9788119821327

Crop monitoring and yield prediction :-

Crop growth monitoring, accurate prediction of yield, and assessment of food production are essential for estimating economic return and managing food security. However, traditional methods of crop yield estimation have been found to be inadequate and can result in poor assessment and inaccurate crop area appraisal. Additionally, these methods are time-consuming, labor-intensive, and expensive due to the need for extensive crop and yield data collection (Reynolds, et al., 2000). Fortunately, technologies such as remote sensing (RS), GPS, and GIS offer significant advantages in this regard. They enable the assessment of temporal and spatial variability of crop dynamics and yield output. By combining RS and GIS with input from other technologies, an efficient solution can be achieved for monitoring crop health and developing models to predict crop yields across different spatial scales.

Various vegetation indices are available to evaluate the extent and vigor of vegetation, crop growth dynamics, stress caused by biotic or abiotic factors, and other useful assessments. Adhav et al., 2021 utilized multiple vegetation indices, including the Normalized Difference Vegetation Index (NDVI), Green Normalized Difference Vegetation Index (GNDVI), Chlorophyll Vegetation Index (CVI), and Difference Vegetation Index (DVI), to determine crop health and identify variations in health conditions.

Precision farming :-

Precision Farming, also known as Precision Agriculture (PA) or site-specific crop management (SSCM), refers to the utilization of technologies and principles to effectively manage the spatial and temporal variability associated with all aspects of agricultural

ISBN : 9788119821327

production. Earl et al., 1996 provided a definition of precision farming as a system that integrates information with crop production, aiming to enhance long-term, site-specific, and overall farm production efficiency, productivity, and profitability, while minimizing unintended impacts on wildlife and the environment. The primary objectives of precision farming include the improved management of inputs such as seeds, fertilizers, pesticides, herbicides, and water, ensuring that the right amounts of these inputs are applied at the right place and time. To achieve these goals, various essential tools and systems, such as GPS, GIS, and RS, are necessary for the timely collection of geospatial information on soil-plant-animal requirements. This information is then utilized to gain valuable insights, which are subsequently employed to prescribe and apply site-specific treatments, ultimately leading to enhanced agricultural productivity, sustainability, and environmental protection (Balafoutis, et al., 2017). Precision farming can be broadly categorized into three stages based on the data collection or site-directed activities that occur before, during, and after the crop growth period. These stages include the Preparatory or Pre-planting stage, Crop growth stage, and Harvesting stage.

Conclusions :-

The utilization of Geographic Information Systems (GIS) in the field of agriculture has experienced a significant surge in recent decades. This growth can be attributed to the advancements in digital technologies, which have enabled GIS to become an indispensable tool for assessing crops, soils, and their surrounding environments. As highlighted in this chapter, GIS is now being employed at every stage of the agricultural value chain. In addition to its well-established uses in

ISBN : 9788119821327

land suitability and use planning, as well as the management of water, soil, and various stresses affecting agricultural systems, GIS has also found new and emerging applications in the realm of digital agricultural tools and technologies. These applications include high-resolution crop monitoring, yield prediction, precision farming, and supply chain management for both primary produce and biomass utilization in energy production.

The multitude of capabilities and insights provided by GIS, particularly its ability to collect and analyze real-time data, has further enhanced its significance in delivering location-based intelligence. This spatial intelligence is crucial for improving the productivity and profitability of farms through the implementation of precision practices. With the combination of current and emerging applications, alongside existing and newer partner technologies, GIS plays a pivotal role in achieving sustainable agricultural productivity.

References :-

1. Adhav V, Das S, Patkar V. (2021). Crop health monitoring using geospatial technologies for nashik district, Maharashtra. International Journal of Modern Agriculture, 10 (2): 1395-1409.
2. Bajaj, S., Mishra , S., Chouhan , D., Bora , S. S., Parashar , S., Bargi, U. S., & Billore , V. (2023). A Comprehensive Review of Application of RS, GIS and GPS in Agriculture India. International Journal of Environment and Climate Change, 13(11), 2597–2608.
3. Balafoutis A, Beck B, Fountas S, Vangeyte J, Wal TV, Soto I (2017). Precision agriculture technologies positively contributing to GHG emissions mitigation, farm productivity and economics. Sustainability, 9 (8):1339.

ISBN : 9788119821327

4. Burrough PA, McDonnell RA. (1998) Principles of Geographical Information Systems. Oxford: Oxford University Press.
5. Chen Y, Yu J, Khan S. (2010). Spatial sensitivity analysis of multi-criteria weights in GIS-based land suitability evaluation. Environmental Modelling and Software, 25(12):1582-1591.
6. Consoli S, Vanella D. (2014). Mapping crop evapotranspiration by integrating vegetation indices into a soil water balance model. Agricultural Water Management, 143:71-81.
7. Delgado JA, Vandenberg B, Neer D, D'Adamo R. (2019). Emerging nutrient management databases and networks of networks will have broad applicability in future machine learning and artificial intelligence applications in soil and water conservation. Journal of Soil and Water Conservation, 74 (6):113A-118A.
8. Delgado JA, Short NM Jr, Roberts DP, Vandenberg B. (2019). Big data analysis for sustainable agriculture on a geospatial cloud framework. Frontiers in Sustainable Food Systems, 3:54.
9. Earl R, Wheeler PN, Blackmore BS, Godwin RJ. (1996). Precision farming: The management of variability. The Journal of the Institution of Agricultural Engineers, 51:18-23.
10. Elhag M. (2016). Evaluation of different soil salinity mapping using remote sensing techniques in arid ecosystems, Saudi Arabia. Journal of Sensors, 1:2016.
11. FAO. (2017). The Future of Food and Agriculture—Trends and Challenges. Rome: FAO.
12. Fortes PS, Platonov AE, Pereira LS. (2005). GISAREG—A GIS based irrigation scheduling simulation model to support improved water use. Agricultural Water Management, 77 (1–3):159-179.

ISBN : 9788119821327

13. Gebeyehu MN. (2019). Remote sensing and GIS application in agriculture and natural resource management. International Journal of Environmental Sciences & Natural Resources, 19 (2):45-49.
14. Gomiero T. (2016). Soil degradation, land scarcity and food security: Reviewing a complex challenge. Sustainability, 8 (3):281.
15. Kumar S, Karaliya SK, Chaudhary S. (2017). Precision farming technologies towards enhancing productivity and sustainability of rice-wheat cropping system. International Journal of Current Microbiology and Applied Sciences, 6 (3):142-151.
16. Kusre BC, Ghosh P, Nath K. (2018). Prioritization of soil conservation measures using erodibility indices as criteria in Sikkim (India). Journal of Earth System Science, 127 (6):1-3.
17. Li M, Zhang XL. (2011). GIS-based evaluation of farmland soil fertility and its relationships with soil profile configuration pattern. Ying Yong Sheng tai xue bao= The. Journal of Applied Ecology, 22 (1):129-136.
18. Leena HU, Premasudha BG, Basavaraja PK. (2016). Sensible approach for soil fertility management using GIS cloud. In: 2016 International Conference on Advances in Computing, Communications and Informatics (ICACCI). IEEE; Sep 21. pp. 2776-2781.
19. Malczewski J, Chapman T, Flegel C, Walters D, Shrubsole D, Healy MA. (2003). GIS multicriteria evaluation with ordered weighted averaging (OWA): Case study of developing watershed management strategies. Environment & Planning A, 35(10):1769-1784.

ISBN : 9788119821327

20. Malczewski J, Rinner C. (2005). Exploring multicriteria decision strategies in GIS with linguistic quantifiers: A case study of residential quality evaluation. Journal of Geography System, 7(2):249-268.
21. Reynolds CA, Yitayew M, Slack DC, Hutchinson CF, Huete A, Petersen MS. (2000) Estimating crop yields and production by integrating the FAO Crop Specific Water Balance model with realtime satellite data and ground-based ancillary data. International Journal of Remote Sensing, 21(18):3487-3508.
22. Shen W, Ni Y, Gao N, Bian B, Zheng S, Lin X (2016). Bacterial community composition is shaped by soil secondary salinization and acidification brought on by high nitrogen fertilization rates. Applied Soil Ecology, 108:76-83.
23. Sikdar PK, Chakraborty S, Adhya E, Paul PK. (2004). Land use/land coverchanges and groundwater potential zoning in and around Raniganj coal miningarea, Bardhaman District, West Bengal: A GIS and remote sensing approach. Journal of Spatial Hydrology, 4(2):1-24.
24. Sishodia RP, Ray RL, Singh SK. (2020). Applications of remote sensing in precision agriculture: A review. Remote Sensing, 12(19):3136.
25. Soomro TR. (2015). GIS enabling smart agriculture. Smart Agriculture: An Approach towards Better Agriculture Management, 8:1-6.
26. Tripathi MP, Panda RK, Raghuwanshi NS. (2003). Identification and prioritisation of critical sub-watersheds for soil conservation management using the SWAT model. Biosystems Engineering, Jul 1;85(3):365-379.

ISBN : 9788119821327

27. Velayutham M, Bhattacharyya T. (2000). Soil resource management. In natural resource management for agricultural production in India'. Special publication during International Conference on Management of Natural Resources for Sustainable Agricultural Production in the 21st Century, Feb 14. pp. 14-18.
28. Weiss M, Jacob F, Duveiller G. (2020). Remote sensing for agricultural applications: A meta-review. Remote Sensing of Environment, 263: 111402.
29. Yang MH, Mouazen AM, Zhao XM, Guo X. (2019). Assessment of a soil fertility index using visible and near-infrared spectroscopy in the rice paddy region of southern China. European Journal of Soil Science, Vol. 71, issue 4, pp. 615-626.
30. Zolekar RB, Bhagat VS. (2015). Multicriteria land suitability analysis for agriculture in hilly zone: Remote sensing and GIS approach. Computers and Electronics in Agriculture, 118: 300-321.

ISBN : 9788119821327

CHAPTER – 3

Principles of Geographic Information Systems

Deepak Chouhan[1] , Aniket Rajput[2]

[1] Scientist (Agricultural Engineering), Krishi Vigyan Kendra, Shahdol (M. P.)

[2] M. Tech (SWE), College of Agricultural Engineering (M. P.)

Abstract

Geographic information systems (GIS) are computer-assisted systems that facilitate the capture, storage, retrieval, analysis, and display of both spatial and nonspatial attribute data. These systems can acquire data from various sources, including survey data, geographical/topographical/aerial maps, satellite data, and archived data. The data can be in the form of locational data, such as latitudes and longitudes, or tabular (attribute) data. The integration of modern technologies has significantly increased the use of computers and information technology in all aspects of spatial data handling. This includes data acquisition, storage and maintenance, processing, analysis, and output. GIS is essentially a computerized database management system that captures, stores, validates, maintains, analyzes, displays, and manages spatially referenced data. Its primary function is to integrate data from diverse sources. A complete GIS comprises hardware, software, data, people, and methods. Advanced GIS capabilities enable the addressing of questions related to location, condition, trends, patterns, and modeling. Various sources, such as manual digitization of paper maps, existing digital data, aerial photographs, remote-sensing satellite imagery, and global positioning systems (GPS), offer promising options for data input in GIS. However, before generating any spatial data, it is crucial to understand the available data types, data analysis procedures, and the capabilities of GIS to ensure realistic outputs.

ISBN : 9788119821327

GIS is widely used as a spatial analysis tool for effective and efficient data generation, management, analysis, and display. Recent advancements in computer technology, remote sensing, GPS, and communication technology have further enhanced the functions of GIS in these areas. Data derived from remote sensing, GPS, and field surveys can be effectively utilized in GIS for mapping, monitoring, and managing land resources, as well as disseminating information in various location-based applications.

Keywords :- Geographic information systems, Hardware, Software

Introduction :-

Geographic Information System (GIS) is a computer-based system utilized for various tasks such as inputting, outputting, storing, manipulating, retrieving, and analyzing spatial data. These systems consist of both computer hardware and software components. GIS is increasingly being employed in a wide range of applications including natural resources management, tourism, transportation, trade, and commerce. Moreover, GIS is integrated with modern technologies such as remote sensing and GPS. The development of GIS began in the late 1950s, but it wasn't until the late 1970s that the first GIS software was introduced by the ESRI lab. Canada played a pioneering role in the advancement of GIS due to innovative efforts dating back to the early 1960s. One of the key advantages of using GIS is the ability to overlay a large number of maps. Traditionally, manual overlays were prepared, which was a cumbersome and error-prone process. In contrast, GIS enables faster retrieval of information through computer-based methods. Instead of browsing through physical maps, information can be retrieved more easily and automatically within a GIS.

Furthermore, GIS allows for the creation of interactive or virtual output. This virtual output is automatically updated whenever the component maps are revised. In contrast, the conventional method involves preparing hardcopy outputs, which are difficult to update. In

ISBN : 9788119821327

such cases, the entire map needs to be redrawn, and annotation becomes cumbersome. Consequently, ambiguity may arise during information retrieval, and certain features may not be properly annotated in paper maps. In contrast, GIS stores information in tables and links it to geographic features, eliminating limitations related to annotation space, color, symbol, and other physical constraints. This ensures that attribute information, such as names, for various features is not lost in paper products. Moreover, in GIS, common boundaries can be digitized once and made available to all layers. This eliminates the need for manual drawing of common boundaries in different maps, which may not match accurately. Once GIS map layers are prepared, any number of maps can be designed, providing flexibility and efficiency. Overall, the use of GIS offers significant advantages over traditional paper maps, as it simplifies information retrieval, enables automatic updates, and provides greater flexibility in map design.

Data capture or input in Geographic Information Systems (GIS) can be a costly endeavor. In comparison, commercially available paper maps may prove to be a more economical option than GIS layers. The primary reason for this cost difference lies in the high expenses associated with data capture in GIS. Utilizing GIS technology necessitates investments in computers, software, and training. Moreover, the handling of GIS requires skilled personnel who are proficient in its operation. In GIS, data must be converted into the native format of the GIS software, which can further add to the overall expenses. Additionally, if suitable import/export functionality is lacking or if the data format is obscure or unknown, the data may become unusable. Therefore, it is crucial for GIS software to possess the necessary functionality required for specific applications. For instance, in transportation applications, the availability of network analysis functions is essential.

ISBN : 9788119821327

The initial definition of GIS regarded it as a spatial database; however, the current inclination is to perceive it in various manners (ESRI, 1996).

• An information technology that stores, analyzes, and presents both spatial and non-spatial data.

• A decision support system that incorporates spatially referenced data within a problem-solving context.

• A tool utilized for the management, analysis, and manipulation of spatially referenced data information within a problem-solving environment.

Components of GIS :- GIS is an acronym that encompasses three interconnected components: geography, information, and systems. Firstly, geography represents the physical world and its spatial characteristics. Secondly, information refers to data and its interpretation, encompassing its significance and utilization. Lastly, systems denote computer technology and the supporting infrastructure. By combining these elements, GIS enables the collection, compilation, storage, retrieval, analysis, manipulation, and integration of environmental, economic, and social data within a unified framework. This is visually depicted in Figure 1 and Figure 2.




Figure 1 :- Integrated elements of GIS

ISBN : 9788119821327




Figure 2 :- The component parts of GIS.

Computer Hardware :- In figure 3 illustrates the fundamental hardware components of a GIS system. The computer is equipped with a hard disk to store both data and programs. However, additional storage options are available through network connections or the utilization of digital tape cassettes, optical CD-ROMs, and other similar devices. To enable the computer programs to utilize maps and documents, a scanner is employed to convert them into a digital format. Finally, a plotter or printer is utilized to visually present the outcomes of the data processing.




Figure 3 :- Major hardware components of Geographical Information System

ISBN : 9788119821327

GIS software :- The rang in GIS software systems is as variable as the hardware it runs, but their functionality is centred on four common types of actions (Figure 4)

- Data input and verification
- Data storage and database management
- Data output and presentation
- Data transformations, analysis, and modelling




Figure 4:- Main software components of GIS.

Data Input :- The data input (Figure 5) encompasses a wide range of possibilities, including the utilization of digital data sets, field observations, or output from various digital sensors. Additionally, it involves the conversion of paper maps or images into digital data sets. This serves as the fundamental framework for constructing a geographic database and forms the basis for all subsequent analysis and output procedures.

Data storage and databases :- The organization and management of data storage and databases (Figure 6) pertain to the manner in which information regarding geographical elements, including their location, connections, and attributes, is structured and arranged. This involves considering how the data should be processed within a computer system and how it is interpreted by system users. The software utilized

ISBN : 9788119821327

to organize and manage the database is commonly referred to as a Data Base Management System (DBMS).




Figure 5 :- Data collection and input




Figure 6:- Components of geographic data based

Data transformation :- Data transformation encompasses two categories of operations: (a) the necessary transformations to rectify errors in the data, update them, or align them with other datasets, and (b) the extensive range of analysis methods that can be applied to the data to obtain answers to the inquiries posed by the GIS. These transformations can be performed on the spatial, topological, and non-

ISBN : 9788119821327

spatial aspects of the data, either independently or in conjunction. Numerous transformations, such as those related to changing scales, adjusting data to new projections, logically retrieving data, and calculating areas and perimeters, have reached a level of maturity that makes them commonplace in virtually every type of GIS. The available transformation methods, their optimal utilization and potential misuse, as well as the ways in which a set of simple transformations can be combined to achieve various forms of geographical or spatial modeling, are all factors to consider (figure 7).




Figure 7:- Data Transformation

Date Output and presentation :- The information can be displayed in various formats, such as maps, tables, and figures (graphs and charts), which can be presented in different ways. These include temporary images on computer screens, printed outputs on printers or plotters, and digitally recorded data on magnetic media (figure 8).




Figure 8:- Data Output

ISBN : 9788119821327

Advantages of GIS :- GIS applications facilitate the organized gathering, updating, processing, and dissemination of land-related data. Information pertaining to geographic location, such as zoning, roads, and parks, can all be effectively managed through GIS. A key requirement of GIS is its ability to handle large volumes of land survey data. Not only can GIS analyze and manipulate spatial information, but it can also seamlessly integrate diverse data types at a rapid pace. Moreover, GIS serves as a valuable tool for monitoring and managing various forms of information. The overlaying of different thematic maps is effortlessly accomplished using the capabilities of GIS. Thematic maps can be created and kept up-to-date by utilizing high-resolution spatial and temporal remote sensing data. The advantages of GIS become particularly evident when transitioning from manual analysis, especially when dealing with extensive datasets. Additionally, GIS has proven to be highly beneficial for complex modeling or when combining datasets with different weightings. Figure 9 visually illustrates the management capabilities of GIS.




Figure 9:- GIS as a management tool

ISBN : 9788119821327

References :-

1. Chrisman, N. R., (2003). Exploring Geographical Information Systems. Wiley, Hoboken, NJ.
2. Heywood, I., Cornelius, S., and Carver, S. (20111). An Introduction to Geographical Information Systems. 4th edn. Pearson, Harlow.
3. Lillesand, T. M., Kiefer, R. W., and Chipman, J. W. (2008). Remote Sensing and Image Interpretation. 6th edn. Wiley, Hoboken, NJ.
4. Lin, H. and Batty, M. (eds.) (2009). Virtual Geographic Environments. Science press, Beijing.
5. Longley, P. A., Goodchild, M. F., Maguire, D. J., and Rhing, D. W. (2010). Geographic Information Systems and Science. 3rd edn. Wiley, Hoboken, NJ.
6. Reddy GPO (2012) Principles and concepts of GIS. In: Reddy GPO, Sarkar D (eds) Remote sensing and GIS in digital terrain analysis and soil-landscape modeling, pp 69–77
7. Star J, Estes J (1990) Geographical information systems: an introduction. Prentice Hall, Englewoods Cliffs

ISBN : 9788119821327

CHAPTER – 4

"Environmental Geoinformatics: Monitoring and Management"

Dr. K. M. Gojiya1*, Er. B. A. Karangiya2 and Prof. M. J. Gojiya3

1Research Associate, Agriculture Research Station (Fruit Crops), Junagadh Agricultural University, Mahuva, Gujarat, India-364290

2Senior Research Assistant, College of Agril. Engg. & Tech., Junagadh Agricultural University, Junagadh, Gujarat, India-362001

3Assistant Professor, College of Agril. Engg. & Tech., Junagadh Agricultural University, Junagadh, Gujarat, India-362001

Corresponding Author Email: kashyapgojiya@gmail.com

Phone Number: 8306678888

Abstract

This chapter explores the pivotal role of geoinformatics in the realm of "Environmental Geoinformatics: Monitoring and Management." Beginning with an introduction to the fundamentals, including Geographic Information Systems (GIS), Remote Sensing, and the Global Positioning System (GPS), the narrative unfolds to elucidate diverse data acquisition methods. From satellite imagery and ground-based monitoring to the challenges and benefits of data integration, the chapter provides a comprehensive understanding of environmental geoinformatics. The applications in environmental monitoring span biodiversity conservation, climate change assessments, and air and water quality analysis. The subsequent section delves into geoinformatics' contribution to environmental management, examining Decision Support Systems, disaster management, and sustainable resource management. As the chapter concludes, it addresses the challenges, future trends, and emphasizes the continued significance of geoinformatics in environmental research and implementation.

Key words*: Geoinformatics, Environmental Monitoring, Decision Support Systems, Sustainable Resource Management*

ISBN : 9788119821327

Introduction

In the intricate dance of the Earth's natural systems, understanding and mitigating the impact of human activities on the environment have become imperatives for sustainable living. Environmental geoinformatics, at the intersection of geography, information science, and environmental sciences, emerges as a powerful ally in this pursuit. This chapter delves into the realm of "Environmental Geoinformatics: Monitoring and Management," exploring the dynamic synergy between advanced geospatial technologies and environmental conservation.

The accelerated pace of global environmental change, driven by factors such as climate change, biodiversity loss, and pollution, demands innovative and comprehensive approaches to monitoring and management. Geoinformatics, encompassing Geographic Information Systems (GIS), remote sensing, and Global Positioning System (GPS), equips researchers, policymakers, and conservationists with a sophisticated toolkit to analyze, visualize, and interpret complex spatial data.

As we embark on this exploration, it is crucial to recognize the pivotal role of geoinformatics technologies in facilitating a deeper understanding of environmental dynamics. Whether unravelling the mysteries of biodiversity hotspots, tracking the subtle shifts in climate patterns, or assessing the quality of air and water, geoinformatics provides a lens through which the intricate tapestry of the Earth's systems can be observed and comprehended.

This chapter will navigate through the fundamentals of geoinformatics, elucidate the diverse methods of data acquisition, and showcase the myriad applications in environmental monitoring. From the microcosm of species distribution modelling to the macrocosm of climate change assessments, geoinformatics emerges as an indispensable tool for informed decision-making and sustainable environmental management.

ISBN : 9788119821327

As we embark on this intellectual journey, let us delve into the innovative realm of environmental geoinformatics, where data transforms into knowledge, and knowledge into actionable insights for the stewardship of our planet.

Fundamentals of Geoinformatics

Geoinformatics, a multidisciplinary field at the intersection of geography and information technology, plays a pivotal role in environmental monitoring and management. This section delves into the fundamental components of geoinformatics, exploring the core technologies of Geographic Information Systems (GIS), Remote Sensing, and the Global Positioning System (GPS).

A. Geographic Information Systems (GIS)

1. **Definition and Core Components:** Geographic Information Systems (GIS) represent a powerful framework for capturing, analyzing, and visualizing spatial and geographical data. At its core, GIS integrates data layers, enabling the creation of comprehensive maps and facilitating spatial analysis. The components include spatial data, software, hardware, and human resources. Spatial data, often in the form of layers, can include information about terrain, land use, and various environmental parameters. GIS software processes and analyzes this data, while hardware includes computers and devices used for data input and output.
2. **Applications in Environmental Monitoring and Management:** GIS applications in environmental contexts are diverse and impactful. From mapping biodiversity hotspots to analyzing land cover changes, GIS aids in decision-making processes related to environmental conservation. It serves as a tool for spatial modeling, risk assessment, and resource management. Environmental monitoring benefits from GIS by providing a spatial context to data, facilitating the identification of patterns and trends critical for informed decision-making.

ISBN : 9788119821327

B. Remote Sensing

1. **Introduction to Remote Sensing Technologies:** Remote sensing involves the acquisition of information about the Earth's surface without direct physical contact. Technologies include satellite and aerial sensors that capture electromagnetic radiation across various wavelengths. These sensors collect data on vegetation health, land cover, and atmospheric conditions. This non-invasive approach allows for large-scale data collection, providing valuable insights into environmental dynamics.
2. **Role in Environmental Data Collection:** Remote sensing plays a pivotal role in environmental data collection by offering a synoptic view of the Earth's surface. It provides real-time and historical data, enabling the monitoring of changes over time. From assessing deforestation patterns to tracking urban expansion, remote sensing contributes to comprehensive environmental assessments. The data collected supports decision-makers in understanding environmental changes and implementing effective conservation strategies.

C. Global Positioning System (GPS)

1. **Overview and Functionality:** The Global Positioning System (GPS) is a satellite-based navigation system that allows users to determine their precise location and track movement. Comprising a network of satellites orbiting the Earth, GPS receivers triangulate signals to provide accurate positional information. This technology has become ubiquitous in various applications, from navigation systems to location-based services on smartphones.
2. **Integration with Environmental Monitoring:** GPS integration enhances the precision and accuracy of environmental monitoring activities. Researchers and field practitioners utilize GPS to georeference data, ensuring spatial context and accuracy.

ISBN : 9788119821327

This integration is crucial in mapping ecosystems, tracking wildlife movements, and precisely locating monitoring stations. The synergy between GPS and environmental monitoring amplifies the effectiveness of data collection and contributes to a more nuanced understanding of ecological processes.

Understanding the fundamentals of GIS, remote sensing, and GPS establishes the groundwork for harnessing geoinformatics in environmental monitoring and management. These technologies, when integrated, provide a holistic approach to studying and safeguarding our planet's ecosystems.

Data Acquisition in Environmental Geoinformatics

Environmental geoinformatics relies on diverse data acquisition methods to capture a comprehensive view of the Earth's systems. This section explores key approaches, including satellite imagery, ground-based monitoring, and the integration of data from multiple sources.

A. Satellite Imagery

1. **Types and Resolution:** Satellite imagery encompasses a range of types, from optical and infrared to synthetic aperture radar (SAR). These satellites vary in their spatial, spectral, and temporal resolutions. High-resolution optical satellites capture detailed images, while SAR satellites penetrate clouds and provide information about surface characteristics. The selection of satellite type depends on the specific requirements of environmental studies.
2. **Interpretation for Environmental Studies:** Satellite imagery interpretation is a crucial skill in environmental geoinformatics. Analysts use these images to monitor land cover changes, assess deforestation, and identify environmental hazards. The data extracted from satellite imagery contribute to environmental modeling, allowing researchers to understand ecosystem dynamics and human impacts on the landscape.

ISBN : 9788119821327

B. Ground-based Monitoring

1. **Sensor Networks:** Ground-based monitoring involves deploying sensor networks across ecosystems to gather real-time data. These sensors can measure parameters such as temperature, humidity, and air quality. Sensor networks offer a fine-grained view of environmental conditions, enabling continuous monitoring and early detection of changes.
2. **In-situ Data Collection Methods:** In-situ data collection involves on-site measurements and observations. Researchers employ field instruments to collect data on soil composition, vegetation health, and water quality. Ground-based methods complement satellite data, providing detailed, localized information essential for accurate environmental assessments.

C. Data Integration and Fusion

1. **Challenges and Strategies:** Integrating data from various sources presents challenges, including differences in spatial and temporal resolutions, data formats, and accuracy. Strategies for overcoming these challenges include standardization of data formats, calibration of sensors, and the development of interoperable data platforms. Overcoming these challenges is essential for creating a cohesive and accurate understanding of environmental processes.
2. **Benefits of Integrating Multiple Data Sources:** The integration of data from diverse sources enhances the richness and accuracy of environmental information. Combining satellite imagery with ground-based measurements provides a more complete picture of ecosystems. This synergy enables researchers and decision-makers to assess environmental changes with greater confidence, leading to more effective conservation and management strategies.

ISBN : 9788119821327

Applications in Environmental Monitoring

Environmental geoinformatics applications span a wide range of fields, contributing to biodiversity conservation, climate change monitoring, and assessments of air and water quality.

A. Biodiversity Conservation

1. **Habitat Mapping:** GIS and remote sensing are instrumental in mapping habitats critical for biodiversity. High-resolution satellite imagery aids in identifying and delineating ecosystems, helping conservationists prioritize areas for protection and restoration.
2. **Species Distribution Modeling:** Environmental geoinformatics facilitates the modeling of species distributions based on environmental variables. This modeling assists in predicting species habitats, supporting conservation efforts, and mitigating the impact of human activities on vulnerable ecosystems.

B. Climate Change Monitoring

1. **Temperature and Precipitation Analysis:** Geoinformatics technologies track temperature and precipitation patterns, providing essential data for climate change studies. GIS and remote sensing contribute to the analysis of temperature trends, precipitation anomalies, and the identification of climate change hotspots.
2. **Sea Level Rise and Glacial Retreat:** Satellite observations play a key role in monitoring sea level rise and glacial retreat. These data contribute to understanding the impact of climate change on coastal regions and high-altitude environments, informing adaptation strategies.

C. Air and Water Quality Assessment

1. **Monitoring Pollution Sources:** Geoinformatics assists in monitoring and identifying sources of air and water pollution. Satellite imagery and ground-based sensor networks provide spatially explicit data that help trace pollutants back to their origins.

ISBN : 9788119821327

2. **Water Body Health Assessment:** The integration of geoinformatics data supports the assessment of water body health. Parameters such as water temperature, nutrient levels, and sedimentation are monitored, aiding in the identification of pollution sources and the development of strategies for water quality improvement.

The integration of data acquisition methods and the application of geoinformatics technologies in environmental monitoring underscore the interdisciplinary nature of contemporary environmental science. These tools empower researchers and policymakers to make informed decisions, contributing to the sustainable management of our planet's resources.

Geoinformatics in Environmental Management

Geoinformatics serves as a linchpin in effective environmental management, offering decision-makers powerful tools to navigate complex challenges. This section explores the pivotal roles of Geoinformatics in Decision Support Systems (DSS), disaster management, and sustainable resource management.

A. Decision Support Systems (DSS)

Role in Policy and Planning: Decision Support Systems (DSS) powered by geoinformatics play a critical role in policy formulation and planning for environmental management. These systems integrate spatial data, analytical tools, and modeling capabilities to provide decision-makers with comprehensive insights. In policy development, DSS aids in identifying optimal locations for conservation areas, predicting the impact of land-use changes, and assessing potential environmental risks. Geoinformatics-driven DSS enhances the effectiveness of policies by ensuring they are grounded in spatially explicit information.

B. Disaster Management

1. **Early Warning Systems:** Geoinformatics contributes significantly to the establishment of early warning systems for

ISBN : 9788119821327

natural disasters. By integrating real-time data from satellites, ground-based sensors, and weather models, these systems can provide timely alerts for events such as hurricanes, floods, or wildfires. The spatial analysis capabilities of geoinformatics enable precise delineation of at-risk areas, enhancing the effectiveness of evacuation and response efforts.

2. **Post-Disaster Assessment and Recovery:** Following a natural disaster, geoinformatics supports rapid post-disaster assessment and recovery. Satellite imagery and GIS data aid in mapping the extent of damage, identifying affected areas, and prioritizing rescue and recovery operations. Geospatial technologies contribute to the reconstruction phase by providing accurate information for rebuilding infrastructure and restoring ecosystems.

C. Sustainable Resource Management

1. **Land Use Planning:** Geoinformatics facilitates sustainable land use planning by offering insights into the spatial distribution of land cover, soil types, and terrain characteristics. GIS tools help optimize urban development, agriculture, and infrastructure projects, ensuring that land use is aligned with environmental conservation goals. Through scenario modeling, decision-makers can assess the potential impacts of various land-use scenarios on ecosystems and biodiversity.
2. **Natural Resource Monitoring and Conservation:** Geoinformatics is instrumental in monitoring and conserving natural resources. Satellite imagery aids in tracking deforestation, soil erosion, and changes in vegetation cover. GIS-based analyses contribute to the identification of critical habitats and corridors for biodiversity conservation. By integrating geospatial data, decision-makers can formulate sustainable resource management strategies that balance human needs with ecological preservation.

ISBN : 9788119821327

The integration of geoinformatics into environmental management practices represents a paradigm shift, offering a holistic and data-driven approach. Decision Support Systems, disaster management tools, and sustainable resource management strategies underpinned by geoinformatics empower societies to make informed choices that preserve the environment for current and future generations.

Challenges and Future Trends

Environmental geoinformatics, while transformative, is not without its challenges. Additionally, anticipating future trends is essential to staying ahead in this dynamic field.

A. Data Privacy and Security:

- ***Challenge:*** The abundance of geospatial data raises concerns about privacy and security. As more sensitive environmental data becomes available, safeguarding it from unauthorized access and ensuring ethical use become paramount.
- ***Future Trend:*** The future involves developing robust data encryption and anonymization techniques. Striking a balance between open access to environmental information and protecting individual privacy will be a focal point.

B. Integration of Emerging Technologies:

- ***Challenge:*** The rapid evolution of technology introduces challenges in seamlessly integrating emerging tools and methodologies into existing geoinformatics frameworks.
- ***Future Trend:*** Artificial Intelligence (AI), machine learning, and blockchain are poised to play pivotal roles in the future of environmental geoinformatics. Integrating these technologies will enhance data analysis, predictive modeling, and data transparency.

ISBN : 9788119821327

C. Policy and Institutional Challenges:

- ***Challenge:*** The development of effective policies and institutional frameworks to govern the ethical use, sharing, and accessibility of geoinformatics data poses a challenge.
- ***Future Trend:*** A growing need for international collaboration and standardization will drive the formation of policies that address global environmental challenges. Institutions will need to adapt to the evolving landscape of environmental data management.

D. Future Directions in Environmental Geoinformatics:

- ***Trend:*** Increased emphasis on real-time monitoring and predictive modeling will define the future of environmental geoinformatics. The integration of data from diverse sources, including citizen science and crowdsourced data, will become more prevalent.
- ***Trend:*** The development of more sophisticated sensor technologies and satellite constellations will enhance data collection capabilities. This will allow for more frequent and detailed observations of the Earth's surface.

Conclusion

Environmental geoinformatics encompasses GIS, remote sensing, and GPS technologies, providing a comprehensive toolkit for environmental monitoring and management. Data acquisition involves satellite imagery, ground-based monitoring, and data integration, offering a nuanced understanding of environmental processes. Geoinformatics applications include biodiversity conservation, climate change monitoring, and air and water quality assessment. In environmental management, geoinformatics contributes through Decision Support Systems, disaster management tools, and sustainable resource management strategies. Geoinformatics enables evidence-based decision-making, fostering sustainable environmental practices. The integration of geospatial technologies enhances the accuracy and

ISBN : 9788119821327

efficiency of environmental monitoring, contributing to effective conservation and resource management. Ongoing research is crucial for addressing challenges and embracing emerging technologies in environmental geoinformatics. Implementation of geoinformatics solutions in policy and practice is vital for translating research into impactful environmental management strategies.

In conclusion, environmental geoinformatics stands as a linchpin in the pursuit of sustainable environmental practices. Addressing challenges and embracing future trends will be essential to harness the full potential of geoinformatics for the benefit of our planet. A continued commitment to research and implementation is necessary to navigate the complexities of the dynamic environmental landscape.

ISBN : 9788119821327

CHAPTER – 5

Uses of GIS, GPS and VRA in Precision Agriculture

Tanveer Ahmad Ahngar[1]*, Ruby Jan[1] and Zahoor Ahmad Dar[2]

[1]Division of Agronomy, Faculty of Agriculture, Wadura, SKUAST-K, Jammu and Kashmir, India

[2]Dryland Agriculture Research Station, Rangreth, SKUAST-K, Jammu and Kashmir, India.

Corresponding author: tanveeragron@gmail.com

Abstract

This succinct abstract provides a glimpse into the transformative applications of Geographic Information System (GIS), Global Positioning System (GPS), and Variable Rate Application (VRA) in precision agriculture. Delve into the pivotal roles of these technologies in enhancing operational precision, spatial intelligence, and resource efficiency. From GPS-guided navigation and GIS-driven spatial analysis to the dynamic customization of inputs with VRA, this chapter explores the integrated framework shaping modern farming practices. Uncover how the seamless collaboration of GIS, GPS, and VRA is revolutionizing precision agriculture, paving the way for sustainable and informed decision-making in every field.

INTRODUCTION

In the dynamic world of modern agriculture, the convergence of technology and cultivation practices has given rise to a transformative approach – precision agriculture. At the forefront of this revolution are three technological pillars that have reshaped the farming landscape: Global Positioning System (GPS), Geographic Information System (GIS), and Variable Rate Application (VRA). Together, they constitute a powerful triad that not only navigates the vast expanses of agricultural

ISBN : 9788119821327

fields but also maps insightful strategies for optimized resource application and sustainable farming practices.

Precision farming is characterized as the utilization of technologies and principles to effectively handle spatial and temporal variability across all aspects of agricultural production (Pierce and Nowak, 1999). Over the past decade, numerous technological advancements have contributed to the evolution of precision farming. The adaptability of precision farming hinges on incorporating and leveraging modern technologies, including advanced farm technologies and site-specific systems. These technologies encompass high-speed internet connectivity and increased farmer awareness.

Precision farming serves as an integrated information and agricultural management system designed to enhance overall farm production efficiency. It emphasizes achieving this improvement in a cost-effective manner while mitigating the adverse impacts of chemical loading on the environment. The central focus of precision farming is the collection of information pertaining to soil and crop conditions, capturing the intricate sequence of these conditions at a spatial level.




Figure: Precision farming integration of technologies

ISBN : 9788119821327

GPS: Global Positioning System (Precision Navigation in Agriculture)

` It is a system for navigation based on satellites that enables the determination of positions anywhere on Earth (Aronoff, 1989). The fields that once relied on the seasoned eye of the farmer have now embraced the precision and efficiency offered by GPS technology. Global Positioning System, originally designed for satellite navigation, has found a crucial application in agriculture. From precise tractor guidance during planting and harvesting to optimizing routes for irrigation and crop monitoring, GPS has become the guiding star for farmers, reducing overlaps, minimizing waste, and maximizing operational efficiency. The utilization of technology has been significantly impacted by the introduction of the Global Positioning System (GPS), not only streamlining spatial data acquisition but also expanding its integration with remote sensing and geographic information systems (GIS). Emerging in the late 1980s, GPS transformed spatial data acquisition, evolving from traditional handheld devices with approximately 10 meters accuracy to highly precise differential GPS (DGPS) devices capable of accuracy within centimetres. However, challenges arise in hilly terrains like Himachal, where small field areas and existing cropping patterns pose limitations for practical implementation of Precision Farming (PF).

GPS technology allows for the recording of in-field variability as geographically encoded data, continuously determining and recording accurate positions. This level of detail enhances database richness for users, particularly in capturing accurate yield data at recorded GPS positions. By integrating GPS receivers with yield monitors, spatial coordinates are obtained for yield monitor data, facilitating the creation of yield maps for each field. The combination of satellite data and GPS-referenced information supports the development of field management strategies for chemical application, cultivation, and harvest.

ISBN : 9788119821327

The advent of precision agriculture, or site-specific farming, is attributable to the integration of the Global Positioning System (GPS) and geographic information systems (GIS). These technologies enable the concurrent collection of real-time data with precise position information, facilitating the efficient manipulation and analysis of extensive geospatial data. In the realm of precision farming, GPS applications are diverse, encompassing farm planning, field mapping, soil sampling, tractor guidance, crop scouting, variable rate applications, and yield mapping.

Geographic Information System (GIS):

It is a computerized system for storing and retrieving data that manages and analyzes spatial information related to crop productivity and agronomic factors. This system has the capability to integrate various types of information and interface with other decision support tools. GIS can present analyzed data in maps, offering a better understanding of interactions among plants, pathogens, and their environment, thereby aiding decision-making based on these spatial relationships. Many affordable GIS software options with diverse functions are currently available. Several farm information systems (FIS), such as LORIS (Local Resources Information System), use simple programs to establish a farm-level database. LORIS, for instance, includes modules for data import, generation of raster files using different gridding methods, storage of information in a database, creation of digital agro-resource maps, and development of operational maps (Schroder et al., 1997). A comprehensive farm GIS incorporates base maps detailing topography, soil types, nutrient levels (such as N, P, K), soil moisture, pH, etc. Additionally, it can store data on crop rotations, tillage practices, nutrient and pesticide applications, yields, and more. GIS proves beneficial in creating maps that illustrate fertility, weed and pest intensity, facilitating the development of maps indicating recommended application rates for nutrients or pesticides.

ISBN : 9788119821327

GIS's effectiveness lies in its ability to overlay various data layers, including ecological conditions, physiognomy, and human pressure indices. This layer-based, thematic system grants flexibility to overlay and assess indices, facilitating the review of site changes. GIS technology is maximized in planning and management applications.

VRA or Variable Rate Application

It is a key component of precision agriculture. It refers to the capability of adjusting the rate of inputs, such as fertilizers, pesticides, and irrigation, based on the specific needs of different areas within a field. Instead of applying a uniform rate across an entire field, VRA allows farmers to customize the application of inputs based on spatial variability, ultimately optimizing resource use and improving overall efficiency.

The implementation of VRA relies on the integration of technologies such as Global Positioning System (GPS) and Geographic Information System (GIS). These technologies provide the necessary spatial data and guidance to enable precise and variable application rates. By analyzing data on soil types, nutrient levels, moisture content, and other relevant factors, farmers can make informed decisions about the amount and placement of inputs required in different areas of their fields.

The advantages of VRA in precision agriculture include increased efficiency, cost savings, and environmental sustainability. Farmers can target specific areas that need more or less of a particular input, reducing waste and minimizing the environmental impact associated with excessive use of fertilizers or pesticides. VRA is a powerful tool that contributes to the overall goal of precision agriculture, which is to optimize resource use, increase productivity, and enhance sustainability in farming practices.

Integration of GIS, GPS, and VRA

The integration of GIS (Geographic Information System), GPS (Global Positioning System), and VRA (Variable Rate Application) in precision

ISBN : 9788119821327

agriculture forms a powerful technological framework that revolutionizes farming practices. Here's how these components synergize:

1. **GIS for Spatial Intelligence:**
    - GIS collects and analyzes spatial data related to the agricultural landscape. This includes information on topography, soil types, crop health, and environmental factors.
    - GIS provides a visual representation of this data through maps, allowing farmers to better understand the variability within their fields.
2. **GPS for Precision Navigation:**
    - GPS technology guides farming machinery with unparalleled precision. Tractors, planters, and harvesters equipped with GPS systems can follow predefined paths, reducing overlaps and optimizing field coverage.
    - GPS enables the accurate geo-referencing of data collected in the field, ensuring precise mapping and analysis.
3. **VRA for Dynamic Customization:**
    - VRA builds on the insights provided by GIS and GPS. It allows farmers to customize the application rates of inputs such as fertilizers, pesticides, and water based on the specific needs of different zones within a field.
    - By integrating GPS data for accurate positioning and GIS data for spatial insights, VRA systems dynamically adjust input application rates in real-time.
4. **Spatial Data Fusion:**
    - GIS serves as the hub for integrating spatial data from various sources. This includes GPS-derived location data, satellite imagery, and other relevant information.
    - The fusion of these data sets within GIS enables a comprehensive understanding of the spatial variability in the field, forming the basis for informed decision-making.

ISBN : 9788119821327

5. **Prescription Mapping:**
    - GIS, GPS, and VRA collaborate in creating prescription maps. These maps guide farming equipment to apply inputs at variable rates across the field according to the specific conditions identified by GIS analysis.
    - Prescription maps are dynamic, allowing farmers to adapt strategies for planting, fertilization, and pest control based on real-time data.
6. **Efficiency and Sustainability:**
    - The integration of these technologies improves overall efficiency in farming operations. Precision navigation reduces overlaps, minimizing input waste.
    - Customized variable rate applications ensure that resources are used optimally, promoting sustainability by reducing the environmental impact of excess inputs.
7. **Data-Driven Decision Making:**
    - The integration of GIS, GPS, and VRA facilitates data-driven decision-making. Farmers can access real-time information about their fields, enabling them to make informed choices for crop management.

In essence, the seamless integration of GIS, GPS, and VRA creates a precision agriculture system where spatial intelligence, precise navigation, and dynamic customization converge to enhance productivity, reduce costs, and promote sustainable farming practices. This integration empowers farmers with the tools needed to navigate the complexities of modern agriculture while minimizing environmental impact.

ISBN : 9788119821327

Major challenges associated with GIS, GPS, and VRA in precision agriculture:

1. Data Integration and Compatibility:

Diverse data formats and lack of standardized interoperability hinder seamless integration of GIS, GPS, and VRA technologies. Difficulty in exchanging information between systems, leading to inefficiencies and incomplete datasets for decision-making.

2. Complexity and Technical Expertise:

Farmers may struggle with the complexity of integrated systems, requiring a level of technical expertise that might be beyond their current skill set. Limited adoption and underutilization of technology due to a lack of understanding and confidence in managing integrated solutions.

3. Costs and Affordability:

The initial investment and ongoing costs associated with GIS, GPS, and VRA technologies can be prohibitive for some farmers. Limited access to precision agriculture benefits, hindering adoption and potential productivity gains.

4. Data Security and Privacy Concerns:

The collection and storage of sensitive spatial data raise concerns about security and privacy. Farmer hesitancy to adopt technologies due to concerns about data misuse or unauthorized access.

5. Limited Connectivity in Remote Areas:

Poor internet connectivity in remote agricultural regions impedes real-time data exchange. Hindered access to real-time information and reduced effectiveness of precision agriculture technologies in remote areas.

6. Resistance to Change:

Psychological barriers and resistance to adopting new technologies among farmers.Slow adoption rates, particularly in regions where traditional farming practices are deeply ingrained.

ISBN : 9788119821327

7. Regulatory and Policy Challenges:

Lack of clear regulations and policies regarding the use of spatial technologies in agriculture.

8. Scalability and Customization:

Challenges in scaling integrated systems for various farm sizes and customizing solutions to suit diverse agricultural operations.

9. Data Accuracy and Quality:

Obtaining accurate and high-quality spatial data is crucial for effective precision agriculture as inaccurate data may lead to flawed decision-making, reducing the overall effectiveness of precision agriculture practices.

10. Collaboration and Communication*:*

Ineffective collaboration and communication among stakeholders, including farmers, technology providers, and policymakers.

Understanding and addressing these challenges is crucial for the successful integration and widespread adoption of GIS, GPS, and VRA in precision agriculture. Finding solutions requires collaboration, policy support, and ongoing efforts to enhance accessibility and usability of these technologies in farming practices.

Strategic Solutions for Overcoming Challenges in GIS, GPS, and VRA Integration in Precision Agriculture

Addressing the major challenges associated with GIS, GPS, and VRA in precision agriculture requires a multifaceted approach involving technology development, education, policy support, and collaboration among stakeholders. Here are potential solutions to the identified challenges:

1. **Data Integration and Compatibility:**

Advocate for standardized data formats and interoperability standards across GIS, GPS, and VRA technologies. Encourage collaboration among technology providers to develop compatible interfaces and promote open-data initiatives.

ISBN : 9788119821327

2. **Complexity and Technical Expertise:**

Develop user-friendly interfaces and intuitive software that require minimal technical expertise. Implement comprehensive training programs for farmers, extension services, and agricultural professionals to enhance their understanding and skills.

3. **Costs and Affordability:**

Explore cost-sharing models, government subsidies, and advancements in technology that may lead to cost reductions over time. Promote awareness of the long-term benefits and return on investment associated with precision agriculture.

4. **Data Security and Privacy Concerns:**

Implement robust data encryption, secure cloud storage solutions, and transparent data-sharing policies. Establish clear guidelines for data usage and storage, ensuring farmers' sensitive information is protected. Build trust through transparent data governance.

5. **Limited Connectivity in Remote Areas:**

Invest in alternative communication technologies, such as satellite-based solutions, to overcome poor internet connectivity in remote agricultural regions. Collaborate with telecommunication providers to enhance infrastructure development in these areas.

6. **Resistance to Change:**

Conduct awareness programs, workshops, and training sessions to educate farmers on the benefits of GIS, GPS, and VRA technologies. Engage in community-based initiatives to address concerns and build trust. Showcase successful case studies of technology adoption.

7. **Regulatory and Policy Challenges:**

Collaborate with policymakers to develop clear guidelines, standards, and incentives that support the responsible adoption of GIS, GPS, and VRA in precision agriculture. Advocate for policies that encourage technological innovation and sustainability.

ISBN : 9788119821327

8. **Scalability and Customization:**

Develop flexible systems that can be easily scaled up or customized to suit various farm sizes and types. Provide adaptable solutions that allow farmers to choose and integrate technologies based on their specific needs.

9. **Data Accuracy and Quality:**

Implement rigorous data validation processes, invest in high-resolution satellite imagery, and regularly update GIS databases for improved accuracy. Promote the use of precision equipment for data collection and emphasize the importance of data quality.

10. **Collaboration and Communication:**

Facilitate platforms for collaboration, such as forums, workshops, and conferences, to encourage information-sharing among different stakeholders. Establish communication channels to address concerns and share best practices. Encourage partnerships between technology providers, farmers, and researchers.

Implementing these solutions requires a coordinated effort from technology providers, policymakers, agricultural extension services, and farmers. By addressing these challenges systematically, the integration of GIS, GPS, and VRA in precision agriculture can be optimized, leading to more effective farming practices.

Conclusion

In conclusion, the integration of GIS, GPS, and VRA has fundamentally reshaped the landscape of agriculture. These technologies, once on the periphery, are now at the heart of precision farming. The ability to harness spatial data, navigate fields with precision, and dynamically adjust resource applications has not only increased efficiency but has ushered in a new era of sustainability. As we reflect on the uses of GIS, GPS, and VRA in precision agriculture, it is evident that these tools are not mere conveniences but indispensable assets. They empower farmers to make informed decisions, optimize

ISBN : 9788119821327

resource use, and contribute to environmentally conscious practices. The journey towards precision agriculture continues, fuelled by advancements in technology and an unwavering commitment to cultivating a more sustainable and productive future.

References

1. Aronoff S. (1989). Geographic information systems: a management perspective. Ottawa: WDL Publications 294 pp.
2. Pierce FJ, Nowak P (1999) Aspects of precision agriculture. Adv Agron **67**:1–85
3. Schroder D, Haneklaus S and Schung E. (1997). Information management in precision agriculture with LORIS. In Precision Agriculture'97, Vol.II: Technology, IT and Management (Ed. J.V. Stafford). BIOS Scientific Publishers Ltd., Oxford, UK. pp.821-826

ISBN : 9788119821327

CHAPTER - 6

"Role of Geospatial Technology in Agriculture"

Dr. K. M. Gojiya1*, Er. B. A. Karangiya2 and Dr. S. K. Chavda3

1Research Associate, Agriculture Research Station (Fruit Crops),
Junagadh Agricultural University,
Mahuva, Gujarat, India-364290

2Senior Research Assistant, College of Agril. Engg. & Tech., Junagadh Agricultural University,
Junagadh, Gujarat, India-362001

3Assistant Professor, C. P. College of Agriculture, Sardar Dantiwada Agricultural University, Sardar
Krushinagar, Dantiwada, Gujarat, India-385506

Corresponding Author Email: kashyapgojiya@gmail.com
Phone Number: 8306678888

Abstract

In the dynamic landscape of agriculture, geospatial technology, encompassing Remote Sensing (RS), Geographic Information System (GIS), and Global Positioning System (GPS), emerges as a transformative force. This chapter explores its multifaceted role in reshaping traditional practices, enhancing precision, and ensuring sustainable development in agriculture. From the fundamentals of RS, GIS, and GPS to applications in crop management, soil and water optimization, pest control, and sustainable practices, this exploration unfolds the profound impact of geospatial technology. Real-world case studies illustrate its success in increasing yields, resource efficiency, and timely pest management. Despite challenges, the chapter envisions the future, discussing trends like AI integration, UAVs, and blockchain, and emphasizes the need for continued research, education, and global collaboration. Geospatial technology stands as a testament to agriculture's adaptability, steering it towards a resilient and efficient future.

ISBN : 9788119821327

Key words: *Geospatial Technology, Precision Agriculture, Remote Sensing, Sustainable Agriculture*

Introduction

In the ever-evolving landscape of agriculture, the integration of advanced technologies has become pivotal in reshaping traditional practices and ushering in an era of unprecedented precision and efficiency. Among these transformative technologies, geospatial technology stands out as a beacon of innovation, offering a suite of tools that harness the power of Remote Sensing (RS), Geographic Information System (GIS), and Global Positioning System (GPS). This chapter delves into the multifaceted role of geospatial technology in agriculture, exploring its profound impact on monitoring, management, and sustainable development within the agricultural sector.

Geospatial technology, a collective term encompassing RS, GIS, and GPS, provides a unique lens through which we can observe, analyze, and optimize various facets of agricultural processes. With the ability to capture, interpret, and manipulate spatial data, these technologies empower farmers, researchers, and policymakers to make informed decisions that transcend the limitations of traditional practices. As we navigate through the intricate web of applications, challenges, and future prospects, it becomes evident that geospatial technology is not merely a set of tools but a transformative force steering agriculture towards a more sustainable and productive future.

The synergy between RS, GIS, and GPS unfolds a realm of possibilities, from precision crop management to efficient resource allocation, enabling a paradigm shift in how we approach the age-old endeavour of feeding the global population (Li-Juan *et al.*, 2005). This chapter aims to unravel the intricacies of geospatial technology, showcasing its applications across diverse agricultural domains, elucidating case studies of successful implementations, and contemplating the challenges and future trends that lie on the horizon.

ISBN : 9788119821327

Join us on a journey through the dynamic intersection of agriculture and geospatial technology, where satellites become the eyes in the sky, maps transform into decision-making tools, and GPS guides the hands that cultivate the fields. The utility of RS, GIS, and GPS in agriculture is not just a technological advancement; it is a testament to the resilience and adaptability of the agricultural sector in the face of a changing world.

As we embark on this exploration, it is our hope that the insights presented in this chapter will inspire further research, foster collaboration, and contribute to the ongoing discourse on the sustainable future of agriculture and allied sciences.

Fundamentals of Geospatial Technology

A. Remote Sensing (RS)

1. Definition and Principles

Remote Sensing (RS) is a technology that enables the acquisition of information about an object, area, or phenomenon without direct physical contact. This is achieved through the use of sensors mounted on platforms such as satellites or aircraft, capturing data from a distance. The underlying principle involves the detection and measurement of electromagnetic radiation reflected or emitted by the target, converting it into valuable information for analysis.

At its core, remote sensing operates on the premise that different objects interact with electromagnetic radiation in distinct ways, allowing us to discern a variety of features and properties. The electromagnetic spectrum, from visible light to microwaves, serves as the palette through which remote sensing paints a detailed portrait of Earth's surface and atmosphere.

2. Types of Remote Sensing Technologies

Remote sensing encompasses various technologies, each tailored to specific wavelengths and applications. These technologies can be broadly categorized into active and passive sensing (Cao and Lam, 2023):

ISBN : 9788119821327

a. **Passive Remote Sensing:** Relies on naturally occurring radiation, such as sunlight, and measures the reflected or emitted energy. Visible and infrared sensors fall under this category, capturing information about surface properties like vegetation health and land cover.

b. **Active Remote Sensing:** Involves emitting energy and measuring the reflected signals. Radar (Radio Detection and Ranging) and LiDAR (Light Detection and Ranging) are common active sensing technologies. Radar, for instance, is adept at penetrating cloud cover and providing insights into topography and surface moisture.

3. Applications in Agriculture

Remote sensing has become an invaluable tool in modern agriculture, offering a wealth of applications that enhance crop management, resource optimization, and environmental monitoring:

a. **Crop Health Assessment:** Satellite imagery and aerial photography enable the assessment of crop health by detecting subtle variations in vegetation vigor. This aids in early identification of stress factors such as pests, diseases, or nutrient deficiencies.

b. **Land Cover Classification:** Remote sensing facilitates the classification of land cover types, providing crucial information for land-use planning, precision agriculture, and ecosystem monitoring.

c. **Yield Prediction and Optimization:** Through the analysis of spectral signatures, remote sensing contributes to yield prediction models. This enables farmers to optimize inputs, manage resources efficiently, and enhance overall productivity.

In essence, remote sensing acts as a sentinel, extending our observational capabilities beyond human vision, and its applications in agriculture continue to evolve, offering unprecedented insights into the dynamic interactions between crops and their environment.

Applications of Geospatial Technology in Agriculture

A. Crop Monitoring and Management

ISBN : 9788119821327

1. Remote Sensing for Crop Health Assessment

Remote sensing technologies play a pivotal role in monitoring and assessing the health of crops. By capturing and analyzing spectral data, remote sensing can identify subtle changes in vegetation, allowing for early detection of stressors such as pests, diseases, or nutrient deficiencies. This proactive approach enables farmers to implement timely interventions, ensuring optimal crop health and yield.

2. Precision Agriculture using GIS and GPS

The fusion of Geographic Information System (GIS) and Global Positioning System (GPS) technologies has given rise to precision agriculture, revolutionizing crop management practices. GIS facilitates the creation of detailed maps that depict variations in soil composition, topography, and crop health. Combined with GPS, farmers can precisely navigate and apply resources, such as fertilizers or pesticides, based on spatial variability within the fields. This targeted approach enhances efficiency, reduces waste, and maximizes overall agricultural productivity (MohanRajan *et al.*, 2020).

3. Yield Prediction and Optimization

Geospatial technology contributes to yield prediction models through the analysis of historical data, environmental factors, and crop health indicators. By integrating remote sensing data, GIS, and GPS, farmers can make informed decisions on resource allocation, leading to optimized yields. This data-driven approach enhances the overall sustainability and economic viability of agriculture.

B. Soil and Water Management

1. Soil Mapping and Analysis with GIS

GIS is instrumental in soil mapping and analysis, providing farmers with detailed spatial information about soil properties and composition. This allows for precise delineation of soil types, nutrient levels, and pH variations across a field. Such information enables tailored soil management strategies, optimizing crop growth conditions and minimizing the environmental impact of agricultural practices.

ISBN : 9788119821327

2. Monitoring Water Resources through Remote Sensing

Remote sensing technology aids in the efficient monitoring of water resources, crucial for sustainable agriculture. Satellites equipped with sensors can detect changes in soil moisture levels, assess water availability, and monitor irrigation patterns. This information empowers farmers to make data-driven decisions regarding irrigation scheduling, ensuring judicious water use and mitigating the risk of water-related stress on crops.

3. Precision Irrigation using GPS Technology

GPS technology enhances precision irrigation by providing accurate location-based information. Automated irrigation systems guided by GPS data can deliver water precisely where and when it is needed, optimizing water usage and promoting water conservation. This approach not only improves crop yields but also contributes to the overall sustainability of water resources in agriculture.

C. Pest and Disease Management

1. Early Detection of Pests and Diseases using Remote Sensing

Remote sensing aids in the early detection of pests and diseases by capturing subtle changes in crop health indicators. This proactive monitoring allows for timely intervention, preventing the spread of diseases and minimizing crop losses. Satellite and drone imagery, coupled with advanced image analysis techniques, enable farmers to identify potential threats before they escalate.

2. GIS Applications in Spatial Analysis of Pest Outbreaks

GIS facilitates spatial analysis of pest outbreaks by integrating information on pest distribution, environmental factors, and crop conditions. By visualizing spatial patterns, farmers can develop targeted pest management strategies, deploying resources effectively to mitigate the impact of outbreaks. This geospatial approach enhances the efficiency of pest control measures.

ISBN : 9788119821327

3. GPS-Enabled Monitoring of Pest Control Measures

GPS technology supports real-time monitoring of pest control measures. By tracking the movement of agricultural machinery and the application of pesticides, farmers can ensure precise and uniform coverage of affected areas. This not only improves the effectiveness of pest control but also minimizes the environmental impact associated with the overuse of pesticides.

D. Sustainable Agriculture Practices

1. Precision Farming for Resource Efficiency

Precision farming, enabled by geospatial technology, promotes resource efficiency by tailoring agricultural practices to the specific needs of each field. Through the integration of RS, GIS, and GPS, farmers can optimize the use of inputs such as water, fertilizers, and pesticides, reducing waste and environmental impact while maximizing crop yields.

2. Environmental Impact Assessment using Geospatial Tools

Geospatial tools facilitate environmental impact assessments in agriculture by analyzing factors such as land use changes, water quality, and biodiversity. GIS allows for the identification of potential environmental risks, guiding the implementation of sustainable practices that minimize negative impacts on ecosystems.

3. Integration of Geospatial Technology in Organic Farming

Geospatial technology can play a crucial role in organic farming by providing insights into soil health, crop diversity, and pest management without the use of synthetic inputs. GIS can assist in planning organic farming practices, optimizing land use, and ensuring adherence to organic certification standards.

In essence, the applications of geospatial technology in agriculture represent a paradigm shift, offering solutions to age-old challenges while fostering sustainable practices that balance productivity with environmental stewardship. The integration of RS,

ISBN : 9788119821327

GIS, and GPS empowers farmers to make informed decisions, driving the agricultural sector towards a more resilient and efficient future.

Challenges and Future Trends

A. Current Challenges in the Adoption of Geospatial Technology in Agriculture

Despite the significant strides made in integrating geospatial technology into agriculture, several challenges persist:

1. Accessibility and Affordability

Limited access to geospatial technology tools, particularly in developing regions, poses a significant barrier. High costs associated with acquiring and implementing these technologies can hinder widespread adoption among small-scale farmers.

2. Technical Capacity and Education

A lack of technical expertise and education on geospatial technology among farmers and practitioners impedes its effective use. Bridging the knowledge gap and providing training programs are essential for maximizing the benefits of these technologies.

3. Data Quality and Interoperability

Ensuring the quality and interoperability of geospatial data remains a challenge. Harmonizing data from various sources and maintaining data accuracy are critical for producing reliable insights and facilitating collaboration across different platforms.

B. Emerging Trends and Innovations

1. Artificial Intelligence and Machine Learning Integration

The integration of artificial intelligence (AI) and machine learning (ML) with geospatial technology is a burgeoning trend. These technologies enhance the analysis of large datasets, enabling more accurate predictions, automated decision-making, and improved resource allocation in agriculture.

ISBN : 9788119821327

2. Unmanned Aerial Vehicles (UAVs) for Precision Agriculture

The use of UAVs, commonly known as drones, is gaining momentum in precision agriculture. Drones equipped with advanced sensors capture high-resolution imagery, providing farmers with real-time data for crop monitoring, disease detection, and precise application of inputs.

3. Blockchain for Supply Chain Transparency

Blockchain technology is being explored to enhance transparency and traceability in agricultural supply chains. By leveraging the decentralized and secure nature of blockchain, stakeholders can track the journey of agricultural products from farm to market, ensuring authenticity and reducing fraud.

C. Potential Future Developments and Advancements

1. Integration of 5G Technology

The rollout of 5G technology holds promise for enhancing connectivity and data transfer speeds in remote agricultural areas. This advancement can support real-time data collection, analysis, and decision-making, further optimizing the efficiency of geospatial technology applications.

2. Quantum Computing for Complex Analyses

Quantum computing has the potential to revolutionize complex spatial analyses in agriculture. Its computational power could enable faster processing of large datasets, facilitating advanced simulations, modeling, and optimization of agricultural systems.

3. Increased Global Collaboration and Data Sharing

As international collaboration and data sharing initiatives gain traction, a more interconnected global network of geospatial information is likely to emerge. This can foster cross-border research, knowledge exchange, and a more comprehensive understanding of global agricultural trends and challenges.

ISBN : 9788119821327

Conclusion

A. Recap of Key Points Discussed in the Chapter

Throughout this chapter, we've explored the multifaceted role of geospatial technology in agriculture. From the fundamentals of remote sensing, GIS, and GPS to their applications in crop monitoring, soil and water management, pest control, and sustainable practices, the transformative impact of these technologies on modern agriculture has been evident.

B. Emphasis on the Transformative Role of Geospatial Technology in Agriculture

Geospatial technology has emerged not only as a set of tools but as a transformative force reshaping the landscape of agriculture. The case studies and success stories highlighted the tangible benefits, from increased crop yields and resource efficiency to timely pest control measures and reduced crop losses. The integration of RS, GIS, and GPS empowers farmers to make informed decisions, steering agriculture towards a more sustainable and productive future.

C. Encouragement for Continued Research and Development in the Field

As we navigate the current challenges and anticipate future trends, it is imperative to emphasize the importance of continued research and development in the field of geospatial technology in agriculture. Addressing accessibility issues, advancing technical education, and exploring innovative solutions will contribute to the widespread adoption and meaningful impact of these technologies across diverse agricultural landscapes.

In conclusion, geospatial technology has the potential to revolutionize agriculture, offering a dynamic toolkit for addressing contemporary challenges and shaping the future of sustainable and resilient food production. By fostering collaboration, embracing emerging trends, and investing in research, we can unlock the full

ISBN : 9788119821327

potential of geospatial technology, ensuring its transformative role in the agricultural sector.

References:

1. Li-Juan, C., Van Paddenburg, A., & Man-Yin, Z. (2005). Applications of RS, GIS and GPS technologies in research, inventory and management of wetlands in China. *Journal of Forestry Research*, *16*(4), 317-322.
2. Cao, C., & Lam, N. S. N. (2023). Understanding the scale and resolution effects in remote sensing and GIS. In *Scale in remote sensing and GIS* (pp. 57-72). Routledge.
3. MohanRajan, S. N., Loganathan, A., & Manoharan, P. (2020). Survey on Land Use/Land Cover (LU/LC) change analysis in remote sensing and GIS environment: Techniques and Challenges. *Environmental Science and Pollution Research*, *27*, 29900-29926.

ISBN : 9788119821327

CHAPTER – 7

Applications of Remote sensing, GIS and GPS in Floriculture

Nagajyothi G N., Pratheeksha C T. and Udaya T V
Ph. D Scholars., Department of Floriculture and Landscaping, University of Horticultural Sciences Bagalkot, Karnataka-587104
Email Id. Nagajyothichary9891@gmail.com

Abstract

Crop monitoring and management is a critical aspect of modern floriculture, and it involves the systematic observation and control of crops to optimize yield, quality, and resource usage. This can be done by "precision farming", is a combination of FARMING that merges with the new technologies for optimum input use by equitable distribution of natural resources, address both economic and environmental issues that surrounds production agriculture today. This farming leverages a wide range of technologies like GPS, GIS, drones, Sensors, VRT, etc., to make farming more efficient, sustainable, and productive. Thus, this can helps to feed the growing population, keeping the environmental degradation to the minimum. In this chapter author address about the Remote sensors, GIS and GPS principles, technologies, assessment and components, that makes the farming more precise and efficient.

Key words: GIS, GPS, Precision farming, Sensor technologies

I. Introduction

The field variations and season variations are very common in production systems. Keeping in view of spatial variation in the fields and using newly available technologies, the mode and blend of fertilizer application varied. Such type of concept was initiated in 1980's as "precision agriculture" with the use of sensors. If the inputs are applied uniformly over the entire field without considering the variations within the fields, it may result either in excess or less that required for

ISBN : 9788119821327

maximum yields in some pockets of field. With this background idea, the concept of precision farming or site specific farming is being evolved starting from farming by the foot, farming by soil (Mandal and Ghosh, 2000).

Precision farming or smart farming (Khosla, 2001), is an approach to farming that utilizes technologies and principles to optimize various aspects of production by assessing spatial and temporal variability to increase the efficiency, productivity, and sustainability of farming operations while reducing waste and environmental impact by several technologies like GPS, RS, VRT (Variable rate technology) etc. (Brisco *et al*., 1998). The major components of precision farming are; Geographical information system (GIS), Geographical positioning system (GPS), Remote sensing, Variable rate technology, NDVI (Normalized Difference Vegetative Index), Nutrient expert system, SSNM (Site Specific Nutrient Management), Bio-intensive farming, Real-time nitrogen management, DRIS (Diagnosis and Recommendation Integrated System) approach, Soil testing and yield monitoring. In this chapter mainly emphasize the relevance and importance of RS, GIS, and GPS technologies specifically in the context of floriculture, showcasing how these tools can address unique challenges and enhance various aspects of flower cultivation.

Objectives

- Promotion of new venture in the 'Floriculture' bringing together various component of floriculture to exploit the variability
- Reduction in cost of cultivation due to site-specific crop management practices
- Increase in production efficiency of inputs due to site-specific management of inputs
- Reduction in soil and environmental pollution
- Soil and ground water conservation
- Improved product quality
- Energy conservation

ISBN : 9788119821327

Significance of applying Remote sensing, GIS and GPS in floriculture

Crop Health Assessment:

- NDVI is a commonly used remote sensing index that measures plant health and greenness. It helps identify areas of stress and potential issues within a field.
- Observing and identifying pests and diseases, often with the help of sensor data, allows for timely intervention through targeted treatments
- Data from monitoring systems can inform decisions on when and where to apply pesticides or herbicides, reducing unnecessary chemical use.

Flowering time and duration management (Zhang *et al*., 2020)

- Capture high-resolution images using various imaging modalities, such as RGB (Red, Green, Blue) cameras, multispectral or hyperspectral imaging, and even thermal imaging to capture different aspects of flowering structures
- Image-based phenotyping refers to the use of images to quantify and analyze various traits and characteristics of plants like plant architecture, flowering intensity, number of flowers, flower size, colour, and spatial distribution.
- Combine remote sensing data with information about pollinator biology and behaviour to understand the relationship between floral resources and pollinator activity.

Monitor the availability and distribution of floral resources in the landscape (Gonzales *et al*., 2022)

- Utilize high-resolution satellite imagery or aerial photography to capture large-scale landscapes and identify areas with flowering patterns and abundant floral resources

ISBN : 9788119821327

- Geospatial analysis to identify areas of high importance for pollinators, such as meadows, wildflower patches, or other natural habitats.

Irrigation Management:

- Sensors measure the water content in the soil, helping farmers determine when and how much to irrigate to prevent over- or under-watering.
- Modern systems can adjust water delivery based on real-time data, weather forecasts, and the specific needs of the crop.

Biomass and Growth Monitoring

- RS technologies can be used to monitor the biomass and growth patterns of floral crops throughout their life cycle.
- This information aids in assessing the effectiveness of cultivation practices and optimizing inputs.

Yield Estimation

- Satellite and aerial imagery, coupled with advanced analytics, assist in estimating floral crop yields.
- Remote sensing data can be used to assess factors such as plant density, flower size, and overall biomass.

Nutrient Management


- VRT enables the precise application of fertilizers based on soil nutrient levels and crop requirements. It minimizes excess nutrient use and reduces environmental impact.
- Fertigation systems combine irrigation and fertilizer application delivering nutrients directly to the root zone, ensuring efficient use.

Crop Planning and Forecasting:

- Historical and real-time data can be used to predict crop yields, helping with market planning and resource allocation.
- Monitoring data can inform decisions on crop rotation and selection for improved soil health and pest management.

ISBN : 9788119821327

- Floriculturists can anticipate changes in weather patterns, optimize planting and harvesting schedules, and minimize the impact of adverse conditions.



Red - low yield, yellow & orange – intermediate yield and green- high yield

Optimizing supply chain management (Odera and Karanja, 2019)

- GIS can assist in selecting optimal locations for floriculture farms based on factors such as climate, soil conditions, and proximity to transportation networks. It aids in planning the distribution of flower farms to ensure efficient production and logistics.
- Helps monitor and manage flower crops by integrating spatial data on weather conditions, soil moisture, and other relevant factors. This information can be used to optimize irrigation schedules, fertilization, and pest control measures**.**
- By integrating GPS and RFID data, stakeholders can gain visibility into the location and condition of flowers during transportation
- Route optimization minimizes transportation costs, reduces delivery times, and lowers environmental impact
- GIS can incorporate market data, such as consumer demographics and preferences, to analyze and forecast demand for specific flower varieties in different regions. This helps in aligning production with market needs.
- GIS technology, combined with temperature and humidity sensors, can monitor the conditions of refrigerated storage and

ISBN : 9788119821327

transportation facilities, ensuring the quality of flowers is preserved.

Understanding the variability

Variation in soil properties: Variability in soil property may be due to soil topography features like slope, length of slope, curvature of slope or soil physical properties like soil depth, percentage of coarse fragments, soil texture and soil structure or chemical factors like soil reaction, electrical conductivity, available nutrients and cation exchange capacity.

Variation in Weed densities: Depends on spatial variations in soil properties and weed seed spread. In the study on weed densities, it was found that weed densities were significantly greater in areas that had higher greater organic carbon and finer texture. In such soils more uniform weed control can be achieved at minimal costs by varying herbicidal applications.

Assessing the variability: Effectiveness of precision agriculture depends on accurate and efficient mapping. Map may depict the spatial distribution of one or more soil properties or processes like growth and yield of crops, weed patches, pathogens or pest populations etc. There are usually three types of maps *viz.,* condition maps, prescription maps and performance maps.

Prescription maps are derived from condition maps and they form the basis for variable rate technology, based on those maps, variable rate of seeds, nutrients, herbicides, irrigation water, fungicides, pesticides etc. are applied to fields in site-specific management.

Performance maps record either the inputs applied to fields like seeds, fertilizers, pesticides etc., or outputs like growth, crop yields and quality of economic produce.

Condition maps (Patil and Bhalerao, 2013) are prepared mainly by: 1. Soil survey; 2. Interpretation of a network of points; 3. High resolution sensing and 4. Modelling

ISBN : 9788119821327

1. **Soil survey:** Is the new agricultural management concept called "farming by soil type", However, the existing soil survey have limited value in explaining the spatial variability within the fields. The value of soil survey in precision agriculture can be improved by intensifying the map scales to finer scale resolution.
2. **Interpolation of point samples:** In this process, sampling is done at definite spatially arranged points. Then by using a range of appropriate statistical procedures, these points are interpolated to produce a spatial estimation for the whole farm area. These spatial statistical procedures can be repeated over time to estimate the temporal variability. Among the factors, soil property mapping are most affected by number of soil samples, the distance between sampling locations and the choice of interpolation procedure.
3. **High resolution sensing:** Technical trend of image sensing are hyper spectral, 3 dimensional and active sensing. Ordinary multi spectral imaging is available for obtaining information on colour, pigments, shape and growth of individual plants and parts, change in water status in plants and soils, soil properties. The hyper spectral image sensing capable of resolving several hundreds of spectral bands in the region from visible to short wave, make it possible to provide exactly more information by analysis of continuous spectral properties in comparison with ordinary multi-spectral analysis. Thermal infrared imagining, a passive spectral imaging method, is effective for early detection of plant stress as well as for measurement of surface temperature of plants and soils. Fluorescence image sensing provides information on changes in cell walls and bleaching of plant pigments. This concept uses to detect the changes in photosynthetic activity caused due to biotic and abiotic stresses in plants.

ISBN : 9788119821327

4. **Modelling:** Is an important tool in precision agriculture to stimulate spatial and temporal variation in soil properties, pests, crop yield and environmental performance of cropping systems. The remote sensing modelling component provides the ability to assess plant water status at high spatial resolution and the crop model provides reasonable yield prediction of water stressed plots using only crop water stress index measurements during the season to indicate inadequate availability of water in plant. A combination of crop modelling and remote sensing can identify crop management zones and causes yield variability.
5. **Image processing**: The image processing method is a simple and fast way to detect weeds with a commercial digital camera. The image-processing algorithm first locates the green objects in the digital image, and then calculates the weed density based on greenness ratio. The image processing, the green objects in each image were identified using greenness method that compared with red, green objects and blue (RBG) intensities.

Components

1. Global Positioning System (GPS)

GPS is a navigation system based on a network of satellites that helps users to record

positional information with an accuracy of between 100 and 0.01 and provides continuous

position information in real time, while in motion. Having precise location information at any

time allows soil and crop measurements to be mapped. GPS allows farmers to locate the exact

position of field information, such as soil type, pest occurrence, weed invasion, water holes,

boundaries and obstructions. There is an automatic controlling system, with light or sound

ISBN : 9788119821327

guiding panel (DGPS), antenna and receiver. GPS is a network of orbiting satellites, sending
precise positional details from space back to earth. The signals are obtained by GPS receivers,
to calculate the exact position and time. This becomes particularly handy when farmers need
to make decisions regarding seed plantation, herbicides, pesticides, fertilizers and irrigational
needs.

Applications of GPS in floriculture

- Using GPS for accurate field mapping and delineation of floral crops. This helps in creating digital maps of flower beds, rows, and other features within the cultivation area.
- Ensuring precise spacing and alignment of floral crops for optimal growth and resource utilization.
- Enhancing logistical operations by tracking the location of harvesting equipment and coordinating activities efficiently.
- Integrating GPS technology into tractors, harvesters, and other equipment for accurate tracking and navigation within flower farms.
- Ensuring precise and efficient movement of equipment during various cultivation operations.
- Optimizing water distribution based on the specific needs of different areas within the flower farm, conserving water resources.
- Integrating GPS data with soil analysis results to create spatially accurate soil maps, aiding in targeted soil management practices**.**
- GPS-enabled drones for aerial surveillance of flower crops helps in monitoring the health and growth

ISBN : 9788119821327

- Associating spatial information with attributes such as crop health, pest incidence, and flowering stages for comprehensive analysis.
- Analyzing spatial variations in yield data to identify factors influencing productivity.
- Enhancing efficiency and reducing errors by guiding workers to specific locations within the flower farm during tasks such as crop inspection, harvesting, and maintenance.
- Implementing precision spraying techniques to target specific areas with pesticides or fertilizers based on spatial data.
- Enhancing traceability by associating spatial information with specific batches of floral products throughout the supply chain for quality control purposes.

In landscape (Nijhuis, 2014):


- Assist landscapers in efficiently navigating through large properties, especially in parks, golf courses, or expansive gardens. It helps optimize the route for mowing, planting, or maintenance tasks.
- For accurately mapping property lines and boundaries
- Enabled drones are employed for aerial surveys, allowing landscapers to capture high-resolution imagery and topographic data. This information aids in the design and planning of landscaping projects.

2. Geographic information system (GIS)

The use of GIS was started in 1960, a computerized mapping system to acquire, store, analyse and display information that is specially referenced to the earth. It is a software that imports, exports and processes spatially and temporally geographically distributed data. This system comprises hardware, software and procedures designed to support the compilation, storage, retrieval and analysis of feature attributes and location data to produce the map. The location

ISBN : 9788119821327

can be expressed in many different ways, such as latitude and longitude, address, or ZIP code. It can include information about the landscape, such as the location of streams, different kinds of vegetation, and different kinds of soil, sites of factories, farms, and schools, or storm drains, roads, and electric power lines, yield, soil survey, crop type, nutrient levels and prone pests. Once analysed, this information is used to understand the relationships between the various elements affecting a crop on a specific site (Singh *et al.,* 2014).



Components of GIS system

A Geographic Information System (GIS) is a complex system designed to capture, store, analyze, manage, and present spatial or geographic data. The components of a GIS system typically include hardware, software, data, people, and methods.

1. **Hardware:**

- Computers: GIS systems require computers to run the necessary software for data processing, mapping, and analysis.
- Input Devices: Devices such as digitizers, GPS receivers, and scanners are used to input spatial data into the GIS.
- Output Devices: Printers, plotters, and monitors are used to display or output maps and analytical results.

ISBN : 9788119821327

2. **Software:**

- GIS Software: This is the core software that allows users to perform various GIS functions, including data input, analysis, visualization, and output. Examples of GIS software include ArcGIS, QGIS, and Google Earth**.**

3. **Data:**

- Spatial Data: The primary data in GIS includes spatial information that represents the location, shape, and characteristics of geographic features. This can include vector data (points, lines, polygons) and raster data (grids of cells representing values).
- Attribute Data: Non-spatial information associated with geographic features, stored in tabular formats. Attributes provide additional details about the features**.**

4. **People:**

- GIS Professionals: Individuals with expertise in GIS, including GIS analysts, technicians, and managers, who use the system to analyze and interpret spatial data.
- End Users: Those who use GIS outputs and maps for decision-making, planning, or research in various fields, such as urban planning, agriculture, environmental science, and more.

Applications of GIS in Floriculture

Geographic Information Systems (GIS) find diverse applications in the field of floriculture, offering a range of tools for spatial analysis, decision-making, and resource management (Mandal *et al.,* 2021).

- Analyzing spatial data to select optimal locations for flower farms based on factors such as topography, soil types, and drainage patterns.
- Planning the layout of flower beds, greenhouses, and infrastructure using GIS to maximize efficiency.

ISBN : 9788119821327

- Implementing precision farming practices by analyzing spatial variability within flower fields. GIS helps in optimizing the application of fertilizers, pesticides, and water resources based on specific spatial characteristics.
- Conducting environmental impact assessments by analyzing spatial data to evaluate the potential environmental effects of floriculture operations.
- Identifying and mitigating the impact on ecosystems, water bodies, and adjacent land.
- Analyzing spatial patterns of disease and pest occurrences using GIS for targeted interventions.
- Developing strategies to minimize the spread of diseases and reduce the impact of pests on flower crops.
- Analyzing spatial data on market locations, transportation routes, and demand patterns to optimize supply chain logistics.
- Planning efficient transportation routes and distribution networks based on geographic considerations.
- Managing water resources efficiently by analyzing spatial data on soil moisture levels, precipitation patterns, and water availability.
- Implementing precision irrigation systems based on spatial information to conserve water.
- Tracking the spatial distribution of different flower varieties, monitoring growth patterns, and managing inventory levels.
- Integrating spatial data on weather patterns, temperature variations, and climatic conditions into GIS for monitoring the impact on flower crops.

In landscape (Nijhuis, 2014):

- GIS in Landscape planning
- GIS can be used to assess and analyze potential sites for landscape projects. Factors such as topography, soil types,

ISBN : 9788119821327


drainage patterns, and proximity to amenities can be considered in the site selection process

- Hydrological modeling to analyze water flow patterns, drainage systems, and watershed characteristics
- To analyze urban heat islands, helping designers understand areas prone to increased temperatures. This information is important for incorporating heat mitigation strategies in landscape design
- Helps in planning pedestrian and vehicular circulation

3. **Remote Sensing**

Remote sensing technology is useful tool for gathering much information simultaneously. A tool for collection, processing and analysis of data to obtain exact information from earth surface without coming into physical contact with it. Remote sensing is the science of obtaining information about objects or areas from a distance, typically from aircraft or satellites, these sensors are generally categorised as aerial or satellite sensors. They can indicate variations in the colours of the field that corresponds to changes in soil type, crop development, field boundaries, roads, water, etc. Data sensors can simply be hand-held devices, mounted on aircraft or satellite-based. Remotely-sensed data provide a tool for evaluating crop health, related to plant stress like moisture, nutrients, compaction, crop diseases and other plant health concerns are often easily detected in overhead images. Remote sensing can reveal in-season variability that affects crop yield, and can be timely enough to make management decisions that improve profitability for the current crop (Siddegowda and Devi, 2021).

ISBN : 9788119821327




Sensor technologies

Various technologies such as electromagnetic, conductivity, photo electricity and ultra sound are used to measure humidity, vegetation, temperature, texture, structure, physical character, humidity, nutrient level, vapour, air etc. Remote sensing data are used to distinguish crop species, locate stress conditions, identify pests and weeds, and monitor drought, soil and plant conditions. Sensors enable the collection of immense quantities of data without laboratory analysis. Location Sensors are the sensors that are used to determine latitude, longitude, and altitude within feet. Optical Sensors use light to measure soil properties. Electrochemical Sensors provide key information like pH and soil nutrient levels. Mechanical Sensors measure soil compaction. They use a probe that penetrates the soil and records resistive forces through use of load cells or strain gauges. Dielectric Soil Moisture Sensors measure the dielectric constant in the soil to assess moisture levels. Airflow Sensors measure soil-air permeability. Satellite-based remote sensing technologies are widely used to guide the global agricultural production from a regional to global scale. Most of the remote sensing technologies are confined to multispectral broad bands, which have limitations for precise quantitative estimation of soil and plant properties primarily because of the low spectral resolution. In India, the requirement for a

ISBN : 9788119821327

marketable remote sensing technology for precision agriculture is the delivery of information with the following characteristics: low turnaround time (24-48 hr), low data cost (~ 100 Rs./acre/season), high spatial resolution (at least 2 m multi-spectral), high spectral resolution.



Types of remote sensing technologies

- **Satellite-based Remote Sensing:** Utilizes orbiting satellites equipped with sensors to capture images and data from space. Provides global coverage and is widely used for monitoring large scale environmental changes.
- **Aerial Photography:** Involves capturing images of the Earth's surface from aircraft using traditional film or digital cameras. Offers high spatial resolution and is often used for detailed mapping and surveying.
- **Unmanned Aerial Vehicles (UAVs or Drones):** Small unmanned aircraft equipped with various sensors, including cameras and multispectral sensors. Enables high-resolution and flexible data collection for smaller areas and specific applications.
- **Lidar (Light Detection and Ranging):** Uses laser beams to measure distances and create detailed three-dimensional maps

ISBN : 9788119821327

of the Earth's surface. Commonly employed for topographic mapping, vegetation analysis, and urban planning.

- **Hyperspectral Imaging:** Captures images in multiple narrow and contiguous spectral bands, providing detailed spectral information. Useful for precise identification of materials and vegetation types.
- **Radar (Radio Detection and Ranging):** Uses radio waves to detect and measure the distance to the Earth's surface. Penetrates through clouds and vegetation, making it suitable for applications like land cover mapping and soil moisture estimation.
- **Thermal Infrared Imaging:** Measures the infrared radiation emitted by objects to assess their temperature. Valuable for applications like detecting heat anomalies, monitoring thermal pollution, and assessing energy balance.
- **Multispectral Imaging:** Captures images in multiple broad and distinct spectral bands.
  Used for agricultural monitoring, environmental assessments, and land cover classification.
- **Passive vs. Active Sensors:** Passive Sensors: Detect natural radiation emitted or reflected by the Earth's surface (e.g., optical sensors). Active Sensors: Emit their own radiation and measure the response (e.g., radar and lidar).
- **Photogrammetry:** Involves the measurement and interpretation of photographs to derive three-dimensional information about the Earth's surface. Used for creating accurate maps, terrain models, and assessing landscape changes over time.
- **Global Navigation Satellite System (GNSS):** Includes systems like GPS, GLONASS, and Galileo, which provide accurate positioning information for ground-based and mobile

ISBN : 9788119821327

applications. Integrated with remote sensing for precise georeferencing and navigation.

Applications of remote sensing/sensors

- **Normalised difference vegetation index (NDVI)**

NDVI value given by Green Seeker. It provides a crude estimate of vegetation health and means of monitoring the change in vegetation over time

$$\text{NDVI} = (NIR - PAR) \,/\, (NIR + PAR)$$

Where;

NIR is near-infrared radiation and PAR is photo synthetically active radiation. NDVI value range = +1 to -1 (+1 for dark green vegetation area and -1 for vegetation less area)

- **Micro-irrigation**

Frequent application of small quantities of water directly above and below the soil surface; through emitters or applicators placed along a water delivery line. Saves upto 70 percentage of water used in conventional methods and high irrigation efficiency more than 90%, no evaporation losses occur. Suitable for Orchards (grape vine, citrus, mango, guava) coconut, vegetable crops, row crops like sugarcane, cotton etc. and flowers (rose, jasmine etc.)

- **Protected cultivation and ornamental nursery crop production (Mahmud *et al.*, 2023)**


It is a cropping technique where the plants were controlled partially or fully as per the requirement of the plant species grown during their growth period. The main purpose of protected cultivation is to create a favorable environment for the sustained growth of plant so as to realize its maximum potential even in adverse climatic conditions. It is observed that farmers have to bear huge financial loss because of wrong prediction of weather and incorrect irrigation method to crops. In this contest with the evolution in wireless sensor technologies and miniaturized sensor devices, it is possible to uses them for automatic

ISBN : 9788119821327

environment monitoring and controlling the parameters of greenhouse. Sensors to assist for spatial data collection, irrigation control model, arrangement for supplying information to farmers, Variable-rate technology model and Green house parameter control system. This system can handle local field survey and collects data of soil water availability, biomass yield, soil compaction, soil fertility, leaf area index, leaf temperature, leaf chlorophyll content, local climate data, insect-disease-weed infestation, plant water status, and yield of grain etc. The total number of sensor nodes and actuators are depends on the size of greenhouse. The protected cultivation with smart framing and IOT driven pay a way for accurate maintenance and production (Ahmed, 2020).

Sensors used in green house: CO2 Sensor, humidity sensor, temperature sensor, soil analysis sensors, substrate sensors, light sensors, wind speed and direction sensors, precipitation sensors and weed sensors (Koning, 2006).

- **Temperature:** thermostat, thermocouples
- **Soil moisture:** Sensor probes, tensiometer
- **Light:** Pyranometer, PAR sensors, light level sensor
- **Surface moisture:** ADP-Agro Tech leaf Wetness Sensor (Siddharam *et al*., 2023)



Temperature and humidity sensors



Light sensor



InfraRed gas analyzer



Soil moisture sensors



ADP-Agro-Tech leaf wetness sensor

ISBN : 9788119821327

Conclusion:

Precision farming is a latest technology of utilizing tools like GIs, GPS and remote sensing for sustainable and equitable distribution of natural resources of our country. If properly used, it can solve the problem of food security; can address both economic and environmental issues that surround production agriculture today. However, this technology is in its infancy in India, there are numerous opportunities of adoption in floriculture to increase the quality and quantity in production of flowers with high grade quality to meet the demand of people's interest. Progressive farmers with guidance from the public and private sectors, and agricultural associations, can adopt it in their fields. Thus, this technology can help to satisfy the growing population by use of resources in precise and accurate manner.

Reference:

1. Ahmed, L. (2020). Automation, Sensor & IoT application in Protected Cultivation Technology.
2. Brisco, B., Brown, R. J., Hirose, T., McNarairn, H. and Staenz, K. (1998). Remote Sensing. 24: 315-327.
3. Gonzales, D., Ibarra, N. H. and Anderson, K. (2022). Remote Sensing of Floral Resources for Pollinators-New Horizons from Satellites to Drones. *Frontiers in Ecology and Evolution.*, 10: 1-16.
4. Khosla, R. (2001). Zoning in on Precision Ag. Colarado State University Agronomy *Newsletter*, *21*(1): 2-4.
5. Koning N. M. (2006). Models and Sensors in Greenhouse Control. *Acta Hort*., pp. 175-182.
6. Mahmud, M. S., Zahid, A. and Das, A. K. (2023). Sensing and Automation Technologies for Ornamental Nursery Crop Production: Current Status and Future Prospects. *Sensors*., 2023, 23: 2-25.

ISBN : 9788119821327

7. Mandal, D., and Ghosh, S. K. (2000). Precision farming – The emerging concept of agriculture for today and tomorrow. *Current Science*, *79*(12), 1644-1647.
8. Mandal, M., Paramanik, B., Sarkar, A. and Mahata, D. (2021). Precision farming in Floriculture. *International Journal of Research –GRANTHAALAYAH, 9*(1): 75-77.
9. Nijhuis S. (2014). Applications of GIS in landscape design research. Geodesign Summit Europe 2014, September 11–12, 2014 Delft University of Technology, Netherlands.
10. Odera, C. A. and Karanja, F. N. (2019). Leveraging GIS in Supply Chain Management for Floriculture Application: Nairobi County, Kenya*. Journal of Geographic Information System.* 11: 389-404.
11. Patil, S. S. and Bhalerao, S. A. (2013). Precision farming: The most scientific and modern approach to sustainable agriculture. *International Research Journal of Science and Engineering. 1*(2): 21-30.
12. Siddegowda, C. J. and Devi, A. J. (2021). A Study on the Role of Precision Agriculture in Agro-Industry. *International Journal of Applied Engineering and Management*. *5*(2): 57-67.
13. Siddharam., Aiswarya L., Kambale, J. B. and Rajesh, G M. (2023). Application of Internet of Things (IoT) in Protected Cultivation: A Review. *International Journal of Environment and Climate Change*. *13*(9): 1518-1529.
14. Zhang, C., Craine, W. A., McGee, R. J., Vandemark, G. J., Davis, J. B., Brown, J., Hulbert, S. H. and Sankaran, S. (2020). Image-Based Phenotyping of Flowering Intensity in Cool-Season Crops., *Sensors.,* 20: 1-17.

ISBN : 9788119821327

CHAPTER - 8

Remote Sensing and Geographic Information System in Forestry

Nikita Kumari[1] and Ajit Tippannavar[2]

[1]M. Sc. Research Scholar, Department of Silviculture and Agroforestry, College of Horticulture and Forestry, Jhalawar (Rajasthan), 326023

[2]M. Sc. Research Scholar, Department of Fruit Science, College of Horticulture and Forestry, Jhalawar (Rajasthan), 326023

*Corresponding author: sonuchoudhary126jat@gmail.com

Abstract

By leveraging remote sensing and GIS technology, quick and efficient tracking of tropical forest conditions and resources is made possible. This allows for the development of targeted policies aimed at promoting sustainable management of these vital ecosystems. The utilization of RS and GIS tools also simplifies the assessment of vegetation dynamics, revealing changes in land cover, wildlife loss, and deforestation rates. Additionally, through geographical and temporal analysis methods, we can accurately identify the impact of afforestation efforts and monitor other cumulative changes. By utilizing GIS and RS data, this study aims to map the current state of Nimbia Forest Reserve spanning from 1986 to 2010. The goal is to not only detect existing changes but also make predictions on potential future changes that could transpire within the next three decades.

INTRODUCTION

In recent years, the effects of natural or man-caused catastrophes have emphasized the need for developing a broader view of many natural processes. The "global monitoring" approach (Goward *et al*., 1987) tends to consider environmental problems from this perspective, and favors the understanding of the complexity underlying

ISBN : 9788119821327

natural phenomena. It's evident that this worldwide strategy requires the implementation of innovative techniques to effectively and efficiently gather, analyze, and present spatial data. This is the main objective behind the Geographic Information System (GIS) approach, which leverages the immense data processing and storage capabilities of computers (Burrough, 1986). This allows for the seamless updating and retrieval of spatial data, as well as the creation of dynamic cartographic models by merging various data layers within the database. Exploring the impact of forest fires is just one of the numerous valuable applications of GIS. Due to the multitude of factors affecting the start and spread of a fire, the use of an integrated analysis technique is imperative. This is especially crucial given the ever-changing nature of this phenomenon. Remote sensing imaging, in particular, proves to be extremely beneficial in such studies, offering a swift assessment of vegetation conditions and the environmental aftermath of fires. The integration of GIS has now become a prevalent practice in the fields of natural, engineering, and social sciences, offering reliable and reproducible methods for collecting, analyzing, and interpreting spatial data. The importance of forests as renewable natural resources cannot be overstated. Not only do they provide us with valuable timber, but they also play a crucial role in creating livable conditions for humans. But their value goes beyond just wood--forests also offer grazing land for animals, serve as habitats for wildlife, provide water resources, and serve as recreational havens. However, it takes efficient management to unlock the full potential of a forest. In today's constantly changing world, forest managers are met with a multitude of challenges. To navigate this complexity, this essay suggests utilizing GIS (Geographic Information System) as a potential solution. Throughout the course of history, forests have held immense value to the global human population, and this value continues to increase as we rely on them for fuel wood and protection. The world's vegetation is in a constant state of flux, with a myriad of factors contributing to the

ISBN : 9788119821327

deforestation of tropical forests (Lambin, 1996). As humans strive to meet their basic needs, they are rapidly altering forests, woodlands, and grasslands at an alarming rate (Pomery, 2004). In today's world, where population growth and demand for natural and man-made forest resources are on the rise, there is a growing necessity for up-to-date and precise spatial information. Geographic Information System (GIS) and Remote Sensing (RS) are now providing new tools for advance ecosystem management.

Remote Sensing (RS) and Geographic Information System (GIS) play crucial roles in forestry management and conservation. Here's how they contribute:

Remote sensing and Geographic Information System (GIS) are two interconnected technologies used in geospatial analysis and mapping.

Remote Sensing entails collecting data about the surface of the Earth without making actual physical contact. The majority of remotely sensed information utilized in spatial analysis and mapping is gathered as selected electromagnetic radiation. Usually, satellite sensors, drones, or cameras and sensors placed on airplanes are used for this. These devices collect data on a range of electromagnetic radiation wavelengths (including visible light, infrared, and microwave) to learn about the health of the vegetation, temperature fluctuations, and land cover, among other features of the Earth's surface.

GIS (Geographic Information System), GIS, on the other hand, is a software-based system that is used to organize, store, examine, and display geographic or spatial data. Users can use it to manage, organize, capture, process, store, and display geographic or spatial data. By integrating various data sources, including maps, satellite images, survey data, and more, GIS enables users to overlay various information layers and make well-informed decisions. These technologies frequently complement one another. The raw data (imagery, elevation data, etc.) is obtained through remote sensing and processed and

ISBN : 9788119821327

analyzed using GIS, which enables users to generate maps, carry out spatial analysis, identify changes over time, model scenarios, and make decisions based on the data collected. To make maps displaying land use patterns, identify regions vulnerable to natural disasters, track changes in vegetation over time, or plan urban growth based on numerous spatial characteristics, for instance, satellite images obtained through remote sensing can be incorporated into a GIS.

Remote Sensing Techniques in Forestry

Satellite imagery: Types, resolution, and applications in forest monitoring

LiDAR technology: Use in forest structure and canopy analysis

Aerial photography: Historical and modern methods for forest assessment

Hyperspectral imaging: Identification and mapping of tree species and health assessment

Thermal imaging: Monitoring forest fires and temperature changes

Remote Sensing in Forestry:

Forest Cover Mapping: Using satellite pictures, remote sensing aids in the mapping and tracking of changes in the forest cover over time. It makes it possible to identify land use changes, afforestation, and deforestation.

Tree Species Identification: Forest tree species can be identified and mapped using remote sensing techniques, particularly with high-resolution imagery and spectral analysis.

Health and Stress Assessment: By looking for indicators of illness, infestation, or stress, remote sensing can be used to monitor the health of forests. This can be achieved by employing specialized sensors to identify particular situations or by evaluating changes in vegetation indices.

ISBN : 9788119821327

Forest Fire Detection: To identify forest fires, track their spread, and evaluate post-fire damage, satellite imaging and heat sensors can be used.

Estimating Timber Volume: LiDAR (Light Detection and Ranging) technology, a type of remote sensing, helps determine the volume of timber and forest biomass, two essential components of sustainable forest management.

How Remote Sensing Works in Forest Management

A bird's-eye perspective of forests is made possible by satellites and aircraft fitted with remote sensors, which also record precise data about the structure, health, and ephemeral changes of the woods. After that, this data is processed, evaluated, and applied to different forest management duties.

Species Mapping: RS is used to map the distribution and health status of various tree species, which is important for the protection of biodiversity.

Deforestation and Forest Degradation Monitoring: Using RS data to monitor rates of deforestation, illicit logging, and changes in land use can aid in the development of conservation strategies and policies. Deforestation can be identified and measured globally thanks to remote sensing. The method assists in locating deforestation hotspots and monitoring changes over time as part of the remote sensing applications in deforestation. Having this knowledge is essential for putting into practice practical deforestation prevention measures.

Importance of Remote Sensing

The use of remote sensing in forestry is transforming our approach to managing our natural resources and creating new avenues for sustainability, economic expansion, and conservation. Drones, satellites, and Light Detection and Ranging (LiDAR) systems are examples of remote sensing technologies that change our understanding of and interactions with these vital ecosystems by providing data necessary for efficient forest management. Accurate and

ISBN : 9788119821327

timely information is necessary for the responsible management of our global forests. One comprehensive, effective, and economical way to collect this data is through remote sensing.

Remote Sensing in Forestry:

Mapping and Monitoring Forest Cover

RS tools, such satellite images, LiDAR, and aerial photography, offer data on forest cover, health, and changes over time. Precise mapping and tracking of forest cover are prerequisites for effective forest management. Large-scale data collection over vast wooded areas is made possible by remote sensing, which facilitates accurate mapping of the vegetation. With the use of this technology, one may discern between various types of forests, track variations in the forest cover, and measure these variations over time. When managing and safeguarding forest resources, foresters can make more informed decisions with the use of this mapping technique.

Forest Health and Disease Monitoring

The use of remote sensing technology is essential for spotting indicators of forest disease and declining tree health. We are able to take preventive measures by anticipating disease breakouts and insect infestations by identifying minute changes in the color of the canopy or in the density of the forest. Additionally, remote sensing aids in recovery efforts by providing information on how well management measures are working. Fire, insects, and disease are among the major natural disturbances that alter forested landscapes. Timely update information ensures inventory databases are current enough to support forest management planning and monitoring objectives.

Insect disturbance

Insect damage monitoring and mitigation are supported by integrated remote sensing and GIS studies, which identify and map insect outbreak and damage locations and characterize disturbance patterns in relation to mapped stand properties.

ISBN : 9788119821327

Using risk and hazard rating systems to model and forecast outbreak trends

supplying information to GIS-based decision support systems for pest control

Fire Detection and Management:

Early detection and effective firefighting are facilitated by the identification and monitoring of fire-prone regions using RS. Wildfires frequently destroy forests, resulting in serious ecological and financial harm. Even in the most remote areas, remote sensing satellites can identify fires and send out real-time alerts. Early detection and effective management of forest fires can be facilitated by the identification of fire-prone regions by thermal and infrared sensors.

Forest Planning and Management

The use of remote sensing in forest planning and management has proven transformational. These technology' abundance of information makes it possible to make well-informed judgments on reforestation, harvesting, and general forest management techniques. In order to preserve the longevity and health of our forests, it also helps in the creation of measures to lessen the effects of diseases, pests, and natural disasters.

Forest Conservation and Biodiversity Assessment

Planning and managing forests with remote sensing has shown to be revolutionary. Making informed decisions on reforestation, harvesting, and general forest management strategies is made feasible by the wealth of information available through these technologies. It also aids in the development of strategies to decrease the effects of illnesses, pests, and natural disasters, all of which are important for maintaining the lifespan and health of our forests.

Landscape Ecology, habitat, and biodiversity

According to Vogt *et al*. (1999), forest inventory and certification processes must take into account landscape ecological features that are connected to habitat and biodiversity in order to practice sustainable

ISBN : 9788119821327

forest management. The landscape's spatial patterns, particular forest conditions associated with habitats, and the biological processes connecting spatial pattern, habitat, and ecosystem functioning are the features of interest. One type of spatial pattern that can be easily obtained by categorizing remotely sensed data is land-cover information. In order to track changes over time and create and validate habitat models for specific species, remote sensing can offer reliable and consistent ways for developing these data layers. Habitat assessment is usually GIS-based; it includes the identification of species useful as indicators of ecological condition (see example by Franklin below) and the selection of data layers likely to be valuable in developing predictive models for the occurrence and distribution of individual species or species assemblages. The process of evaluating habitats has been considerably improved by the use of remotely sensed data in conjunction with other spatial information integrated inside a GIS framework. Enhancements in the management of forests also hinge on a deeper comprehension of the ecological processes involved in the cycles of carbon, nutrients, and water. Key inputs for models of carbon flux, nitrogen uptake and fertilization effect, and drought and water stress indicators come from remotely sensed data (Lucas and Curran 1999).

Wildlife Habitat Assessment

Remote sensing is essential for evaluating wildlife habitats. Critical information about vegetation kinds, forest structure, and changes throughout time is made possible by technology, and this information helps us understand the demands of wildlife habitats and how they respond to changes in their surroundings. This information is crucial for protecting biodiversity and keeping ecosystems robust and functional.

Hydrological and Watershed Analysis

In wooded environments, remote sensing provides vital tools for hydrological and watershed study. It makes surface water body

ISBN : 9788119821327

monitoring, water quality evaluation, and hydrological pattern prediction easier. In forest ecosystems, an understanding of these dynamics can aid in the effective and sustainable management of water supplies.

Geographic Information System (GIS) in Forestry:

Spatial Analysis: GIS assists in the analysis of spatial data related to forests, such as land cover types, topography, soil types, and proximity to water bodies, supporting decision-making processes. It integrates various layers of spatial data (forest boundaries, terrain, and soil types) to perform complex analyses for informed decision-making.

Planning for Forest Management: Geographic Information System (GIS) aids in the planning and efficient management of forest resources, such as wood harvesting, wildlife preservation, and recreation areas.

Habitat Modeling: Geographic Information System (GIS) makes it easier to identify

appropriate habitats for various species, supporting conservation efforts.

Monitoring and Reporting: GIS enables continuous monitoring of forest changes and provides a platform for reporting these changes to stakeholders.

Integration of Remote Sensing and GIS in Forestry:

Data Fusion: Combining RS data with GIS allows for better accuracy and more comprehensive analyses.

Decision Support Systems: Integrating RS and GIS creates decision support systems that aid in real-time decision-making for forest management and conservation.

Predictive Modeling: Predictive models can project future changes by merging GIS data with previous forest changes from RS, which supports proactive management measures. Fundamentally, RS and GIS are essential forestry tools because they offer precise, timely, and spatially explicit information that is necessary for conservation, sustainable forest management, and policy development.

ISBN : 9788119821327

Biodiversity Conservation: GIS allows the integration of data related to biodiversity, endangered species, and ecological corridors to make informed decisions for conservation efforts within forests.

Forest Inventory and Biomass Estimation: By combining data from remote sensing with measurements made on the ground, GIS makes it easier to create forest inventories by giving detailed information about the composition and structure of the forest. For the purpose of estimating biomass and taking stock of forests, remote sensing technology is essential. It contributes to the evaluation of forest resources by assisting in the estimation of tree heights, diameters, and volumes. Furthermore, the estimation of carbon stock a critical component of studies on climate change and the development of carbon sequestration programs depend on data from remote sensing systems.

Detailed forest inventory data

Forest inventory records are based mostly on stand boundaries generated from the manual interpretation of aerial images. Stand boundaries are vector-based representations of uniform forest feature units. A set of characteristics, usually comprising species composition, stand height, stand age, and crown closure, characterize these stand polygons. The inventory database can be updated with change (e.g., harvest) information for quality assurance, audit, and bias identification using digital remotely sensed data. Additionally, it can detect biases in the forest inventory databases caused by map sheet boundaries, vintage, or interpreter preferences, as well as add more attribute information. More thorough data on trees and stands must be gathered in order to manage forests sustainably for a variety of timber and non timber values.

Modeling and Prediction: GIS-based models are useful for simulating scenarios, forecasting forest development, and predicting the effects of many influences on forests, such as climate change and human involvement. The use of remote sensing data in GIS systems improves

ISBN : 9788119821327

the capacity for forestry analysis, visualization, and decision-making. Together, these technologies support conservation initiatives, sustainable forest management techniques, and our knowledge of the dynamics of forest ecosystems.

Spatial technologies and their applications

GIS applications in general

The applications of GIS, GPS, and RS technologies, either separately or in combination, cover a wide range of fields and complexity levels, according to Anon. Plotting maps for field usage, selecting sampling locations, and analyzing the distribution of soil types in connection to productivity and yields are a few examples of basic applications. Higher-level applications leverage the analytical powers of RS and GIS software. This can include vegetation classification for estimating crop productivity or environmental implications, modeling of surface water drainage patterns, or tracking animal movement patterns. The degree of integration between GIS applications and other forest management and financial systems determines the classification of these applications into different groups. Among these categories

- Data collection and maintenance.
- Map production.
- Data viewing and query.
- Decision support systems

GIS application in forest management

Map production:

Forest managers rely on a vast array of maps to successfully carry out their daily tasks. While these maps are commonly used to locate specific areas, such as plantations, they also provide a wealth of additional information. This can include important details such as roads, rivers, compartment boundaries, planted species, and compartment size. Furthermore, these maps may also feature other useful elements like neighboring areas, conservation zones, water

ISBN : 9788119821327

sources, fire breaks, infrastructure, and even topographical features like contours. These maps are essential tools that enable forest managers to make informed decisions and effectively manage their resources.

Fire management:

Another significant management challenge is the impact of fire on forest resources. Prescribed burning, wildlife control, ire prevention, and post-ire recovery measures are examples of management activities. In this situation, the modeling powers of GIS have shown to be highly useful. GIS has been utilized by forest fire managers for fuel mapping, meteorological condition mapping, and grading of fire danger. The vegetation cover, animals, plants, soil, streambed, air quality, microclimate, and even overall climate are all significantly impacted by forest fires. The devastation to property and lives as well as the loss of timber are readily apparent. Forest fires also result in the loss of the forest's recreational value and the devastation of wildlife habitat. The capacity for anger anticipation was essential for overseeing authorized burning actions In order to estimate the ire intensity based on variables such slope, height, site exposure, wind speed, relative humidity, cloud cover, temperature, and living and dead fuel moisture, described how ire behavior models were constructed using fuel models. Nonetheless, these models are not spatial and are usually applied to forecast fire behavior over a broad region. By using a raster-based GIS, it was possible to make the ire behavior models more sensitive to spatial diversity within the park. Several fuel and ire intensity models were then implemented using the GIS's mathematical modeling skills and a few lookup tables, along with input layers that were saved there.

Harvest planning:

Thorough planning of harvesting operations is necessary for good forest management practices. Identifying depots, extraction routes, felling orientations, and sensitive areas like wetlands are all part of the harvest planning process. A fundamental planning tool for these operations is a map. Maps are used by other tactical harvest planning

ISBN : 9788119821327

services to show scheduled felling over a period of years and to combine extraction routes and felling regions, allowing for the efficient use of resources and harvesting equipment.

Resource management:

Data collection for forest inventories and change monitoring are essential to forest management operations, according to Wulder and Franklin. However, a GIS can enhance these endeavors by integrating models to direct, say, timber harvesting, silviculture, and fire control operations, or forecast the availability of fuel wood and other resources. Priorities including protecting wildlife habitat, guaranteeing recreational opportunities, and reducing harvesting's visual effects are also becoming more and more important. Certain applications, like the production of timber, address a particular management issue, while others, like the production of timber along with habitat protection, show how a variety of management concerns can be linked through the use of GIS.

Uses of GIS in forest management:

• Where is it located? There are numerous methods to identify the location of forest resources on Earth, including place names, postal codes, zip codes, and geographic references like latitude and longitude.

• Status: Where is it located? a specific amount of non-forested land separated from a road or river.

• Trends: What's altered since then? Finding out what has changed over time in a studied forest or land use area is helpful.

• Patterns: Are there any spatial patterns? Check to see whether there is a landslide in the forest.

• Modeling: What if?

Determine what happens, if a road network is added in a forest For the purpose of delivering accurate and trustworthy information about natural resources, like forests, remote sensing (RS) is an indispensable instrument. A system for gathering, storing, analyzing, and managing data and related qualities that are spatially referred to the earth is

ISBN : 9788119821327

known as a geographic information system (GIS). To name a few uses, they can be applied to asset management, resource management, and scientific research. To aid in the management of forest resources, forest managers in Nigeria and around the world are increasingly turning to computerized forest inventories and geographic information systems. Several data sources, including field surveys, aerial photography, and satellite images, are employed to satisfy the various information needs for forest conservation, restoration, and management.

GIS and Remote Sensing Applications in Forestry

1. Mapping and Monitoring Forest Cover
2. Forest Fire Detection and Monitoring
3. Forest Inventory and Biomass Estimation
4. Forest Planning and Management
5. Deforestation Monitoring
6. Forest Health and Disease Monitoring
7. Wildlife Habitat Assessment
8. Watershed and Hydrological Analysis
9. Forest Conservation and Biodiversity Assessment

Forest managers have been using remote sensing more often, largely due to improved imaging integration with databases and GIS technologies, as well as technological advancements that better meet their information demands (Wulder and Franklin 2003). The most significant forest data gathered from remotely sensed imagery falls roughly into one of the following categories:

extensive monitoring of the health of the forest and natural disturbances; thorough data from the forest inventory, such as within-stand attributes;

The fundamentals of geographic information systems (GIS) and how they apply to the management of forestry; the use of GIS databases for forest inventory and management; the modeling of habitats and the

ISBN : 9788119821327

conservation of animals through GIS approaches; decision support system for GIS system.

Conclusion

Our understanding, management, and conservation of our forest resources are changing as a result of the use of remote sensing in forest management. Gathering data about things or regions from a distance is known as remote sensing, and it usually makes use of satellite or aerial sensors. Remote sensing is essential to forestry because it allows for the monitoring of forests, health assessments, detection of changes in land cover, and biomass estimation. Large-scale data collecting is made possible by it, and it facilitates the decision-making process for the conservation and management of forests. On the other hand, a geographic information system (GIS) collects, stores, processes, and displays spatial or geographic data. It combines several kinds of data layers onto one platform, including topography, vegetation, soil types, and land cover. GIS is helpful for forestry in. For example, satellite imagery collected through remote sensing can be processed and analyzed using GIS software to create detailed maps showing forest cover changes over time, assess the impact of wildfires or insect infestations, or plan for sustainable forest management practices. For studies on forest fires, an integrated analysis of geographical variables is highly beneficial. A source of vegetation data is provided via remote sensing, and the creation of fire hazard models is made possible by GIS processing. With the increasing complexity of forest management brought about by increased environmental and societal demands, GIS is probably going to become more and more important. More options for information access, even in more remote locations, will be made possible by advancements in wireless communication and wider band width. For the analytical resolution of competing suitabilitys and resource allocation decisions, more powerful techniques are required.

ISBN : 9788119821327

Applications for forestry represent the entire potential of GIS technology, in a sense.

Reference

1. Burrough, P. A. (1986). Principles of geographical. *Information systems for land resource assessment. Clarendon Press, Oxford*.
2. Chuvieco, E. and Congalton, R. G. (1989). Application of remote sensing and geographic information systems to forest fire hazard mapping. *Remote sensing of Environment*, **29**(2): 147-159.
3. Ezenwenyi, J. U. and Chukwu, O. (2020). Trends in Management of Tropical Forests: Application of Remote Sensing and Geographic Information System. In *Handbook of Research on the Conservation and Restoration of Tropical Dry Forests* (pp. 284-305). IGI Global.
4. Goward, S. N., Dye, D., Kerber, A. and Kalb, V. (1987). Comparison of North and South American biomes from AVHRR observations. *Geocarto International*, **2**(1): 27-39.
5. Lambin, E. F. (1996). Change detection at multiple temporal scales: seasonal and annual variations in landscape variables. *Photogrammetric engineering and remote sensing*, **62**(8): 931-938.
6. Lucas, N. S. and Curran, P. J. (1999). Forest ecosystem simulation modelling: the role of remote sensing. *Progress in Physical Geography*, **23**(3): 391-423.
7. Pomery, C. (2004). DNA and Family History: How Genetic Testing Can Advance Your Genealogical Research. Dundurn.
8. Rai, P. K. (2013). Forest and land use mapping using Remote Sensing and Geographical Information System: A case study on model system. http://www.taccire.sua.ac.tz/handle/123456789/164.
9. Sonti, S. H. (2015). Application of geographic information system (GIS) in forest management. *Journal of Geography & Natural Disasters*, **5**(3): 1000145.

ISBN : 9788119821327

10. Tudun-Wada, M. I., Tukur, Y. M., Hussaini, Y., Sani, M. Z., Musa, I. and Lekwot, V. E. (2014). Analysis of forest cover changes in Nimbia Forest Reserve, Kaduna State, Nigeria using geographic information system and remote sensing techniques. *International Journal of Environmental Monitoring and Analysis*, **2**(2): 91-99.
11. Vogt, D. J., Larson, B. C., Gordon, J. C. and Fanzeres, A. (Eds.). (1999). *Forest certification: Roots, issues, challenges, and benefits*. CRC Press.
12. Wulder, M. A. and Franklin, S. E. (2003). Remote sensing of forest environments, introduction: The transition from theory to information. *Remote sensing of forest environments: concepts and case studies*, pp: 3-12.

ISBN : 9788119821327

CHAPTER – 9

Application of RS, GIS and GPS in Fruit Science

Anju Yadav[1] Dhaneshra Kumari Gurjar[2]

[1]Ph.D. Scholar, Department of Fruit Science, College of Horticulture and Forestry, Jhalawar (Raj), 326023

[2]Ph.D. Scholar, Department of Fruit Science, College of Horticulture and Forestry, Jhalawar, Agriculture University, Kota (Raj), 326023

*Corresponding author: yadavaj109@gmail.com

Abstract

Fruit crops play a crucial role in enhancing land productivity, creating job opportunities, boosting exports, improving the economic well-being of farmers and entrepreneurs, and ensuring food and nutritional security. To effectively manage existing crops and expand fruit crop cultivation, having an updated and accurate database is essential for systematic planning and decision-making. Remote sensing (RS) emerges as a sophisticated tool that facilitates the collection and updating of information, aiding in the development of scientific management plans for fruit crops. The utilization of remote sensing, geographic information system and global positioning system is rapidly demonstrating its potential in various applications, such as detecting crop biomass, assessing soil properties, monitoring soil moisture and nutrient content, counting green fruits, estimating crop yields, and identifying damage caused by biotic and abiotic stresses. These technologies serve as reliable sources of information for the identification, classification, mapping, monitoring, and planning of natural resources, as well as for disaster mitigation and management. They hold great promise for site-specific management, precision horticulture, market planning, and export activities. In numerous instances, RS, GIS and GPS have proven to be influential tools for environmental monitoring.

ISBN : 9788119821327

Keywords: Remote sensing, Geographic Information Systems, Global Positioning System, Fruit crops, database

Introduction

Modern horticulture has witnessed a significant transformation with the integration of advanced technologies such as Remote Sensing (RS), Geographic Information Systems (GIS), and Global Positioning System (GPS). Remote Sensing plays a pivotal role in fruit crop monitoring by providing valuable information on crop health, stress detection, and yield estimation. Satellite and aerial imagery, equipped with various spectral bands, enable precise and timely assessment of vegetation indices, helping farmers make informed decisions regarding irrigation, fertilization, and pest control. Geographic information systems serve as a powerful tool for spatial analysis and decision-making in fruit crop management. GIS facilitates the integration of diverse data layers, such as soil types, weather patterns, and topography, offering a holistic view of the orchard environment. Through spatial modeling and mapping, GIS aids in optimizing resource utilization, predicting disease spread, and designing efficient orchard layouts. Global positioning system technology enhances precision agriculture practices in fruit cultivation. GPS-enabled devices provide accurate location data, allowing farmers to precisely navigate through orchards during various field operations. This technology is crucial for implementing variable rate applications, optimizing planting patterns, and automating machinery, leading to improved efficiency and reduced environmental impact.

The integration of RS, GIS, and GPS in fruit crop management fosters a data-driven approach, enhancing the overall sustainability and productivity of orchard operations. The successful implementations of these technologies, demonstrating their real-world impact on fruit crop cultivation. Additionally, challenges and potential future advancements in this integration are discussed, aiming to guide researchers,

ISBN : 9788119821327

practitioners, and policymakers toward the continued development and adoption of these technologies in fruit farming.

There has been a slow evolution of information technologies in fruit crop production as innovators combined their prior field experience with an attitude of learning by doing while looking for practical applications for farm management. The application of aerial or satellite imaging, along with Global polar satellite (GPS) and (GIS) Geographical information system, is the first step towards the successful application of Site-specific crop management (SSCM) in fruit crops. The use of Site-specific crop management (SSCM) for horticultural crops such as fruits has potential for increasing net returns and optimizing resource use. The delineation of orchards and spatial analysis using geospatial technology can provide additional information for management decision making, such as the determination of fruit yield, the quantification and scheduling of precise and proper fertilizer, irrigation needs, and the application of pesticides for pest and disease management.

Remote Sensing

Remote sensing is the process of acquiring data or information about objects or substances not in direct contact with the sensor by gathering its inputs using electromagnetic radiation or acoustical waves that emanate from the targets of interest. This involves sensing and recording reflected or emitted energy and processing, analyzing and applying that information.

Types of Sensors used in Remote Sensing Sensor:

These are the devices that receive electromagnetic radiation and convert it into signal that can be recorded or displayed as either numerical data or an image. There are many types of sensors used in fruit crops 1) Microwave radiometer 2) Gravimeter 3) Spectrometer 4) Camera 5) Solid scanner 6) Optical mechanical scanner 7) Laser water depth meter 8) Laser distance meter 9) Radar 10) Satellites also act as sensors. In thermal remote sensing the invisible radiation patterns of

ISBN : 9788119821327

objects are converted into visible images and these images are called thermo grams or thermal images. Thermal images can be acquired using portable, hand-held or thermal sensors that are coupled with optical systems mounted on an airplane or satellite. This technology is a non-invasive, non-contact and nondestructive technique used to determine thermal properties and features of any object of interest and therefore it can be used in many fields, where heat is generated or lost in space and time.

Platforms for Remote Sensing System

Platforms are nothing but vehicles or carriers of remote sensing. Mainly there are three types of platforms used in remote sensing.

Ground Based: These include the use of vehicles, portable masts and towers. These are mounted with cameras, and sensors at different heights.

Airborne: It includes the use of aeroplanes, helicopters and balloons. It is not cost effective one and occupies only small areas.

Space Borne: In space-borne remote sensing, sensors are mounted onboard a spacecraft (space shuttle or satellite) orbiting the earth. Space-borne or satellite platform are onetime cost effected but relatively lower cost per unit area of coverage. It can acquire imagery of entire earth without taking permission. Remote sensing process mainly involves the use of visible to the microwave region. As the atmosphere appears to be transparent in the visible region most of the satellite remote sensing studies are designed in visible region. Sensor can detect the one object from the other by the means of the electromagnetic radiation absorbed.

Principles and Process of Remote Sensing System

In much of remote sensing, the process involves an interaction between incident radiation and the targets of interest.

ISBN : 9788119821327

i) **Energy Source or Illumination:** The first requirement for remote sensing is to have an energy source which illuminates or provides electromagnetic energy to the target of interest. Example: sun act as source of energy.

ii) **Radiation and the Atmosphere**: As the energy travels from its source to the target, it will come in contact, interact and passes through the atmosphere. This interaction may take place a second time as the energy travels from the target to the sensor.

iii) **Interaction with the Target**: Once the energy makes its way to the target through the atmosphere it interacts with the target depending on the properties of both the target and the radiation.

iv) **Recording of Energy by the Sensor:** After the energy has been scattered back or emitted from the target. It requires a sensor (remote - not in contact with the target) to collect and record the electromagnetic radiation.

v) **Transmission, Reception, and Processing:** The energy recorded by the sensor has to be transmitted often in electronic form to a receiving and processing station at which the data was processed.

vi) **Interpretation and Analysis:** The processed image is interpreted, visually and/or digitally or electronically to extract information about the target, which was illuminated.

vii) **Application**: The final step of remote sensing process is applying the information extracted from image about the target in order to find the solution to particular problem.

There are two basic types of sensors:

1) Passive sensors

2) Active sensors

Passive Sensors: It record radiation reflected from the earth's surface. The source of this radiation must come from outside the sensor. In most of the cases, it is solar energy. Passive solar sensors can only capture data during daylight hours. The thematic mapper (TM) sensor system on the landsat satellite is a passive sensor.

ISBN : 9788119821327

Active Sensors: These are different from passive sensors. Unlike passive sensors, active sensors require the energy source to come from within the sensor. For example, a laser-beam remote sensing system is an active sensor that sends out a beam of light with a known wavelength and frequency. This beam of light hits the earth and is reflected back to the sensor. It records the time it took for the beam of light to return. Topographic LIDAR laser beach mapping data included on this CD-ROM were collected with an active sensor.

Types of Remote Sensing System

1. Visual Remote Sensing System

The best example of a Visual remote sensing system in human beings is human visual system in general sense. The sensors in human visual system have two types of photosensitive cells, known as the cones and rods in retina of the eyes. There are three types of cones each being sensitive to one of the red, green, and blue regions of the visible spectrum. The rods are sensitive only to the total light intensity. Hence, everything appears in shades of grey when there is insufficient light. As the objects events being observed are located far away from the eyes the information needs a carrier to travel from the object to the eyes. In this case, the information carrier is the visible light, a part of the electromagnetic spectrum. The part of the scattered light is intercepted by the eyes, forming an image on the retina after passing through the optical system of the eyes. The signals generated at the retina are carried through the nerve fibres to the brain i.e. the central processing unit (CPU) of the visual system.

2. Optical Remote Sensing System

In optical remote sensing optical sensors detect solar radiation reflected or scattered from the earth form images resembling photographs taken by a camera high up in space. The wavelength region usually extends from the visible and near infrared VNIR to the short-wave infrared SWIR. Different materials such as water, soil, vegetation, buildings and roads reflect visible and infrared light in

ISBN : 9788119821327

different ways. They have different colours and brightness under the sun. The interpretation of optical images requires the knowledge of the spectral reflectance signatures of the various materials (natural or man-made) covering the surface of the earth.

3. Infrared Remote Sensing System

Infrared remote sensing makes use of infrared sensors to detect infrared radiation emitted from the earth's surface. The middle-wave infrared (MWIR) and long-wave infrared (LWIR) are within the thermal infrared region. These radiations are emitted from warm objects such as the earth's surface. They are used in satellite remote sensing for measurements of the earth's land and sea surface temperature. Thermal infrared remote sensing is also often used for detection of forest fires, volcanoes and oil fires.

4. Microwave Remote Sensing System

There are some remote sensing satellites which carry passive or active microwave sensors. The active sensors emit pulses of microwave radiation to illuminate the areas to be imaged. The images of the earth surface are formed by measuring the microwave energy scattered by the ground or sea back to the sensors. These satellites carry their own "flashlight" emitting microwaves to illuminate their targets. The images can thus be acquired day and night. Microwaves have an additional advantage as they can penetrate clouds. The mages acquired even when the clouds covering the earth surface. A microwave imaging system which can produce high resolution image of the earth is the synthetic aperture radar (SAR).

5. **Satellite Remote Sensing System:**

In this, many remote sensing images are acquired by earth observation satellites. These remote sensing satellites are equipped with sensors looking down to the earth. They are the eyes in the sky constantly observing the earth as they go round in predictable orbits. The orbital platforms collect and transmit data from different parts of the electromagnetic spectrum, which in conjunction with larger scale

ISBN : 9788119821327

aerial or ground-based sensing and analysis provides researchers with enough information to monitor trends. The other uses include different areas of the earth sciences such as natural resource management, agricultural fields such as land usage and conservation and national security and overhead, ground-based and stand-off collection on border areas.

6. **Airborne Remote Sensing System**

In airborne remote sensing downward or sideward looking sensors are mounted on an aircraft to obtain images of the earth's surface. An advantage of airborne remote sensing, compared to satellite remote sensing, is the capability of offering very high spatial resolution images (20 cm or less). The disadvantages are low coverage area and high cost per unit area of ground coverage. It is not cost-effective to map a large area using an airborne remote sensing system. Airborne remote sensing missions are often carried out as one-time operations, whereas earth observation satellites offer the possibility of continuous monitoring of the earth.

Geographic Information Systems (GIS)

Geographic information systems are computer-based systems that deal with virtually any type of information about features that can be referenced by geographical location. These systems are capable of handling both locational data and attribute data about the features.

Components of GIS

GIS comprises of the following six components

1. Hardware: It refers to powerful computers, printers, output devices, such as monitors, input devices, and large amounts of disk space to store the large GIS data sets.

2. Software: It refers to the free and commercial software packages which allows GIS users to perform GIS functions such as the manipulation, storage, query and analysis of spatial and non-spatial information

ISBN : 9788119821327

3. Data: It refers to all spatial and non-spatial information stored in a digital format on the computer. A GIS can visualize the data as a map, graph or other graphic, which helps the user to understand and choose through massive amounts of data.

4. Methods: It includes the formulas, statistics, analysis and algorithms that are used to turn data into information, which aids easy interpretation of the information.

5. People: It refers to the users of GIS. This includes both the general public as well as the skilled GIS professionals.

6. Network: It refers to both the computer and social network.

Global Positioning System (GPS)

GPS (Global Positioning System) is a satellite-based navigation system that enables farmers to map and measure their fields with high precision. This data can be used to create detailed field maps, which can help farmers to identify variations in soil type, moisture content, and other factors that can affect crop growth and yield. GPS can also be used to guide precision equipment such as tractors, sprayers, and harvesters, enabling farmers to apply inputs at precise locations in the field.

Application of RS, GIS and GPS in Fruit Crops

Research on remote sensing in fruit crops through remotely piloted aircraft systems (RPAS), commonly known as drones, has gained significant traction. These devices are increasingly utilized for monitoring horticultural crops, particularly fruit crops. By analyzing images captured by RPAS, various plant physical characteristics, including water content, cellular structure, and plant pigment such as chlorophyll, can be derived.

Chlorophyll content, a crucial parameter for assessing tree vitality, is closely linked to essential plant processes such as photosynthetic potential, nutrient content, plant stress, maturity, and senescence. The machine learning algorithm Random Forest demonstrated the highest performance in a chlorophyll extraction

ISBN : 9788119821327

model. Applying this model to RPAS images allows for the creation of chlorophyll maps throughout the growing season in fruit orchards. Chlorophyll's significance in determining orchard productivity is evident. The relationship between canopy chlorophyll content and corresponding harvest indicators indicates that regions with low canopy chlorophyll content early in the growing season and high chlorophyll content later on are likely to experience reduced productivity.

Remote sensing in estimating water requirements for crops: The widely adopted method for this purpose is the FAO approach, which considers climatic variables within reference evapotranspiration and the crop type characterized by the crop coefficient (Kc). In citrus trees, the crop coefficient is primarily influenced by tree ground covers (GC). In extensive areas, the estimation of tree ground covers (GC) can be achieved through remote sensing tools. Once tree water requirements are determined, this information can be integrated into geographic information systems. Accurate fruit crop yield estimation, with a focus on tree variability, is essential for precision orchard management. A burgeoning research trend involves the application of spectroscopic analysis and image sensing technology to estimate site-specific fruit yield. This advancement contributes to the evolving landscape of research in precision agriculture.

Vegetation indices: serve as remote sensing indicators, utilizing reflectance values at various wavelengths incorporated into equations to derive specific parameters. Different indices, such as the Normalized Difference Vegetation Index (NDVI) for vegetation cover and the Structure Insensitive Pigment Index (SIPI) for stress, are employed as indicators for various parameters.

Precision fertilization in citrus orchards: In citrus plants Zaman *et al*. (2005) investigated the application of variable-rate nitrogen in Florida, employing ultrasonically-sensed tree size for precision. Real-time measurements of tree canopy sizes were conducted in a standard

ISBN : 9788119821327

17 ha Valencia grove, utilizing an automated ultrasonic sensor system equipped with Differential Global Positioning System (DGPS). Prescription maps for variable nitrogen fertilizer application were created based on ultrasonically scanned tree sizes at an individual tree level, utilizing Arc View GIS and Mid tech Field ware. Leaf samples from trees with diverse canopy sizes, conventionally fertilized at a uniform rate of 270 kg N/ha/y, underwent analysis for nitrogen concentration. An analysis of 2980 tree spaces in the grove revealed a skewed size distribution, with 62% falling in the 0- to 100m^3/tree volume classes and a median volume of 79 m^3/tree. The tree volumes ranged from 0 to 240 m^3/tree. Regression analysis indicated that trees with excess leaf nitrogen (>3%) typically had canopies less than 100 m^3, potentially leading to lower fruit yields and quality. Furthermore, wasted fertilizer nitrates could leach beyond the root zone to groundwater.

To address excess fertilization in smaller trees, a granular fertilizer spreader with hydraulically powered split-chain outputs, controlled by a Mid Tech Legacy 6000 controller, was employed for variable-rate nitrogen application in half of the grove. This approach achieved a 38% to 40% reduction in granular fertilizer cost, implementing variable N rates on a per-tree basis ranging from 135 to 270 kg N/ha/y. The use of remote sensing to assess canopy nitrogen status through the canopy chlorophyll concentration index (CCCI) for fruit crops like apple and pear to remove the influence of the orchard under storey, test datasets of high spatial resolution imagery.

Detection of **soil salinity using thermal imaging**

The utilization of thermal imaging for detecting soil salinity is essential due to the significant environmental degradation caused by high soil salinity levels, hindering crop growth and overall regional production. Thermal imageries can serve as a valuable source for extracting soil salinity information, as the emitted radiance provides insights into subsoil conditions.

ISBN : 9788119821327

Nursery monitoring in fruit crops through thermal imaging

Nurseries are located where plants are propagated and cultivated to reach a usable size, utilizing open fields, container fields, tunnels, or greenhouses. Early identification of seed quality and monitoring seeds is crucial, necessitating a tool that can distinguish between viable and nonviable seeds before utilization. This tool not only aids in recognizing individual seeds' viability but also allows for timely adjustments in cultural practices to prevent stress from becoming well established.

Researchers now acknowledge that the physiological attributes of seeds are not visually discernible. Consequently, thermal infrared imaging techniques have been proposed as a potential alternative to traditional methods for assessing seed viability. The thermal profile of seeds becomes instrumental in detecting subtle temperature changes, which vary with viability. This capability facilitates the classification and separation of viable seeds from non-viable ones.

Spectral characteristics of fruit crops

Utilizing images from the visible and near-infrared (NIR) bands within the electromagnetic spectrum is crucial. Consequently, any multispectral image incorporating NIR bands can effectively distinguish fruit crops from other land covers.

The spectral signatures of Lingon blueberry and bearberry plants indicate that their shrubs exhibit very low reflectance compared to grasses and forest trees. This is attributed to their medium stature and their ability to blend easily with surrounding shadows. Blueberry plants are distinguishable in the 680–700 nm (red) band width and the 800-900 nm (NIR) band width range. Red blush citrus is discernible in the 600-650 nm (green) band width, demonstrating high absorption in the 950-1,000 nm (NIR) band width range with a steep absorption rate. Additionally, it exhibits the highest reflection rate in the 1,000-1,150 nm (NIR) band width range (Panda *et al.,* 2010).

ISBN : 9788119821327

Exploring the potential of high resolution worldview- 3 imagery for estimating yield of mango

The use of high-resolution WorldView-3 (WV3) satellite imagery for mango yield estimation involves integrating both geometric (tree crown area) and optical (spectral vegetation indices) data through an artificial neural network (ANN) model. The stratified sampling technique (SST) was employed to select eighteen trees from each orchard, and these trees were subsequently ground-truthed for yield (kg/tree) and fruit number per tree. From the WV3 imagery, spectral reflectance data and tree crown area (TCA) were extracted for each sampled tree.

Upon combining data from all trees across three orchards in two growing seasons, the trunk cross-sectional area (TCA) emerged as the most crucial predictor for both fruit yield (kg/tree) and fruit number. Following TCA, the normalized difference vegetation index (NDVI) and the red-edge band were identified as significant contributors (Rahman *et al*., 2018).

Assessment of Quality in Fruits Crops

Grape wine quality was evaluated based on vine vigor, a parameter directly linked (Johnson *et al.,* 2001). Utilizing remotely sensed vegetation index imagery, sub-block management zones were established in a three-hectare commercial chardonnay wine-grape vineyard. Ground-based measurements later confirmed distinct differences between low and high-vigor zones concerning biomass (mainly shoot vigor), vine water status, and crucially, fruit and wine characteristics. Harvesting based on vigor zones allowed the extraction of unique wine lots from a block traditionally treated as a single unit, providing winemakers with enhanced flexibility in the final blending process.

ISBN : 9788119821327

Identification of pest and diseases in fruit crops

The use of hyper spectral remote sensing has shown potential in distinguishing between healthy and malformed mango panicles based on spectral reflectance data. Various indices derived from hyper spectral reflectance demonstrated promise, with the red edge technique being the most effective for discrimination. Malformed panicles were characterized by higher red edge values and red edge 'position' (red shift), which contrasts with the general trend of a blue shift in REP and lower REV indicating diseased or stressed plants. Additionally, hyper-indices like the normalized difference vegetation index (NDVI) and structure-insensitive pigment index (SIPI) proved useful in differentiating malformed panicles from healthy ones (Nagaraja *et al*., 2014).

Remote sensing techniques can reduce pest monitoring costs in orchards. Eike *et al*. (2009) evaluated the feasibility of detecting spider mite damage in orchards by measuring visible and near infrared reflectance of 1153 leaves and 392 canopies in 11 peach orchards in California. Pairs of significant wavelengths, identified by Partial Least Squares regression, were combined into normalized difference indices. These indices were evaluated for correlation with mite damage. Eight spectral regions for leaves and two regions for canopies (at blue and red wavelengths) were significantly correlated with mite damage. Index values were linearly correlated with mite damage (R2=0.47), allowing identification of mite hotspots in orchards. However, better standardization of aerial imagery and accounting for perturbing environmental factors will be necessary for making this technique applicable for early mite detection.

Physiological stress in trees causes changes in photosynthetic pigments such as chlorophyll, carotenoids and other factors which can easily be detected through changes in spectral reflectance. With improvements in spatial, spectral and temporal resolution of remote sensing, multispectral imagery remains advantageous due to its real-

ISBN : 9788119821327

time or near real-time imagery for visual assessment. With this imagery technique it is possible to detect infected trees and healthy trees. Sindhuja *et al.,* (2013) reported that there was a difference in average reflectance. Values of the healthy trees in the visible region were lower than those in the near infrared region, while the opposite was the case for HLB-infected trees. This could help in taking up of immediate control measures, so that spread of disease could be minimized.

Maturity Evaluation in fruit crops

An investigation focused on employing thermographs to assess mechanical damage, bruising, and maturation levels in apples, specifically the Jonagold and Cox varieties. Thermal imaging proves effective in estimating fruit maturity, even enabling the differentiation of various varieties at the same ripeness stage. Consequently, thermal imaging systems are well-suited for scrutinizing the maturity and quality of produce. In-depth thermal analyses of the transpiration behavior of the fruits and its interaction with the environment enable the identification of the optimal harvest date and the customized design of postharvest machinery.

Detection of bruises using thermal imaging

It has become a viable solution for identifying the most common types of damage, such as bruises and scratches, on the surfaces of fruits and vegetables during transportation and handling. Bruising is characterized by the damage to fruit tissue caused by external forces, resulting in physical texture changes or chemical alterations in color, smell, and taste. The advancement in imaging technology has made it possible to perform online detection of bruises on fruits.

Thermal imaging presents a promising alternative for bruise detection. The initial study on bruise detection using thermal imaging focused on apple and Satsuma mandarin fruits. Artificial bruises were induced by pressing and scratching, and thermal images were captured

ISBN : 9788119821327

with a thermal camera operating within the wavelength range of 8-14μm. The temperature difference between bruised and intact skin typically fell within the range of 0.2 ^{o}C to 1.0 ^{o}C, with a measuring duration of at least 10 minutes. Notably, the temperature at the bruised areas was lower than that of the unbruised skin.

Mapping of orchards and estimation of cultivable land area

In the Pulwama district of Kashmir valley, Mushtaq and Asima (2014) utilized remote sensing and agro-meteorological land-based observations to estimate the cultivatable land area of apple orchards. They considered various parameters, including terrain features such as elevation, slope, and aspect, for a more comprehensive characterization of apple orchards. Landsat and AWIFS digital data were employed to monitor and estimate the acreage under apple orchards. The majority of apple orchards (89.82%) were concentrated in the elevation range of 1500-2000 m, with no orchards below 1500m and 10% above 2000m. Dense orchards were found in the elevation range of 1500-2000 meters. In another study by Sharma and Panigrahi (2010) in the Shimla district of Himachal Pradesh, high-resolution remote sensing data from the Indian Remote Sensing (IRS) satellite P6 was utilized to map apple orchards and generate a block-wise database on apple plantations. The research achieved over ninety percent accuracy using remote sensing data. Key findings indicated that Shimla is a major apple-growing block with 376.30 sq. km. under apple orchards. The elevation range of 2000-3000m and a slope of 21-40° predominantly supported orchards in Shimla district. In a recent study by Verma *et al*. (2020) in Lucknow district, multi-temporal Landsat 8 Operational Land Imager (OLI) images from 2015 to 2017 were used to classify mango orchards. The mango orchard area was estimated, revealing an increase in acreage from 2015 (2807 ha) to 2017 (2966 ha).

ISBN : 9788119821327

Assessment of Land Suitability for Pomegranate Cultivation: In the Sangola tahsil of Solapur district, Maharashtra's south-western region, Yegade *et al*. (2013) conducted a study to evaluate the appropriateness of the land for pomegranate production. Utilizing multi criteria logic and GIS applications, the researchers assessed various factors, including slope, soil drainage, soil texture, soil depth, soil type, groundwater availability, and the existing land-use in the study area. The results of the land evaluation indicated that 4986 ha of the total pomegranate crop area in the study region were situated in highly suitable areas, with an additional 3255 ha falling within moderately suitable areas. A substantial portion of 5596 ha was classified as marginally suitable areas. This study provides valuable local-level insights that farmers can leverage to make informed decisions about selecting optimal cropping patterns based on the suitability of the land.

Orchard crop inventory: In a study conducted by Chaudhari *et al.* (2019), a research methodology was developed to assess the area of three significant fruit crops – Banana, Mango, and Citrus across 20 districts in Gujarat, Madhya Pradesh, Uttar Pradesh, and Bihar. The researchers selected an appropriate bio-window for analyzing various crop types and employed pixel-based hybrid classification methods, combining unsupervised ISODATA clustering and supervised MXL classification. Additionally, object-based classification of high-resolution remote sensing data from Resourcesat LISS III and/or LISS IV, Cartosat – 1 PAN was performed. The study included accuracy assessment and a comparison with departmental reported statistics, revealing an overall classification accuracy exceeding 80% for all the crops. This research demonstrated the effectiveness of satellite remote sensing data in estimating the crop area of major horticultural crops.

Detection of water stress: Abiotic stress such as deficiency of moisture can also be identified through remote sensing technology. Several techniques such as high spatial resolution multispectral and thermal airborne imagery were employed to monitor crown

ISBN : 9788119821327

temperature and the Photochemical Reflectance Index (PRI) in peach orchards. Different irrigation regimes included sustained and regulated deficit irrigation strategies, were utilized to induce stress. The results depict that a difference in reflectance pattern between well irrigated and stressed plants can be observed (Saurez *et al*., 2010).

High resolution remote sensing imagery in fruit crops

Crop yield is perhaps the most important piece of information for crop management in precision farming. It integrates the effects of various spatial variables such as soil properties, topography, plant population, fertigation, irrigation, and pest infestations. A yield map can therefore be an indispensable input for site-specific operations either by itself or in combination with other spatial Information. The yield monitor data can only be used for after-season management, whereas some problems such as nutrient deficiencies, water stress, or pest infestations should be managed during the growing season. Remote sensing imagery obtained during the growing season has the potential not only for after-season management, but also for within-season management. Additionally, yield maps derived from remote sensing imagery can be used as an alternative when yield monitor data are not available. Traditional satellite imagery has been used for yield estimation over large geographic areas but this type of imagery has limited use for assessing the variation in yield within fields because of its coarse spatial resolution. Therefore, airborne multispectral and hyper spectral imagery and high resolution satellite imagery have been used for mapping within-field yield variability and other precision agriculture and horticulture applications. Airborne multispectral imaging systems provide image data at fine spatial resolutions and at narrow spectral bands and have the real-time monitoring capability (Lee *et al*., 2010). The potential of the narrow-band TBVI (two-band vegetation index) derived from airborne hyper spectral imagery to predict citrus fruit yield was examined in Japan (Ye *et al.,* 2008). The

ISBN : 9788119821327

microwave backscatter response of pecan tree canopy samples to estimate pecan yield in situ using terrestrial radar was studied (James *et al*., 2013).

Estimation of cultivatable land area in apple fruit crop

Apple orchard cultivatable land area was estimated using remote sensing and agro-metrology land based observation in Pulwama district of Kashmir valley Mushtaq and Asima (2014). They have taken different parameters viz., terrain parameters like elevation slope and aspect facilitates better characterization of apple orchards. These parameters can be further utilized for apple orchard area expansion. Landsat and AWIFS digital data were used to monitor and estimate acreage under apple orchards.

Application of RS, GIS and GPS after Harvest in Fruit Crops

1. Employing Vibration Response Technique for Assessing Pear Fruit Firmness during Storage

The laser doppler vibrometer (LDV) technology was employed to estimate the texture firmness of pears. This technique involved detecting responses to imposed vibrations on intact fruit using a shaker. The vibrations transmitted through the fruit to the upper surface were measured with a laser doppler vibrometer. The response signals were processed using a fast Fourier transform algorithm, extracting the desired results for firmness evaluation.

2. Utilizing a Laser Doppler Vibrometer for Remote Sensing of Fruit Textural Changes

Changes in fruit texture during ripening were remotely sensed by assessing surface vibrations. The method was tested on fruits exhibiting a variety of firmness and textural characteristics, including ethylene-treated kiwifruit, apples stored at 10 or 20 °C, and persimmons stored at 10 °C. In each instance, fruits were positioned on a stage capable of generating sine wave vibrations ranging from 5 to 2,000 Hz. The

ISBN : 9788119821327

vibration transmitted through the fruit to the upper surface was accurately measured without direct contact using a Doppler laser vibrometer. To obtain the genuine vibrational signal of the fruit mass, the perceived fruit surface signal was corrected by subtracting the stage vibration, determined based on an accelerometer signal (Muramatsu *et al*., 1999).Application of Vibration Response Technique for the Firmness Evaluation of Pear Fruit during Storage.

References

a. 269–275. https://doi.org/10.5194/ISPRS-ARCHIVES-

b. 269–275. https://doi.org/10.5194/ISPRS-ARCHIVES-

c. 269–275. https://doi.org/10.5194/ISPRS-ARCHIVES-

d. 269–275. https://doi.org/10.5194/ISPRS-ARCHIVES-

e. Chaudhari K., N. Nishant, G. Upadhyay, R. More, N.

f. Chaudhari K., N. Nishant, G. Upadhyay, R. More, N.

g. Chaudhari K., N. Nishant, G. Upadhyay, R. More, N.

h. Chaudhari K., N. Nishant, G. Upadhyay, R. More, N.

i. Chaudhari, K.N., Nishant, G., Upadhyay, R., More, N., Singh, S.P.,

j. Vyas and Bhattacharya, B.K. (2019). Crop inventory of orchard crops in India using remotely sensed data. International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences-ISPRS Archives, 42(3/W6), pp.269-275.

k. Eike, L., Adam, H., Minghua, Z., Walter, J.B. and Cecil, D. (2009). Remote sensing of spider mite damage in California peach orchards. *Int. J App. Earth Observation and Geoinformation*, 11(3):244-255.

l. Information Sciences - ISPRS Archives42(3/W6), pp.

m. Information Sciences - ISPRS Archives42(3/W6), pp.

n. Information Sciences - ISPRS Archives42(3/W6), pp.

o. Information Sciences - ISPRS Archives42(3/W6), pp.

ISBN : 9788119821327

p. inventory of orchard crops in India using remotely
q. inventory of orchard crops in India using remotely
r. inventory of orchard crops in India using remotely
s. inventory of orchard crops in India using remotely

2. James, A., Hardin, P.R. and Weckler, C.L. (2013). Microwave backscatter response of pecan tree canopy samples for estimation of pecan yield in situ using terrestrial radar. *Computers and Electronics in Agriculture,* 90:54-62.
3. Johnson. L.F., Bosch, D.F., Wiiliums, D.C. and Lobitz, B.M. (2001). Remote sensing of vineyard management zones: Implications for wine quality. *Applied Engineering in Agriculture*, 17(4):557-560.
4. Lee, W.S., Alchanatis, V., Yang, C., Hirafuji, M., Moshouen, D. and Lif, C. (2010). Sensing technologies for precision specialty crop production. *Computers and Electronics in Agriculture*, 74:2-33.
5. Muramatsu, N., Sakurai, N., Wada, N., Yamamoto, R., Keiichi, T. and Toshikazu A. (1999). Remote sensing of fruit textural changes with a laser doppler vibrometer. *Journal of the American society for horticultural science*, 125(1):120-127.
6. Mushtaq, G. and Asima, N. (2014). Estimation of apple orchard using remote sensing and agro-meteorology land based observation in Pulwama district of Kashmir valley. *International Journal of Remote sensing and Geosciences*, 3(6):2319-3484.
7. Nagaraja, A., Sahoo, R.N., Usha, K., Singh, S.K., Sivaramanae, N. and Gupta, V.K. (2014). Spectral discrimination of healthy and malformed mango panicles using spectroradiometer. *Indian Journal of Horticulture*, 71(1):40-44.
8. Panda, S.S., Gerrit, H. and Joel, O.P. (2010). Remote sensing and geospatial technological applications for site-specific management of fruit and nut crops: A review. *Remote sensing,* 2:1973-1997.

a. Photogrammetry, Remote Sensing and Spatial
b. Photogrammetry, Remote Sensing and Spatial

ISBN : 9788119821327

c. Photogrammetry, Remote Sensing and Spatial
d. Photogrammetry, Remote Sensing and Spatial

9. Rahman, M.M., Andrew, R. and Mila, B. (2018). Exploring the potential of high resolution worldview-3 imagery for estimating yield of mango. *Remote sensing*, 10:1-18.
    a. sensed data, International Archives of the
    b. sensed data, International Archives of the
    c. sensed data, International Archives of the
    d. sensed data, International Archives of the
10. Sharma, A. and Panigrahy, S. (2010). Apple orchard characterization using remote sensing and GIS in Shimla district of Himachal Pradesh. National Horticulture Technology Mission Programme.
    a. Singh, S. P. Vyas and B. K. Bhattacharya (2019). Crop
    b. Singh, S. P. Vyas and B. K. Bhattacharya (2019). Crop
    c. Singh, S. P. Vyas and B. K. Bhattacharya (2019). Crop
    d. Singh, S. P. Vyas and B. K. Bhattacharya (2019). Crop
    e. XLII-3-W6-269-2019
    f. XLII-3-W6-269-2019
    g. XLII-3-W6-269-2019
    h. XLII-3-W6-269-2019
11. Sindhuja, S., Joe, M., Sherrie, B. and Reza, E. (2013). Huanglongbing (Citrus Greening) detection using visible, near infrared and thermal imaging techniques. *Sensors,* 13(6):2117-2130.
12. Suarez, L., Zarco, T.P.J., Gonzalez, V., Berni, J.A., Sagardoy, R. and Morales, B. (2010). Detecting water stress effects on fruit quality in orchards with time-series PRI airborne imagery. *Remote Sensing of Environment,* 114(4):286-298.
13. Verma, H.C., Rajan, S. and Ahmed, T. (2020). Mapping and area estimation of mango orchards of Lucknow region by applying knowledge based decision tree to Landsat 8 OLI satellite images.

ISBN : 9788119821327

International Journal of Innovative Technology and Exploring Engineering (IJITEE), 9(3):2278-3075.

14. Ye, X., Sakai, K. and Asada, S. (2008). Application of narrow-band two-band vegetation index in estimating fruit yield in citrus. *Biosystems Engineering*, 99:179-189.
15. Yedage, A.S., Gavali, R.S. and Jarag, A.P. (2013). Land assessment for horticulture (pomegranate) crop using GIS and fuzzy decision analysis in the Sangola taluka of Solapur district. International *Journal of Remote Sensing and GIS*, 2(3):104-113.
16. Zaman, Q.U., Schumann, A.W. and Miller, W.M. (2005). Variable rate nitrogen application in florida citrus based on ultrasonically-sensed tree size. *American Society of Agricultural Engineers*, 21(3):331-335.

ISBN : 9788119821327

CHAPTER – 10

Application of RS, GIS and GPS in Plant Pathology

Ravichandra[1*], Sudarshan Varma[2] and Channabasava[3]

[1]*Department of Plant Pathology, CoA, KSNUAHS, Shivamogga, Karnataka, India*

[2]*Department of Soil Science, CoA, KSNUAHS, Shivamogga, Karnataka, India*

[3]*Department of Genetics and Plant Breeding, CoA, UAS Raichur, Karnataka, India*

[*]Corresponding Author: ravichandragk13@gmail.com

Abstract

Agriculture is the basis of every economy worldwide. Crop production is one of the major factors affecting domestic market and is also a major prerequisite of economic development in any country. A lot of issues are responsible for estimated crop production varying in different parts of the world. Among them plant diseases acts as a major threat to farmers as they lead to major destruction in overall food supply. To provide effective measures for detection and avoidance of the destruction requires an early identification of type of plant disease present. As conventional disease diagnostic methods are time consuming, demand trained experts there is need for new, automated sensor-based methods for detection and quantification of plant diseases. RS, GIS and GPS These sensors assess the optical properties of plants within different regions of the electromagnetic spectrum and are able to utilize information beyond the visible range thus allow us to quantify disease symptoms that are invisible to naked eye. They enable the detection of early changes in plant physiology due to biotic stresses, because it can modifies the transpiration rate, leaf shape, tissue color, and also variation in the interaction of solar radiation with plants. Currently the most promising techniques are sensors that measure reflectance, temperature or fluorescence.

ISBN : 9788119821327

INTRODUCTION

Remote Sensing (RS), Geographic Information Systems (GIS), and Global Positioning System (GPS) technologies have become integral tools in modern plant pathology for efficient disease monitoring, analysis, and management.

1. Remote Sensing (RS):

Remote sensing involves collecting information about an object or area without direct physical contact. In plant pathology, RS utilizes satellite or airborne imagery to assess vegetation health, detect stress, and identify disease patterns. This technology allows for large-scale monitoring, providing valuable insights into the spatial distribution of diseases across fields.

2. Geographic Information Systems (GIS):

GIS integrates spatial data, such as maps and satellite images, to analyze and interpret relationships between different geographic features. In plant pathology, GIS is used to map disease distribution, assess environmental conditions, and identify factors contributing to disease spread. GIS helps researchers and farmers make informed decisions for disease management.

3. Global Positioning System (GPS):

GPS technology provides accurate location data, enabling precise mapping and monitoring of disease occurrences. This is crucial for understanding the spatial dynamics of diseases, tracking their progression over time, and implementing targeted control measures. GPS is often used in conjunction with GIS for detailed spatial analysis.

Applications of RS, GIS and GPS

Disease Mapping: RS and GIS facilitate the creation of disease distribution maps, helping to identify disease hotspots and patterns. This information aids in targeted interventions.

Monitoring Environmental Factors: RS can monitor environmental variables like temperature, humidity, and vegetation indices, providing insights into conditions favorable for disease development.

ISBN : 9788119821327

Precision Agriculture: GPS allows for the precise mapping of disease-affected areas, enabling farmers to apply control measures only where needed. This reduces resource use and minimizes environmental impact.

Early Detection: RS can detect subtle changes in vegetation health that may indicate the onset of diseases before visible symptoms appear. Early detection is crucial for timely intervention.

Decision Support Systems: Integrating RS, GIS, and GPS data into decision support systems helps researchers and farmers make informed decisions for disease management, optimizing resource allocation and reducing economic losses.

In summary, the integration of RS, GIS, and GPS technologies in plant pathology enhances the efficiency and effectiveness of disease monitoring, analysis, and management practices, ultimately contributing to improved crop health and yield.

Agriculture is the basis of every economy worldwide. Crop production is one of the major factors affecting domestic market and is also a major prerequisite of economic development in any country. A lot of issues are responsible for estimated crop production varying in different parts of the world. Among them plant diseases acts as a major threat to farmers as they lead to major destruction in overall food supply. To provide effective measures for detection and avoidance of the destruction requires an early identification of type of plant disease present. As conventional disease diagnostic methods are time consuming, demand trained experts there is need for new, automated sensor-based methods for detection and quantification of plant diseases (Mutka *et al*., 2015).

Currently, new non-invasive methods for diagnosing plant diseases using sensor technologies, robotics, computer vision and machine learning are rapidly developing. These sensors assess the optical properties of plants within different regions of the electromagnetic spectrum and are able to utilize information beyond

ISBN : 9788119821327

the visible range thus allow us to quantify disease symptoms that are invisible to naked eye. They enable the detection of early changes in plant physiology due to biotic stresses, because it can modifies the transpiration rate, leaf shape, tissue color, and also variation in the interaction of solar radiation with plants. Currently the most promising techniques are sensors that measure reflectance, temperature, or fluorescence.

Imaging techniques is an alternative, non-destructive efficient method to assess plant disease over a large area. It provide a better means to quantify disease stress than visual assessment methods. Various spectroscopic and sensor-based methods like fluorescence imaging, multi or hyperspectral imaging, infrared spectroscopy, fluorescence spectroscopy, visible spectroscopy have been developed for early detection, identification and quantification of plant diseases (Singh *et al*., 2020).

Imaging techniques that detect the electromagnetic spectrum outside of visible light allow us to quantify disease symptoms that are not visible by eye, increasing the range of symptoms we can observe and potentially allowing for earlier and more thorough symptom detection.

Plant leaf disease detection and classification

Data collection:

Our collected data contains different images of the leaf diseases that were taken from the internet, and these images are stored as a database in the computer, and they are used also to train our along with the healthy leaf images. Below are samples of images of the three diseases that we picked for our dataset

ISBN : 9788119821327




Fig 1: Input image

The process of plant disease detection system basically involves five steps:

1. Image Acquisition

The first important step in the analysis of plant diseases is to obtain high-quality images that meet the objectives of research. In this phase, images of plant leaves are gathered using digital media like camera, mobile phones etc. with desired resolution and size. The images can also be taken from web.

The right choice of sensors and platforms, the correct setting of the spatial and spectral resolution, lighting scheme, scan rate, frame rate and exposure time are prerequisites for obtaining accurate results. The formation of database of images is completely dependent on the application system developer. The image database is responsible for better efficiency of the classifier in the last phase of the detection system.

ISBN : 9788119821327




Fig 2: Image acquisition in different angles, during all growth stages of crop, different locations and seasons

2. Image Preprocessing:

The next step is image preprocessing, is to improve image data by removing background, noise and also suppressing undesired distortions. It enhances image features for processing and analysis. The images stored in RGB format are resized to standard size. These resized RGB images are then converted to HSV format.

Preprocessing of hyperspectral image includes includes calibration and spectrum correction. The goals of the calibration process are to standardize the spectral and spatial axes of the hyperspectral image, evaluate accuracy and reproducibility of the acquired data under different operating conditions, eliminate curvature effect and instrumental errors.

ISBN : 9788119821327




Fig 3: Image preprocessing

3. **Segmentation:**

Image segmentation is applied to simplify the illustration of image with segments so that it can be more meaningful and easily analyzed. Image segmentation is performed to segment the disease affected and unaffected portions of the leaf. As the premise of feature extraction, this phase is also the fundamental approach of image processing.

K-Means clustering method is used for partitioning of images into clusters in which at least one part of cluster contain image with major area of diseased part. The k Means clustering algorithm is applied to classify the objects into K number of classes according to set of features (Al-Hiary et al., 2011). Other methods includes threshold-based, watershed algorithm, edge detection.

ISBN : 9788119821327




Fig 4: Image Segmentation

4. Feature Extraction:

After segmentation the infected region, various features are extracted to describe the infected region. Hence, in this step the features from this area of interest need to be extracted. These features are needed to determine the meaning of a sample image.

Features can be based on colour, shape, and texture. Recently, most of the researchers are intending to use texture features for detection of plant diseases. HSV color features are important to sense image environment, recognize objects and convey information. Texture is one of the most important feature which can be used to classify and recognize objects. There are various methods of feature extraction that can be employed for developing the system such as gray-level co-occurrence matrix (GLCM), color cooccurrence method, spatial grey-level dependence matrix, and histogram based feature extraction. The GLCM method is a statistical method for texture classification.

5. Classification:

After extracting color and texture features, classification is final step. This phase implies to determine if the input image is healthy or

ISBN : 9788119821327

diseased. If the image is found to be diseased, some existing works have further classified it into a number of diseases. classification is performed by using Support Vector Machine (SVM) classifier, KNN classifier and ESEMBLE classifier. They proposed the use of digital image processing techniques to successfully detect and classify four major plant leaf diseases, namely Bacterial Blight, Cercospora Leaf Spot, Powdery Mildew, and Rust, as well as to automatically detect the symptoms of diseases as early as they appear on the growing stage. (Oo and Htun, 2018).




Fig 5: Uploading the Input images




Fig 6: Segmentation of the input images




Fig 7: Classification - Result detection disease




Fig 8: Classification - Result accuracy

<u>Optical sensors for assessment of plant disease symptoms</u>

- **Digital imaging (RGB)**
- **Hyperspectral imaging (HSI)**

ISBN : 9788119821327

- **Chlorophyll fluorescence imaging (CFI)**
- **Thermal imaging (TI)**

1. <u>RGB imaging</u>

- RGB stands for Red, Green, and Blue, the three primary colors used to represent the full spectrum of colors visible to the human eye. It is a commonly used technique in plant disease detection and monitoring, because detection by this method is simple, cost-effective, and also due to wide availability of RGB cameras.
- Involves capturing images using cameras equipped with red, green, and blue sensors, and then combining these channels to produce a full-color image. Several software packages, such as ASSESS 2.0, "Leaf Doctor," Scion Image software, and custom-made modules are used.




Fig 9: Overview of current sensor technologies used for the automated detection and identification of host-plant interactions. These sensors can be implemented in precision agriculture applications and plant phenotyping on different scales from single cells to entire ecosystem

ISBN : 9788119821327



RGB



HSI



CFI



TI

- Disease severity can be extracted as diseased pixels or as a percentage after the background is masked from the object of interest.
- Uniform focus, sharpness, and illumination, imaging angle (leaf orientation) and distance between the object and the sensor (pixel size) are crucial for high accuracy and reliable results and also have are additional influences on the image quality

ISBN : 9788119821327




Fig 10: RGB imaging




Fig 11: Segmentation od RGB image

2. **<u>Hyperspectral imaging (HSI)</u>**

- Hyperspectral imaging is a fast and nondestructive sensing technology.
- It is a relatively new technology that involves the acquisition of electromagnetic spectra at every pixel in an image, thus combining spatial and spectral information.
- Hyperspectral remote sensing is also known as high spectral resolution remote sensing. According to the international remote sensing community, a spectral resolution of $10^{-1}\lambda$ is defined as multispectral, and a spectral resolution of $10^{-2}\lambda$ is defined as hyperspectral.

ISBN : 9788119821327

Principles of Hyperspectral Imaging Technology and Sensors

- Hyperspectral data frame is formed as a three-dimensional image cube. The X-Y dimension expresses the spatial position information of the image. The third dimension (λ) is the spectral/wavelength dimension, which is composed of several bands in the spectral space.
- By making a multidimensional section of the spectral image cube, different types of spectral features can be obtained, such as the spectral characteristics at any point pixel.
- In addition to RGB bands, the hyperspectral sensor can also cover the visible band of the electromagnetic spectrum from 400 nm to 700 nm, the near-infrared band from 700 nm to 1000 nm, and the shortwave infrared band from 1000 nm to 2500 nm.
- The spatial resolution has a strong influence on the detection of plant disease or plant-pathogen interaction. It depends on distance between the sensor and the object. Sensors with a spatial resolution of approximately 1m is more suitable(Mahlein, 2016).

ISBN : 9788119821327




Fig 12: Structure of hyperspectral image data cube




Fig 13: Hyperspectralrange in Electromagnetic spectrum

Photo Response of Pathogen-Infected Plants

Light can interact with plant tissue in the following ways :

i. light transmission through a leaf

ii. light that is absorbed by leaf chemicals (e.g., pigments, water, sugars, lignin, and amino acids)

iii. light reflected from internal leaf structures or directly reflected from the leaf surface




Fig 14: Interaction between natural light and plant leaf surface

ISBN : 9788119821327




Fig 15: Leaf pathogen interaction during pathogenesis

Changes in reflectance due to plant pathogens and plant diseases can be explained by impairments in the leaf structure and chemical composition of the tissue during pathogenesis that is highly specific.




Fig 16: Spectral signatures of healthy barley leaves (green) and diseased barley leaves with net blotch (yellow), brown rust (brown), and powdery mildew (blue), respectively, at 10 days after inoculation.

ISBN : 9788119821327

The reflectance characteristic of a plant results from the biochemical compounds present in the leaves, and the physical characteristics of leaves (Mishra et al., 2017). The interaction between light and plants also depends on the wavelength. In the visible wavelength range (400–700 nm), the surface of the plant has a low reflectivity due to the absorption of light by photosynthetic pigments (chlorophylls, anthocyanins and carotenoids). In the near infrared (700–1100 nm), the reflectance increases due to light scattering in the intercellular space. In the short wave infrared range (1100–2500 nm), healthy plants have a low reflectance due to the absorption of light by water, proteins and other carbon components. The green color of the leaf is consistent with the characteristic reflection peak at 550 nm.




Fig 17: Comparison of (a) RGB image, (b) multispectral imaging, and (c) hyperspectral imaging.

Specifically, RGB images are obtained on the R/G/B channels, multispectral images are obtained at a small number of discontinuous wavelengths, and hyperspectral images are obtained at a large number of continuous wavelengths.

3. <u>Chlorophyll fluorescence imaging (CFI)</u>

- Visible disease symptoms do not provide all of the available information about plant health and may not be the best indicator for plant disease severity, especially early during infection. Plants that are experiencing biotic and abiotic stresses exhibit changes in chlorophyll fluorescence emission.

ISBN : 9788119821327

- Light-energy absorbed by plants is distributed over three competing processes: **photosynthesis, thermal dissipation and chlorophyll fluorescence emission**. An increase in Chl-FI thus implies a decrease in photosynthesis and/or thermal dissipation, and vice versa.




Fig 18: Origin of chlorophyll fluorescence

- Under normal conditions, most chlorophyll fluorescence is emitted from photosystem II (PSII) at 685 nm. When plants experience stress, this results in altered patterns of chlorophyll fluorescence emission, which can be observed with fluorescence imaging.

➢ **Chlorophyll fluorescence parameter to measure altered patterns of fluorescence emission are**

- One of the most widely studied parameters based on chlorophyll fluorescence is Fv/Fm, also known as the maximum quantum efficiency of PSII. This parameter is calculated from Fm, the maximum fluorescence of a dark-adapted leaf, and Fv, the difference between Fm and the minimum fluorescence from dark-adapted leaf (F0). While non-stressed plants maintain a consistent Fv/Fm value, various studies have shown that plants

ISBN : 9788119821327

experiencing biotic or abiotic stresses have decreased Fv/Fm values. Fv / Fm value decreases as disease progresses. Changes in this parameter occur before visible disease symptoms occurs.

- Non-photochemical quenching (NPQ or q_N) is a measure of energy that is dissipated as heat from photosynthetic reaction centers
- ΦPSII, the effective quantum yield of PSII




Fig 19: Effect of the infection of leaves by a virulent strain of *Pseudomonas syringae* on conventional chlorophyll fluorescence parameter F_V/F_M

Arabidopsis thaliana control plants (left column) and plants 6 h (middle column) and 24 h (right column) after infiltration by the virulent strain DC3000 of *P. syringae*. The control plants were untreated or infiltrated with 10 mM $MgCl_2$. The white arrows indicate which leaf and which half of the leaf was treated in the experiments. (A) Colour photographs. The leaves remained attached to plants. (B**)** False colour image showing F_V/F_M—maximum quantum yield of PSII.

Overall, chlorophyll fluorescence imaging has benefits in providing physiological information about plant health, which in some cases allows for detection of disease prior to visible symptoms appearing.

ISBN : 9788119821327




Fig 20: Working of chlorophyll fluorescence imaging system

4. **Thermal imaging (TI)**

Depending on the nature of infection, pathogens have different effects on the temperature of infected plant tissues. Temperature is negatively correlated with transpiration rate. Thus, pathogens that induce stomatal closure in plants often lead to decreased transpiration rates and increased leaf temperature.

Infrared thermography imaging can detect relative differences in leaf surface temperature by detection of infrared radiation from leaves. Different stages of infection have different effects on leaf temperature. Initially causes localized decreases in surface temperature, which are thought to be due to suppression of stomata closure early during infection. At later stages when the pathogen has caused areas of necrosis, the temperature of the infected leaf tissue increases to levels that are higher than that of uninfected tissue, which may be due to the inability of damaged tissue to perform natural cooling through transpiration.

Infrared thermography (IRT) assesses plant temperature and is correlated with plant water status, the microclimate in crop stands, and with changes in transpiration due to early infections by plant pathogens. Emitted infrared radiation in the thermal infrared range from 8 to 12 mm can be detected by thermographic and infrared

ISBN : 9788119821327

cameras and is illustrated in false color images, where each image pixel contains the temperature value of the measured object. In plant science, IRT can be used at different temporal and spatial scales from airborne to small scale applications

Thermographic and infrared cameras will illustrate the false color images, contains the temperature value of the measured object. Thermal IR measurements showed a higher radiance in the diseased area compared to that of the healthy one. The temperature difference allowed the discrimination between the infected and healthy leaves before the appearance of visible necrosis on leaves.




Fig 21: Infrared thermal imaging measurement




Fig 22: Monitoring of rose leaf colonization by *Peronospora sparsa* and symptom development of downy mildew in early stages (5 and 7 days after inoculation) of the disease by thermographic imaging.

ISBN : 9788119821327

Conclusion

RS, GIS and GPS are alternative, non-destructive efficient methods to assess plant disease over a large area. It provide a better means to quantify disease stress than visual assessment methods. Various spectroscopic and sensor-based methods like fluorescence imaging, multi or hyperspectral imaging, infrared spectroscopy, fluorescence spectroscopy, visible spectroscopy have been developed for early detection, identification and quantification of plant diseases.

The power of imaging lies in its ability to offer early detection of diseases, often before visible symptoms to the human eye. This early warning system allows farmers and researchers to take timely and targeted actions, mitigating the spread of pathogens and minimizing crop losses. By adopting these technologies, agricultural practices become more sustainable, as farmers can optimize resource utilization and reduce the reliance on harmful chemicals, thus promoting a healthier ecosystem.

REFERENCES:

1. CHAERLE, L., HAGENBEEK, D., DE BRUYNE, E., VALCKE, R. AND VAN DER STRAETEN, D., 2004, Thermal and chlorophyll-fluorescence imaging distinguish plant-pathogen interactions at an early stage. *Plant Cell Physiol.*, **45**(7): 887-896.
2. GRANUM, E., PÉREZ-BUENO, M.L., CALDERÓN, C.E., RAMOS, C., DE VICENTE, A., CAZORLA, F.M. AND BARÓN, M., 2015, Metabolic responses of avocado plants to stress induced by *Rosellinia necatrix* analysed by fluorescence and thermal imaging. *Eur. J. Plant Pathol.,* **142**: 625-632.
3. MAHLEIN, A.K., 2016, Plant disease detection by imaging sensors–parallels and specific demands for precision agriculture and plant phenotyping. *Plant Dis.*, **100**(2): 241-251.
4. MUTKA, A.M. AND BART, R.S., 2015, Image-based phenotyping of plant disease symptoms. *Front. Plant Sci.*, **5**(1): 734.

ISBN : 9788119821327

5. SINGH, V., SHARMA, N. AND SINGH, S., 2020, A review of imaging techniques for plant disease detection. *Artif. Intell. Agric.*, **4**(1): 229-242.
6. TAKTI, M. AND EL AZHARI, M.E.H., 2022, Plant disease detection using image processing.
7. WAN, L., LI, H., LI, C., WANG, A., YANG, Y. AND WANG, P., 2022, Hyperspectral Sensing of Plant Diseases: Principle and Methods. *Agronomy*, **12**(6): 1451.
8. ZHU, W., CHEN, H., CIECHANOWSKA, I. AND SPANER, D., 2018, Application of infrared thermal imaging for the rapid diagnosis of crop disease. *IFAC-Pap.*, **51**(17): 424-430.

ISBN : 9788119821327

CHAPTER – 11

Application of RS, GIS and GPS in Soil Resource Mapping

Sudarshan Varma

Ph.D. Scholar, Department of Soil Science, College of Agriculture, Keladi Shivappa Nayaka University of Agricultural and Horticultural Sciences, Shivamogga, Karnataka, India

Ph. No.: 9980807313

Email: sudarshanvarma95@gmail.com

Vilakar, K.

Ph.D. Scholar, Department of Soil Science, College of Agriculture, Keladi Shivappa Nayaka University of Agricultural and Horticultural Sciences, Shivamogga, Karnataka, India

Ph. No.: 9441684845

Email: kayithvilakar@gmail.com

Ravichandra

Ph.D. Scholar, Department of Plant Pathology, College of Agriculture, Keladi Shivappa Nayaka University of Agricultural and Horticultural Sciences, Shivamogga, Karnataka, India

Ph. No.: 9035440352

Email: ravichandragk13@gmail.com

ABSTRACT

The integration of Remote Sensing (RS), Geographic Information Systems (GIS), and Global Positioning System (GPS) technologies has significantly revolutionized soil resource mapping methodologies. This chapter encapsulates their pivotal roles and applications in this domain. RS techniques capture multispectral data, aiding in the identification and

ISBN : 9788119821327

analysis of soil properties from aerial and satellite imagery. GIS platforms provide a spatial framework for organizing, storing, and analyzing these diverse datasets, facilitating the creation of accurate soil maps. GPS technology ensures precise geolocation data, enhancing the spatial accuracy of soil information collection. Together, these technologies enable efficient soil characterization, classification, and mapping, crucial for agricultural planning, land management, and environmental conservation. The fusion of RS, GIS, and GPS speeds up soil resource assessment, offering valuable insights into soil variability, erosion risks, fertility levels, and suitability for different land uses, thereby enabling informed decision-making in sustainable land management practices.

Keywords: Remote Sensing, Soil Mapping, Geographic Information Systems, Global Positioning System, Spatial Analysis

Introduction

Remote Sensing (RS), Geographic Information Systems (GIS), and Satellite-based Positioning Systems (commonly known as GPS) stand as pivotal tools for the assessment and management of natural resources. RS involves gathering information about objects, phenomena, or processes through non-contact methods, typically from satellites or airplanes, using sensors across the electromagnetic spectrum. GIS facilitates the input, organization, analysis, and presentation of data collected via RS and other sources. GPS instruments accurately determine an object's location in terms of longitude, latitude, and altitude.

ISBN : 9788119821327

On a global scale, these technologies offer a cost-effective approach to studying interactions within the biosphere, geosphere, and atmosphere. At a smaller scale, space technology contributes valuable insights into developing land and water resources. For instance, monitoring changes in forest cover through RS and devising afforestation plans using GIS exemplify their application on macro and micro levels.

The abundance of remotely sensed data from diverse sensors on various platforms, offering a wide range of spatiotemporal, radiometric, and spectral resolutions, establishes remote sensing as one of the most comprehensive sources of data for large-scale applications and research. This extensive data has become a crucial input for numerous environmental process models

Overall, the integration of RS, GIS, and GPS technologies has revolutionized the assessment, management, and sustainable utilization of natural resources, providing crucial data and insights at varying scales of analysis and application.

Concepts of soil in soil resource Mapping soil

Soil: Soil is a three-dimensional natural entity, shaped by natural forces and human intervention, comprised of earth materials, organic matter and conducive for outdoor plant growth. Its upper boundary typically extends to the air or shallow water, while its lower limit often reaches hard rock or earthy material lacking evident biological activity.

Pedon: Pedon represents the smallest distinct unit of soil, often hexagonal in shape, encompassing a volume of soil. Its surface area ranges from 1 m^2 to 10 m^2, depending on the soil's characteristics and variability.

ISBN : 9788119821327

Profile: A profile refers to the vertical section of a pedon that exhibits the arrangement and nature of horizons. In soil resource inventories, profiles are examined to describe and analyze soil horizons, typically up to a depth of 2 meters or until bedrock is encountered.

Polypedon: A polypedon is a collection of several similar pedons or an inferred soil entity, serving as the basis for soil mapping and classification, often representing a soil series.

Mapping unit: A mapping unit comprises areas identified and labelled similarly in terms of soil series, associations, or types on a soil map. Each distinct area delineated on the map constitutes a separate delineation, differing in certain soil properties from other units on the map.

Taxonomic unit: Taxonomic units are aimed at classifying soils beyond the level of soil series, considering diagnostic horizons, soil temperature and moisture regimes, particle size, and mineralogical classes within the control section, often focusing on soil properties between 25 cm and 100 cm in depth. Taxonomic units include orders, suborders, great groups, subgroups, and families.

Base maps: Base maps serve as the foundation for delineating soil boundaries. In traditional soil surveys, toposheets and village maps were used, while modern surveys utilize base maps generated from aerial photographs and satellite data (photographic or digital).

Standard soil survey: The standard soil survey primarily focuses on observing and documenting the physical attributes of soils in their natural environment while analyzing their chemical properties within a laboratory setting. This process involves categorizing soils into distinct units based on well-defined criteria and outlining their boundaries on standardized maps (Anonymous, 1970). These surveys are classified

ISBN : 9788119821327

into three types based on the scale of the base map, the depth of soil observation, and the precision of mapping (Tables 1 and 2).

Table 1. Types of Soil Survey

	Reconnaissance survey (Low intensity/ small scale mapping)	**Detailed survey (High intensity/ large scale mapping)**	**Detailed Reconnaissance survey(Medium intensity)**
Base map	SOI toposheets 1: 50,000	Village maps/1: 4000/ 1:8000	This combines both reconnaissance and detailed soil surveys.
Soil profile	3 to 6 km	640 acres	A reconnaissance surveyis conducted throughout the project area and a detailed survey is conducted in intensive cultivation areas.
Soil observation (through auger)	¼ km to 1 km	¼ to ½ km	
Mapping unit	Soil series/soil association[1] / soil complex[2]	Types[3] and Phases[4]of the soil series	
Level of mapping	Taluk/District	Village/ block	

[1]Soil association consists of two or three soil series;

[2]Soil compiled consists of more than 3 soil series;

[3]Type means surface texture; and

[4]Phase includes solum depth, slope, erosion, gravelliness, stoniness, salinity and alkalinity classes.

Soil resource data

The site characteristics, morphological properties, physical properties, chemical properties and exchangeable properties (as

ISBN : 9788119821327

mentioned in Table 3) of the soil are essential in mapping land suitability and capability classification (FAO, 1988).

Table 2. Suggested Methods of Soil Survey

Maps	Scale	Base maps / RS tools	Level of use
Explanatory maps	1:1M & above	1:1 M toposheets satellite imageries	State / Country region
Small scale maps	1:250,000 scale	Toposheets / Satellite imageries	State / District / Agroclimatic region
Medium scale	1:50,000 scale	Toposheets /Aerial photographs / satellite imageries	Taluk / District
Large scale	1:10,000 to 1: 25,000	Village maps / AerialPhotographs/ Merged LISS III data	Village / Block

Table 3. Soil characteristics, morphological and analytical properties

Site Characteristics	Morphological	Physical	Chemical	Exchangeable
• Geology • Geomorphology • Drainage • Slope • Erosion • Land use • Natural vegetation • Depth of groundwater table	• Horizon thickness colour • Mottles • Texture • Structure Calcareousness • Concretions Abundance and size of roots and	• Particle size fractions • Bulk density • Moisture capacity at field capacity and	• pH • EC • $CaCO_3$ • Organic carbon • Total Nitrogen	• CEC • Ex. Cations • Ex. Acidity

ISBN : 9788119821327

• Stoniness • Gravelliness • Presence or absence of salinity and alkalinity	pores Permeability Presence of clay films/slickensides	permanent wilting point		

Use of aerial photographs in soil mapping

Aerial photography serves as a crucial tool in soil mapping, offering various types of images such as black and white, colour infrared (IR), and colour infrared (CIR) photographs. The scale of these aerial photographs varies from 1:40,000 to 1:60,000 for preliminary soil mapping and 1:10,000 to 1:25,000 for more detailed assessments. They provide a 3D view through stereoscopes, aiding in the identification of terrain features like slopes, drainage patterns, hills, valleys, and plains within a geological area.

Different landforms such as hills, pediments, pediplains, valleys, and alluvial plains can be delineated using photographic elements like slope, erosion, tone, texture, and land use. This assists in identifying physiographic units for each landform, which are then thoroughly examined to understand the soil composition. Aerial photography supports the identification of these units and their characteristics.

The process involving aerial photography for soil mapping involves steps depicted in Fig. 1, which may necessitate orthorectification if rectified aerial photographs are not initially utilized.

ISBN : 9788119821327




Fig. 1. Methods of soil mapping using aerial photographs

Use of Satellite data soil mapping

Additionally, satellite data is employed for soil mapping, available in various scales such as 1:1 million, 1:250,000, 1:50,000, and 1:25,000. Satellite images, particularly false colour composites (FCC) from the summer season, are preferred for soil mapping purposes. For detailed mapping, PAN-merged LISS (Linear Imaging Self-Scanning Sensor) images are utilized.

Similar to aerial remote sensing, satellite image interpretation involves delineating major landforms using elements like texture, tone, shape, size, association, and pattern, typically done on a light table. These interpretations help identify image interpretation units, and subsequently, soil compositions are determined through fieldwork followed by soil analysis, as shown in Fig. 2.

Digital image processing techniques like supervised and unsupervised classification are applied in soil mapping using satellite data. Supervised classification involves using training sets (clusters of

ISBN : 9788119821327

pixels with known compositions obtained through fieldwork) to generate soil maps. In unsupervised classification, a cluster map showcasing pixels with similar digital numbers (DN) is prepared, and fieldwork is conducted to assess the soil composition for each cluster. This ground truth information is then input into the computer system to generate accurate soil maps (Karale, 1992).




Fig. 2. Visual interpretation of satellite imageries for soil mapping

Soil maps

Soil survey maps: Maps are generated through standard methodologies using various tools like toposheets, village maps, aerial photography, or satellite data. These maps are published with appropriate scales after

ISBN : 9788119821327

undergoing necessary rectification processes. Generalized soil maps are created by consolidating the delineations from existing soil survey maps to form broader map units using cartographic techniques.

Schematic soil maps: Schematic soil maps usually at smaller scales (1:1 M and above), are compiled by utilizing existing maps related to geology, geomorphology, climate, land use, etc., often with limited field investigations. These maps prove beneficial in less developed areas before organized field surveys.

Digital soil maps: Digital soil maps created through scanning and digitizing existing soil maps using ground control points, serve as valuable information layers for generating various thematic maps either manually or through computer-based Geographic Information Systems (GIS).

Thematic maps: Thematic maps such as soil suitability maps or soil quality maps, are crafted for specific applications using GIS tools.

The selection of the method for soil resource mapping, which involves creating base maps using remote sensing tools like aerial photographs or satellite data, depends on factors such as available resources, funding, and time constraints. Remote sensing methods are often preferred over conventional methods due to their time and cost efficiency. Drawing from past soil surveys conducted across various projects, the following methods are recommended for different surveys to enhance effectiveness and efficiency without compromising accuracy.

Applications/interpretation of Soil Maps

Soil maps serve diverse purposes across different situations and applications. These applications demonstrate the wide-ranging utility of soil maps across numerous fields and industries. Some of the various uses include:

ISBN : 9788119821327

Land Capability Classification'

Land capability classification involves categorizing soils based on inherent soil characteristics, external land features, and environmental factors that restrict agricultural use. The classification employs eight classes, denoted by Roman numerals I to VIII, arranged in ascending order of limitations for land use. Classes I to IV are suitable for agriculture under specific and proper management, while classes V to VIII are better suited for wildlife sanctuaries and recreational purposes.

Within each land capability class, subclasses are represented by small letters such as 'e' for erosion, 's' for soil limitations, 'c' for climatic limitations, and 'w' for wetness. These subclasses identify specific soil groups with varying limitations and hazards. Moreover, land capability units are formed by grouping one or more soil mapping units that share similar potentials and ongoing limitations for land use.

Land Irrigability Classification

The classification of Land Irrigability focuses on predicting soil behaviour when subjected to altered water conditions resulting from irrigation practices. It takes into account soil irrigability classes (A to E), alongside considerations of topography and drainage. Using Arabic numbers 1 to 6, land irrigability classes are designated, with higher numbers indicating increased limitations.

Classes 1 to 4 are considered suitable for irrigation, while Class 5 is temporarily categorized as unsuitable pending further investigations. Class 6 includes lands permanently unsuitable for irrigation. Subclasses in land irrigability represent lands with similar limitations for sustained irrigation use. Lowercase letters like "s," "t," and "d" signify whether the limitations stem from soil properties, topography, or drainage. Lands

ISBN : 9788119821327

with multiple major deficiencies are denoted by relevant letters after the irrigability class.

Fertility Capability Classification

This system involves categorizing soils based on their fertility constraints in a way that can be qualified. It considers both the physical and chemical properties of the soil in the Fertility Capability Classification, aiding in grouping soils with similar fertility limitations and responses to fertilizers. The classification system comprises type, substrata type, and condition modifiers. Type is determined by the surface texture of the soil, denoted by letters (C, L, S, O). Substrata Type refers to the subsoil texture between depths of 20 and 50 cm (C, L, S, R). Condition modifiers represent physical and chemical properties influencing soil and fertilizer interactions. These modifiers include: g (gleying), d (dry), e (Low CEC), a (aluminium toxicity), h (acidic condition), i (Fe-P fixation), x (X-ray amorphous), v (Vertic characteristics), k (potassium deficiency), b (basic reaction), s (salinity), n (nitric), and c (clay with cat properties).

Land Suitability Classification

Land suitability classification assesses the appropriateness of a particular type of land for specific uses. It involves analyzing soil survey data, economic and social factors, existing land uses, and the potential need for change. This process results in separate classifications for various types of land uses relevant to the area. The classification system typically includes orders, classes, subclasses, and units. Two primary orders, suitable (S) and non-suitable (N), are identified. Within these orders, classes further distinguish suitability levels: S1 denotes highly suitable land, S2 signifies moderately suitable land, and S3 represents marginally suitable land. Subclasses within these classes indicate specific

ISBN : 9788119821327

limitations, similar to land capability subclasses. Each suitability unit within a subclass reflects differing management requirements.

Soil suitability models tailored for specific crops rely on the criteria defining soil site characteristics within existing management conditions. However, assessing soil suitability for a particular crop is based on its limiting characteristics, which might lead to underestimating a soil's overall suitability for that crop. It's important to note that the suitability of land for specific purposes depends on the scale, purpose, and intensity of the study. Depending on these factors, the classification system may adopt all or a limited number of categories (Zhang, 1994).

Soil Productivity Rating

A soil appraisal system has been developed to assess both the current and potential productivity of soil, based on a modified Storie Index. This system evaluates eight factors—moisture (H), drainage (D), depth (P), texture/structure (T), base saturation (N), soluble salt content (S), organic matter (O), and mineral reserves (A)—assigning ratings on a scale of 0-100. These individual percentages are then multiplied cumulatively to derive the productivity index (P). Similarly, a potentiality index (P`) is calculated, taking into account management interventions. The ratio of P` to P, known as the co-efficient of improvement (CI), indicates the extent to which productivity can be enhanced through implemented management measures.

$$P \text{ (or) } P` = H/100 \times D/100 \times P/100 \times T/100 \times N/100 \times S/100 \times O/100 \times A/100 \times 100$$

$$\text{Co-efficient of Improvement (CI)} = P`/P$$

Soil ratings are categorized into different ranges based on their index scores: 65-100 indicates excellent quality, 35-64 is considered

ISBN : 9788119821327

good, 20-34 is average, 8-19 is poor, and below 8 is deemed extremely poor. These ratings can be visually represented on maps depicting productivity and potentiality indexes. These productivity ratings serve as valuable guides for selecting the most suitable land use options among field crops, fodder crops, and tree crops. For instance, if a soil unit has productivity ratings of 60 for field crops, 80 for fodder crops, and 50 for tree crops, it suggests that the soil exhibits a higher potential for producing fodder crops compared to field or tree crops.

Soil Quality Maps

Soil maps derived from various parameters such as depth, erosion, texture, pH, and more are crucial in land management practices. These maps aid in crop selection and soil reclamation methods like liming for acidic soil, or gypsum application for alkaline soil.

Remote sensing and GIS in watershed characterization and management

Remote sensing and Geographic Information System (GIS) technology play vital roles in characterizing and managing watersheds. A watershed, defined by terrain topography, directs runoff to a specific point and varies in size depending on the base map's scale used for delineation. Satellite data of varying resolutions, along with topographic drainage maps, help delineate watersheds effectively. Techniques like stereo aerial photographs and satellite remote sensing are valuable in this process. Digital Elevation Models (DEMs), derived through GIS analysis from topographic contour information, assist in automated ridgeline and drainage network delineation. Watershed characteristics encompass topography, geology, soils, vegetation, land use, climate and exclude socio-economic factors.

ISBN : 9788119821327

Watershed prioritization

Prioritizing watersheds becomes essential to implement schemes considering catchment area size, issue severity, financial constraints, and local demands. Quantitative models estimating soil erosion help in prioritizing watersheds based on erosion estimates.

Remote sensing and GIS in soil erosion modelling

Soil erosion prediction has been a challenge for researchers, leading to the development of various models categorized as empirical, semi-empirical, and physical process-based models. Empirical models like the Universal Soil Loss Equation (USLE) rely on observations and statistics to predict soil erosion on hillslopes. Semi-empirical models, lying between empirical and physical process-based ones, use spatially lumped water and sediment equations. Physical process-based models, such as WEPP, aim to represent erosion's essential mechanisms, accounting for various factors' interactions and their spatial and temporal variations.

Universal soil loss equation (USLE)

The USLE, a widely used empirical equation for predicting soil erosion, was developed specifically for croplands on hillslopes. The equation's components include factors for rainfall, soil erodibility, slope length and steepness, cover management, and erosion control practices, providing a predictive estimate of soil erosion. This equation is as follows:

$$A = R.K.L.S.C.P$$

Where, A is the average annual soil loss (mass/area/year); R is the rainfall erosivity index; K is the soil erodibility factor; L is the slope

ISBN : 9788119821327

length factor; S is the slope gradient factor; C is the vegetation cover factor and P is the conservation protection factor.

Modified Universal soil loss equation (MUSLE)

It was proposed by Willams (1975). The modified version of the universal soil loss equation can be given as follows-

$$Sye = Xe.K.L.S.Ce.Pe$$

Where, Sye is the event sediment yield

$$Xe = (Qe, qp)\ 0.56$$

Where, 0.56 is an empirical co-efficient; Qe is the runoff amount and qp is the peak run-off rate obtained during the erosion and K.L.S.Ce & Pe as defined for USLE.

Morgan, Morgan and Finney Model (MMF)

The Morgan, Morgan and Finney (MMF) model aims to predict annual soil loss by incorporating advancements in erosion understanding into a more complex framework while retaining the simplicity of the Universal Soil Loss Equation (USLE). This model integrates recent insights into both the water phase and sediment phase of erosion processes. It operates through six equations requiring 15 input parameters. By comparing rain splash detachment predictions and runoff transport capacity, it determines the annual soil loss by identifying the limiting factor between detachment and transport.

Physical Contrast based model

In contrast to empirical models like USLE, which have limited applicability within specific ecological conditions and cannot account for deposition, various process-based models such as WEPP, EUROSEM, and LISEM have been developed. These models offer a broader scope,

ISBN : 9788119821327

addressing larger areas and watersheds, and considering deposition processes.

Sediment Yield Index (SYI) model

The Sediment Yield Index (SYI) model, developed by AISLUS for watershed prioritization in River Valley Projects, predicts soil erosion based on soil attributes, land use, and terrain slope characteristics (Spanner *et al*. 1983). Remote sensing tools, such as aerial photographs and satellite data, play a crucial role in mapping landscape features influencing erosion. These tools aid in studying erosional factors like soil type, slope gradient, land cover, rainfall interception, and vegetation dynamics through multi-temporal satellite images. Remote sensing data, including Digital Elevation Models (DEMs) derived from optical and microwave remote sensing, provide vital inputs for soil erosion modelling.

Geographic Information Systems (GIS) have emerged as powerful tools for handling spatial and non-spatial data, facilitating input visualization, model interaction, and erosion risk assessment. They play a significant role in soil erosion inventory, modelling, and assessment by integrating remote sensing data and providing valuable spatial information. While the USLE remains a frequently used method for assessing erosional soil loss, numerous studies demonstrate the potential of remote sensing and GIS techniques in quantitatively assessing soil erosion, highlighting their utility in erosion modelling and risk assessment without being limited by ecological conditions or scale.

Digital Database on Soils

Numerous organizations possess extensive data on soils, including maps and comprehensive attributes like soil properties,

ISBN : 9788119821327

geographical positioning, land usage, and lithology. However, a well-organized digital repository at the state or national level is lacking, which necessitates the establishment of a centralized digital database. Advancements in Geographic Information Systems (GIS), Relational Database Management Systems (RDBMS), Decision Support Systems, and rapid information technology development have led to the emergence of a new discipline called Soil Information Systems.

Soil and Terrain Digital Database (SOTER)

One notable system is the Soil and Terrain Digital Database (SOTER), which systematically organizes natural resource data for easy access, amalgamation, and analysis. This database focuses on areas characterized by distinct landforms, morphology, slopes, parent materials, and soils at a 1:1 million scale (SOTER UNITS). Each unit, integrated with a Geographic Information System, contains a computerized database encompassing topography, landform, soils, climate, vegetation, and land use attributes.

National Natural Resource Information System (NRIS)

The Department of Space, Government of India, has devised the National Natural Resource Information System (NRIS) aimed at providing essential information to decision-makers. NRIS consolidates data on land, soil, water, forests, and socio-economic parameters through remote sensing and conventional techniques. It functions as a network of GIS-based nodes spanning watersheds, blocks, districts, states, and the country, encompassing both spatial and non-spatial data inputs.

ISBN : 9788119821327

National Informatics Centre (NIC)

The National Informatics Centre (NIC) through NICENET provides spatial databases accessible via GISNIC software. These databases aid in information retrieval and presentation, assisting in the creation of thematic maps and extraction of attribute data from existing databases.

Soil information system for soil health card (SISSHC)

Another significant system, the Soil Information System for Soil Health Card (SISSHC), developed by AISLUS, advocates for implementing soil health cards in watershed management programs. These cards aim to raise awareness for improved soil and land resource utilization. The National Bureau of Soil Survey and Land Use Planning (NBSS&LUP) completed soil resource mapping in various states at scales of 1:2, 50,000 and 1:5000,000. This extensive soil resource study generated voluminous data from field studies and laboratory analyses, including around 75,000 soil profiles, stored in digital formats.

Efficient dissemination of this wealth of resource information in various formats is crucial for its utilization by planners, development agencies, and environmental improvement initiatives. The NBSS&LUP, identified as a sub-centre of the Agricultural Research Information Centre (ARIC) by ICAR, contributes to AGRIS, which covers soil science literature. This collaborative effort aids in facilitating access to vital information for agricultural planning and research without duplication.

Recent Advances: Hyper Spectral Remote Sensing

Conventional broad-band sensors like SPOT, Landsat MSS, IRS LISS III, and LISS IV are limited in mapping soil properties due to their

ISBN : 9788119821327

bandwidth of 100 to 200 μm, which cannot adequately resolve specific features of terrestrial materials. In contrast, hyperspectral sensors possess high spectral resolution across a wide electromagnetic spectrum range, enabling the identification of chemical compositions of imaged targets such as rocks, soils, or vegetation. Hyperspectral sensors capture reflected electromagnetic energy from the Earth's surface spanning from the visible wavelength to the near-infrared and mid-infrared regions (0.3μm to 2.5μm) in numerous narrow contiguous bands, typically in the order of 10nm. These narrow bandwidths offer detailed spectral responses for each pixel, providing precise information about constituents and presenting a clear advantage over multispectral imaging. This high spectral resolution helps detect subtle variations in materials' spectral responses, aiding in their identification.

Recent studies have demonstrated relatively strong correlations between soil reflectance and specific physical and chemical soil properties. It's also observed that the environmental conditions during soil formation influence soil reflectance. If these relationships between soil reflectance and its chemical and physical properties can be quantitatively established under specific environmental conditions, it would significantly enhance the extraction of valuable soil information from sensor data obtained by current and future earth observation satellite systems.

In recent years, there has been a growing emphasis on launching hyperspectral satellites for detailed regional characterization of land surface features. This shift aims to enhance the capacity to extract soil-related information from remote sensing data efficiently. Contrasting the traditional method involving soil sampling and laboratory analysis, remote sensing based on images or reflectance offers an efficient, rapid,

ISBN : 9788119821327

and cost-effective means to identify spatial differences in crop and soil conditions within fields. It holds the potential to identify intricate spatial patterns in soil properties across fields and optimize soil sampling strategies for quantifying these patterns. Several soil properties, such as surface condition, particle size, organic matter, soil colour, moisture content, iron and iron oxide content, and mineralogy, have been observed to influence their spectral behaviour.

Conclusion

Remote sensing and GIS have emerged as powerful tools for generating spatial information about natural resources. The extensive and often uncontrolled use of land has led to significant soil degradation. There is an urgent need for a dependable inventory of soils and other resources. Remote sensing data has proven to be instrumental in mapping soils and various resources. The advancement of a new generation of technology offering higher spatial resolution, expanded spectral coverage, improved revisit capabilities, and stereo viewing has opened new possibilities across different application domains. GIS technology is driving rapid transformations in the analysis and management of natural resources. Integration of GIS with remote sensing, GPS, and computer technology is introducing innovative approaches for data collection, storage, processing, analysis, and modelling.

References

1. Anonymous. 1970. Soil survey manual, IARI, New Delhi.
2. FAO. 1988. Guidelines for soil description. FAO. Rome.
3. Karale, R.L. 1992. Remote sensing with IRS 1A in soil studies: Developments, status and prospects. In: Natural Resources

ISBN : 9788119821327

Management – A new Perspective. (ed.) R.L. Karale. National Natural Resources Management System (NNRMS), Department of Space, Govt. of India, Bangalore, pp. 128-143.

4. Spanner, M.A., Strahler, A.H. Estes, J.E., 1983. Soil loss prediction in a Geographic Information System Format. In Papers Selected for Presentation at the Seventeenth International Symposium on Remote Sensing of Environment. 89-102. 2–9June 1982. Buenos Aires, Argentina. Ann Arbor, Mich.
5. Steininger, M.K., 1996. Tropical secondary forest regrowth in the Amazon: age, area and change estimation with Thematic Mapper data. International Journal of Remote Sensing. 17, 9–27.
6. Williams, J.R. 1975. Sediment yield prediction with Universal Equation using runoff energy factor. In: Present and prospective Technology for predicting sediment yields and sources, Agricultural Research Service, US Department of Agriculture. 244–252.
7. Zhang, L. 1994. A comparison of the efficiency of the three models to estimate water yield changes afterforced catchment conversion, M.Sc. (Forest) Thesis, University of Melbourne, Australia.

ISBN : 9788119821327

CHAPTER – 12

Optical Sensing Technology (OST) for Detection of Plants Diseases

Satendra Kumar Sharma (Ph.D., Scholar) **

Nentia Chib (Ph.D., Scholar)

Deepak Kumar (Prof./Chief Scientist)

Division of Plant Pathology

Sher-e-Kashmir University of Agricultural Science & Technology,

Jammu, 180009

Email of Corresponding Author: - ss0944619@gmail.com

ABSTRACT

Optical sensing technology, leveraging light to measure various phenomena, presents a non-contact approach that extends beyond human visual capabilities, detecting wavelengths outside the visible spectrum. In agricultural contexts, its application for plant disease detection proves promising, offering rapid and non-destructive alternatives to molecular techniques. This technology utilizes different sensors capturing specific electromagnetic spectrum regions, facilitating the detection of phenotype changes caused by diseases. In particular, image sensors play a pivotal role in disease detection by capturing high-quality images essential for accurate downstream processing. RGB image sensors, mirroring human biological visual sensors, detect red, green, and blue colors from a visible light spectrum, analyzing changes in plant tissues and abnormalities. They have found diverse applications in disease recognition, quality assessment of food items, and crop mapping. Additionally, they have been instrumental in assessing resistance to specific diseases and predicting grain yield. Moreover, aerial RGB

ISBN : 9788119821327

imaging has demonstrated effectiveness in monitoring plant diseases at a high resolution. Advancements in supervised classification using RGB aerial imagery have shown promise in evaluating disease impact, quantifying metabolite concentrations, and even determining microbial toxin contamination in plants. Further advancements involve combining RGB with thermal imaging for enhanced crop management, predicting plant traits, and monitoring different irrigation regimes. Machine learning techniques coupled with these imaging modalities have shown potential in accurately predicting plant traits, aiding in precise crop management strategies. The systematic selection of appropriate image features for classification and segmentation using pattern recognition algorithms has significantly enhanced classification accuracies. The evolution of digital image analysis resources and databases has further supported the utilization of optical sensing technologies in agriculture.

Keywords: Plant disease, detection, RGB, Remote sensing, precision, phenotyping, Optical sensors

Introduction

Optical sensing technology is a non-contact technology that utilizes light to measure objects or phenomena. This can be different by employing various light sources, such as lasers, and different sensing technologies, such as time-of-flight, structured light, and self-mixing interference. Additionally, it can also detect various properties of light, such as wavelength, frequency, polarization, absorbance, luminescence, and refractive index, and convert them into electric signals.

Optical design technology that combines various lenses can achieve high precision and stability even in low reflective or transparent conditions and sensors are capable of measuring both the frequency and

ISBN : 9788119821327

intensity of light, which can be interpreted using multivariate statistical techniques. Although it operates similarly to human vision, it can detect wavelengths of light beyond the visible spectrum that are not detectable by the human eye. Different sensors can measure specific regions of the electromagnetic spectrum, from ultraviolet (UV) to long-wave infrared (LWI). Optical sensing technologies can detect the phenotype changes caused by the diseases, presenting a rapid and non-destructive alternative to molecular techniques for plant disease detection. Additionally, it increases the objectivity of field-based visual assessment. Optical sensing methods can be classified by their platforms and associated scale of imagery, as well as by their spectral characteristics.

VARIOUS IMAGE SENSORS USED FOR DISEASE DETECTION

Imaging techniques are used for capturing, processing, storing, and representing visual information for investigation, analysis, or broadcasting. it has various applications in precision agriculture and can be a time-saving and cost-effective tool for farmers [1-2]. However, the most in crucial aspect of imaging techniques is the initial step of capturing high-quality images, which are necessary for accurate downstream image processing.

Acquisition of high-quality images for disease detection is essential and can be achieved using an appropriate image sensor. Modern imaging techniques are digital it is easier to store, process, and manipulate digital information than analog. Any image-capturing equipment or camera captures a band of light reflected from an object or landscape. The optical component of the camera adjusts the focus (focal length) and span of the scene (aperture) to capture a plane of the image. The principle is to capture reflected light from a focal scene and optically

ISBN : 9788119821327

transfer it to a light-sensitive grid of electronic pixels, known as an image sensor. Image sensors are available in various pixel configurations ranging from 80 × 60 to 1280 × 1024 pixels. There are different types of image sensors depending on the image sensor's sensitivity to a particular spectrum of light (electromagnetic waves), viz., thermal, RGB sensors, multispectral/hyperspectral sensors, etc. These sensors are equipped with surrounding electronics or Read-out Integrated Circuit (ROIC) that transform a weak photocurrent from grid-organized pixels to a final digital bit-map image. The camera's resolution depends on the number of pixels on the image sensor, which determines the quality of the image captured.

"Role of RGB Imaging Sensors in Agricultural Disease Detection"

Human eyes have three types of sensors sensitive to different colors in a scene. RGB (Red, Green, Blue) image sensors are specifically engineered to capture these three colors from the visible light spectrum reflected in a scene [3]. These sensors operate by focusing on the scene using an array of photodiodes arranged into clusters. Each cluster comprises photodiodes covered with Red, Green, Blue, and Clear filters. For example, a standard 8 × 8 RGB array typically contains 16 photodiodes for red, green, blue, and clear light, respectively (refer to Figure 1). When light passes through the red filter, only red light reaches the photodiode, generating an electrical signal representing red light. Similar signals are produced for green and blue light by other photodiodes. Specialized electronics then combine these signals into a square wave signal, subsequently processed and transmitted for analysis by a computer or IoT device.

ISBN : 9788119821327




Figure 1: Illustrates a standard RGB imaging process.

Table 1. RGB sensors used in the early detection of plant disease

S.no	Disease	Pathogen/causal agent	Plant host	Reference
1	Citrus Peel Diseases (Canker, greasy spot, scab, melanose and insect damage)	Multiple diseases	Grapefruits (Citrus × paradisi)	[4]
2	Olive leaf spot disease	*Spilocaea oleaginea*	Olive (*Olea europaea*)	[5]
3	Cercospora leaf spot	*Cercospora beticola*	Sugar beet (*Beta vulgaris*)	[6]
4	Stripe (yellow) rust	*Puccinia striiformis f. sp. tritici*	Winter Wheat (*Triticum aestivum*)	[7]
5	Potato late blight	*Phytophthora infestans*	Potato (*Solanum tuberosum* L.)	[8]
6	Chlorosis	Iron deficiency	Soybean (*Glycine*	[9]

ISBN : 9788119821327

			max)	
7	Tomato plant diseases	Multiple diseases	Tomato (*Solanum lycopersicum*)	[10]
8	Phymatotrichopsis root rot disease	*Phymatotrichopsis omnivora*	Alfalfa (*Medicago sativa*)	[11]
9	Aphanomyces root rot	*Aphanomyces euteiches*	Lentil (*Lens culinaris*)	[12]
10	Verticillium wilt	*Verticillium dahliae*	Olive (*Olea europaea*)	[13]

RGB-based imaging has seen a growing application in disease and pest recognition, quality assessment of diverse food items (both fresh and processed), crop and weed mapping, determination of secondary metabolite concentrations, non-invasive estimation of nutrient status (e.g., nitrogen content), and chlorophyll content estimation in micro propagated plants [14-15]. Employing sensors sensitive to red (wavelength 550-750 nm), green (wavelength 500-549 nm), and blue (wavelength 400-499 nm) regions of the spectrum, RGB imaging generates image data [16]. These sensors commonly quantify images based on three primary color values (red, green, and blue), resulting in comprehensive responses across spectral bands and a wide range of colors [17].

Plant optical properties rely on light transmission through tissues, pigment, sugar, and water absorption, as well as light reflection. Digital images produced by these sensors enable the analysis of plant tissues and anomalies at the field or farm level [18, 19]. In recent years, RGB indices have been utilized to evaluate losses caused by fungal

ISBN : 9788119821327

pathogens, determine resistance levels to specific diseases (such as wheat's yellow rust), and predict subsequent grain yield [20-21]. Furthermore, RGB sensors prove effective in acquiring high-resolution aerial images (from aircraft or satellites) that can be instrumental in plant disease monitoring.

RGB imaging serves various functions in plant pathology, notably in evaluating the concentration of metabolites produced by plant pathogens like mycotoxins, with significant implications for ensuring food quality and human health. For example, the use of RGB imaging facilitated the identification of deoxynivalenol, a toxin found in plants affected by *Fusarium graminearum*. This method has proven cost-effective, and user-friendly, and offers a viable alternative to traditional approaches in assessing pathogen effects on plant growth and microbial toxin contamination [22].

In olive plants, early detection of Verticillium wilt using RGB indexes showcased substantial differences between untreated and treated plants [23]. Furthermore, researchers, such as Goluguri and colleagues, extensively reviewed image classifiers and deep learning techniques employed in identifying various rice diseases. RGB imaging has been pivotal in detecting both biotic and abiotic stress factors in plants [25]. For instance, in soybean plants afflicted with iron deficiency chlorosis, Bai and co-workers utilized RGB imaging and statistical models for field-based assessment [25].

Plant responses to stress, whether biotic or abiotic, result in changes to their morphological and physiological traits, which can be assessed through imaging techniques. Imaging methods were used to evaluate leaf area, water consumption, and water potential in drought-stressed grapevines [26]. Recently, RGB sensors have been instrumental

ISBN : 9788119821327

in phenotyping plant root systems, crucial for understanding plant health [27]. Parameters such as water potential and chlorophyll content play vital roles in determining plant health under stress conditions. RGB reflectance was used to predict leaf water potential in potatoes and estimate chlorophyll content in micro-propagated potato plants [28][29]. Efforts have been made to develop distinct models for each image type, recognizing that thermal and red-green-blue (RGB) images possess varying resolutions, potentially impacting model performance when trained concurrently. In a recent study, researchers examined remote sensing indices obtained from both thermal and RGB images. They utilized stepwise multiple linear regression (SMLR) and an integrated adaptive neuro-fuzzy inference system with a genetic algorithm (ANFIS-GA) to monitor various traits in potato varieties under different evapotranspiration levels (100%, 75%, and 50%). Notably, the ANFIS-GA models exhibited superior predictive capabilities for traits like fresh weight, dry weight, water content, and total tuber yield during both the calibration and testing phases. Thus, the combined application of RGB and thermal imaging indices with ANFIS-GA models proves effective in managing potato crop growth and production under diverse irrigation regimes [30]. Distinct features extracted from RGB images, including color, Gray levels, texture, dispersion, connectivity, and shape parameters, hold significant utility in image classification and segmentation using various pattern recognition algorithms and machine learning approaches. Hence, a meticulous selection of appropriate image features is pivotal for achieving improved classification accuracies [31]. Moreover, resources such as 'Quantitative Plant' (https://www.quantitative-plant.org/) and databases now support

ISBN : 9788119821327

digital image analysis, offering tools and platforms for image-based research and analysis in plant science [32].

Thermal Sensors: Advancements in Agricultural Imaging for Plant Disease Monitoring

Temperature is a crucial physical parameter in many biological and industrial processes, and thermography has revolutionized several scientific disciplines, including plant science. Infrared (IR) thermal imaging, developed for military purposes, is commonly used in many fields such as engineering, aerospace, medicine, agriculture, and veterinary sciences [33-34]. Thermal imaging sensors use materials like InGaAs, which are more sensitive to shortwave infrared (wavelength ≈ 0.78 μm to 1.7 μm) and not the visible spectrum of light. Therefore, thermal cameras capture pictures of heat, not of visible light as RGB sensors. All objects, including ice, emit heat (IR waves) [35]. The measurement of emitted radiations (IR) from an object is generally referred to as thermography. Objects above -273.15°C emit in the infrared (IR) range of the electromagnetic spectrum. Thermal imaging converts the invisible radiation patterns of an object into analyzable, visible images. An IR thermal imaging system comprises a thermal camera with an IR detector, a signal processing unit, and a computer for image acquisition. The InGaAs sensor array is primarily made up of a 2-D photodiode array. Each photodiode absorbs the incident IR radiation and transforms it into an electric current, which is read out and calibrated by the ROIC [36]. The Photodiode array has three layers viz., indium phosphide (InP) substrate, InGaAs absorption layer, and an ultrathin InP cap (Figure 2). The InP cap is an indium bump bonded to an ROIC, to which the camera preamplifier electronics get connected.

ISBN : 9788119821327

Further, the off-chip electronics manipulate collected electrical voltages to a bit-map to form a digital image [37]. Various models of thermal cameras are available that work in different temperature ranges and a comprehensive listing of these models is available. The thermal imaging technique has been successfully utilized for the assessment of seed viability and estimation of water loss status in crops.

The thermal imaging technique has been successfully utilized for the assessment of seed viability, estimation of water loss status in crops, scheduling irrigation, yield estimation, determining maturity indices, and detecting diseases in crop plants [38]. IR thermography can detect small differences in plant temperature, attributable to different types of biotic and abiotic stressors [39-40]. Among biotic stress factors in plants, thermography has successfully detected diseases caused by different classes of pathogens such as bacteria, fungi, viruses, and phanerogamic parasites, in their early stages of infection. Moreover, the technique has been adopted to detect of insect pests [41]. In addition, the thermal imaging technique was used to detect the presymptomatic monitoring of wheat stripe rust [42]. Yang and colleagues also developed a fast infrared-based thermal image processing technique for monitoring tea diseases in tea orchards [43]. Charlie and colleagues reported the pre-symptomatic detection of tobacco mosaic virus (TMV) disease in tobacco plants using both chlorophyll fluorescence imaging (Chl-FI) technique and thermography [44].

ISBN : 9788119821327




Figure 2. A typical IR (infra-red) image sensor used for thermography




Figure 2. A typical IR (infra-red) image sensor used for thermography

ISBN : 9788119821327




Figure 3. Tea disease detection based on fast infra-red thermal image processing technology [45]

Disease Detection Using Hyperspectral and Multispectral Image Sensors"

Hyperspectral and Multispectral Image Sensors

Imaging sensors are generally categorized based on spectral resolution (i.e., the width of measurable wavebands), spatial scale, and the type of detector (imaging or non-imaging). Multi-spectral imagers capture spectral information of objects/scenes in several broad wavebands. A typical multispectral camera can provide image data in R, G, B, near-infra (NIR), and even UV wavebands. Multispectral camera systems use a composite hybrid set of spectral filters and image sensors to capture multiple images over multiple bands of the electromagnetic spectrum [46]. On the other hand, a hyperspectral camera spans an even wider range of spectral wavebands. Hyperspectral image data is usually organized as large matrices with spatial x- and y-axes and spectral

ISBN : 9788119821327

waveband information in the z-axis [47]. Furthermore, multispectral image (MSI) sensors can only quantify the reflectance of the scene (plant tissue in this case) in a few broad possibly non-contiguous, bands, whereas hyperspectral image sensors (HSI) can measure the spectrum more continuously (in narrow contiguous bands) from the scene. Therefore, HSI processes a larger amount of information than MSI and can handle a large amount of data at a particular time frame requiring higher computing configurations of the associated system to process the data and interpret results [48]. Figure 4 diagrammatically present the detailed differences between MSI and HIS.




Figure 4. A typical multispectral image sensor system

ISBN : 9788119821327

Table 3. Hyperspectral image sensors used in early detection of plant disease

S. No.	Disease	Pathogen	Plant host	Reference
1	Dark leaf and pod spot	*Alternaria Spp.*	Brassica napus (oilseed rape)	[49]
2	Tobacco mosaic	*Tobacco mosaic virus*	*Nicotiana tabacum (tobacco)*	[50]
3	Powdery mildew	Erysiphe necator	*Vitis vinifera* (wine grapes)	[51]
4	Leaf rust	Puccinia triticina	*Triticum Spp.* (wheat)	[52]
5	Powdery mildew	*Blumeria graminis*	*Hordeum vulgare* (barley)	[53]
6	Charcoal rot	*Macrophomina phaseolina*	*Glycine max* (soybean)	[54]
7	Cercospora leaf spot	*Cercospora beticola*	*Beta vulgaris* (sugar beet)	[55]
8	Downy mildew	*Plasmopara viticola*	*Vitis vinifera* (wine grapes)	[56]
9	Gummy stem blight	*Mycosphaerella melonis*	*Cucumis sativus* (cucumber)	[57]

ISBN : 9788119821327

Multispectral image sensors used in early detection of plant disease

Multispectral image sensors have been instrumental in early plant disease detection because they can capture spectral changes in diseased plants. Which are often linked to histological and biochemical shifts. This change can now be analyzed through modern applications such as the PROSPECT Model. However, handling and processing of data from these sensors can be intricate and labour-intensive.

To address this complexity, machine learning methods have emerged as invaluable tools for analyzing numerous spectral bands. They are increasingly utilized for monitoring plant diseases across laboratories, greenhouses, and field settings [58][59]. Multispectral imaging, covering wavelengths from 395 to 970 nm, has been employed to assess the seed health of *Spinacia oleracea* L. Additionally, it has been successfully utilized to differentiate between aflatoxin-contaminated and non-contaminated hazelnuts, as well as contaminated ground red chili pepper flakes [60][61]. In various instances, multispectral imaging has proven effective in disease classification. For instance, it achieved a remarkable 95.6% accuracy in identifying aflatoxin contamination in hazelnuts and screening barley resistance against powdery mildew disease [62]. Narrow-banded LEDs facilitated the acquisition of multispectral images at ten spectral bands, aiding in the screening and diagnosis of powdery mildew in barley plants. A handheld, cost-effective multispectral imaging sensor was developed for in-field detection and classification of the whitefly, a vector of mosaic and brown streak diseases in cassava plants [63]. Moreover, multispectral imaging has been applied to detect early symptoms of leaf spot disease in Winter oilseed rape and diagnose grapevine trunk disease (Esca) in vineyards.

ISBN : 9788119821327

These methods, whether airborne or ground-based, showcased efficiency in detecting foliar symptoms of Esca in vineyards over multiple years [64][65][66]. These varied applications highlight the versatility and effectiveness of multispectral imaging techniques in diverse agricultural settings, enabling early disease detection, classification, and monitoring across various crops.

Table 4. Multispectral image sensors used in early detection of plant disease

S. No.	Disease	Pathogen	Plant host	Reference
1	Soybean Rust	*Phakopsora pachyrhizi*	Soybean (*Glycine max*)	[67]
2	Tomato spot wilt disease	Tomato spotted wilt virus	Peanut (*Arachis hypogaea*)	[68]
3	Mosaic and brown streak disease	*Bemisia tabaci* (whitefly) Insect vector	Cassava (*Manihot esculenta*)	[63]
4	Barley's powdery mildew	*Blumeria graminis* f.sp. *hordei*	Barley (Hordeum vulgare cv. Ingrid wild type)	[62]
5	Citrus greening	*Candidatus liberibacter*	Citrus fruits	[69]
6	Leaf spot in oilseed rape	*Pyrenopeziza brassicae*	Winter oilseed rape (*Brassica napus* L.)	[64]

ISBN : 9788119821327




Figure 6. Multispectral disease detection techniques involve using multiple spectral bands or wavelengths to detect and identify diseases in agriculture.

Advancements and Future Prospects in Disease Detection Technologies for Agriculture

Traditionally, assessing plant health relied on observing the plant's physical appearance, limited to the visible range of the electromagnetic spectrum. However, advancements in image acquisition sensors, such as IR sensors, multispectral sensors, and hyperspectral sensors, coupled with enhanced computing capabilities, have expanded our ability to explore beyond the visible spectrum. These technologies now enable the early detection of plant diseases well before they are observable by the human eye [47].

Sethy *et al*., provided a comprehensive review, of the progress in image acquisition, preprocessing, segmentation, feature extraction, and classification techniques for assessing paddy. They proposed a standardized framework for and emphasized the critical impact of image quality, training set size, and feature selection on performance [70]. Additionally, they recommended the use of support vector machines (SVMs) was recommended for precise disease classification and image quantification [71].

The integration of digital imaging, aerial mapping, and advanced technologies such as image processing, wireless sensor networks

ISBN : 9788119821327

(WSNs), Internet of Things (IoT), and Artificial Intelligence (AI) is reshaping modern agriculture [72]. This convergence is paving the way for smart agriculture, facilitating automated monitoring of various plant-related parameters such as soil, water, nutrition, diseases, pests, irrigation, and greenhouse conditions through interconnected sensor nodes communicating over the internet [72]. Cloud-based solutions are anticipated to further streamline field data collection, leveraging cutting-edge technologies like data analytics and AI in smart agriculture [73]. This transformative blend aims to foster a safer environment, robust economy, contented farmers, and improved accessibility to affordable and nutritious food. Advancements in technology bring us closer to achieving real-time monitoring, identification, and classification of plant diseases. Diverse image features can be efficiently utilized for classification and segmentation using various pattern-recognition algorithms and machine-learning approaches [47]. AI-based decision support systems hold promise in enabling precision agriculture by offering early recommendations for plant protection interventions, such as fungicide spraying. With the increasing use of smartphones among farmers, a user-friendly mobile application could serve as a valuable tool, empowering stakeholders with valuable insights into plant health from a plant pathologist's perspective, at least for RGB images.

Reference

1. Vibhute, A. and Bodhe, S. K. (2012). Applications of image processing in agriculture: a survey. *International Journal of Computer Applications*, 52(2).

ISBN : 9788119821327

2. Narvaez, F. Y., Reina, G., Torres-Torriti, M., Kantor, G. And Cheein, F. A. (2017). A survey of ranging and imaging techniques for precision agriculture phenotyping. *IEEE/ASME Transactions on Mechatronics*, 22(6): 2428-2439.
3. Vit, A. and Shani, G. (2018). Comparing RGB-D sensors for close-range outdoor agricultural phenotyping. *Sensors,* 18(12): 4413.
4. Zhao, X., Burks, T. F., Qin, J. and Ritenour, M. A. (2009). Digital microscopic imaging for citrus peel disease classification using color texture features. *Applied Engineering in Agriculture*, 25(5):769-776.
5. Al-Tarawneh, M. S. (2013). An empirical investigation of olive leave spot disease using auto-cropping segmentation and fuzzy Cmeans classification. *World Applied Sciences Journal*, 23(9): 1207-1211.
6. Zhou, R., Kaneko, S. I., Tanaka, F., Kayamori, M. and Shimizu, M. (2014). Disease detection of Cercospora Leaf Spot in sugar beet by robust template matching. *Computers and electronics in agriculture*, 108: 58-70.
7. Zhou, B., Elazab, A., Bort, J., Vergara, O., Serret, M. D. And Araus, J. L. (2015). Low-cost assessment of wheat resistance to yellow rust through conventional RGB images. *Computer Electron. Agric*., 116: 20-29.
8. Sugiura, R., Tsuda, S., Tamiya, S., Itoh, A., Nishiwaki, K., Murakami, N., Shibuya, Y., Hirafuji, M. and Nuske, S. (2016). Field phenotyping system for the assessment of potato late blight resistance using RGB imagery from an unmanned aerial vehicle. *Biosystems Engineering*, 148: 1-10.

ISBN : 9788119821327

9. Bai, G., Jenkins, S., Yuan, W., Graef, G. L., and Ge, Y. (2018). Field-based scoring of soybean iron deficiency chlorosis using RGB imaging and statistical learning. *Frontiers in plant science*, 9:1002.
10. Hlaing, C. S. and Zaw, S. M. M. (2018). Tomato plant diseases classification using statistical texture feature and color feature. In 2018 IEEE/ACIS 17th International Conference on Computer and Information Science (ICIS) (pp. 439-444). IEEE.
11. Mattupalli, C., Moffet, C. A., Shah, K. N. and Young, C. A. (2018). Supervised classification of RGB aerial imagery to evaluate the impact of a root rot disease. *Remote Sensing*, 10(6): 917.
12. Marzougui, A., Ma, Y., Zhang, C., McGee, R. J., Coyne, C. J., Main, D. and Sankaran, S. (2019). Advanced imaging for quantitative evaluation of Aphanomyces root rot resistance in lentil. *Frontiers in plant science*, 10: 383.
13. Sancho-Adamson, M., Trillas, M. I., Bort, J., Fernandez-Gallego, J. A. and Romanyà, J. (2019). Use of RGB vegetation indexes in assessing early effects of Verticillium wilt of olive in asymptomatic plants in high and low fertility scenarios. *Remote Sensing*, 11(6): 607.
14. Tillet, N. P., Hague, T. and Miles, S. J. (2001). Comput. *Electron. Agric*., 32: 229-246.
15. Dutta Gupta, S., Ibaraki, Y. and Pattanayak, A. (2013). Development of a digital image analysis method for real-time estimation of chlorophyll content in micro-propagated potato plants. *Plant Biotechnology*. Rep., 7: 91-97.
16. Lowe, A. Harrison, N. and French, A. P. (2017). Hyperspectral image analysis techniques for the detection and classification of the early onset of plant disease and stress. *Plant Methodsm* 13: 80.

ISBN : 9788119821327

17. Susstrunk, S., Buckley R. and Swen, S. (1999). Standard RGB color spaces. In: Proceedings of the Seventh Color Imaging Conference: Color Science, Systems and Applications Putting It All Together. pp. 127- 134. Scottsdale, Arizona, USA: The Society for Imaging Science and Technology.
18. Mahlein, A. K. (2016). Plant disease detection by imaging sensors–parallels and specific demands for precision agriculture and plant phenotyping. *Plant disease*, 100(2): 241-251.
19. Kefauver, S. C., El-Haddad, G. Vergara-Diaz, O. and Araus, J. L. (2015). RGB Picture vegetation indexes for high-throughput phenotyping platforms (HTPPs). *Remote. Sens. Agr*. Ecosyst. Hydrol., Xviii. 9637, 96370J.
20. Maloney, P. V., Petersen, S., Navarro, R. A., Marshall, D., McKendry, A. L., Costa, J. M., and Murphy, J. P. (2014). Digital image analysis method for estimation of Fusarium-damaged kernels in wheat. *Crop Science*, 54(5): 2077-2083.
21. Vergara-Diaz, O., Kefauver, S. C., Elazab, A., Nieto-Taladriz, M.T. and Araus, J. L. (2015). Grain yield losses in yellow-rusted durum wheat estimated using digital and conventional parameters under field conditions. *Crop Journal*., 3: 200-210.
22. Cambaza, E., Koseki, S. and Kawamura, S. (2019). Why RGB imaging should be used to analyze *Fusarium graminearum* growth and estimate deoxynivalenol contamination. *Methods and protocols*, 2(1): 25.
23. Sancho-Adamson, M., Trillas, M. I., Bort, J., Fernandez-Gallego, J. A. and Romanyà, J. (2019). Use of RGB vegetation indexes in assessing early effects of Verticillium wilt of olive in asymptomatic

ISBN : 9788119821327

plants in high and low fertility scenarios. *Remote Sensing*, 11(6): 607.

24. Goluguri, N. R. R., Suganya D. K. and Vadaparthi, N. (2020). Image classifiers and image deep learning classifiers evolved in detection of *Oryza sativa* diseases: survey. *Artificial Intelligence Review*, 1-38.
25. Bai, G., Jenkins, S., Yuan, W., Graef, G. L. and Ge, Y. (2018). Field-based scoring of soybean iron deficiency chlorosis using RGB imaging and statistical learning. Frontiers in plant science, 9:1002.
26. Briglia, N., Montanaro, G., Petrozza, A., Summerer, S., Cellini, F. and Nuzzo, V. (2019). Drought phenotyping in Vitis vinifera using RGB and NIR imaging. *Scientia Horticulturae*, 256: 108555.
27. Bodner, G., Alsalem, M. and Nakhforoosh, A. (2021). Root System Phenotyping of Soil-Grown Plants via RGB and Hyperspectral Imaging. *In Crop Breeding* (pp. 245-268). Humana, New York, NY.
28. Zakaluk, R. and Ranjan, R. (2008). Predicting the leaf water potential of potato plants using RGB reflectance*. Canadian biosystems engineering*, 50: 7.1-7.12.
29. Yadav, S. P., Ibaraki, Y. and Gupta, S. D. (2010). Estimation of the chlorophyll content of micro-propagated potato plants using RGB-based image analysis. *Plant Cell*, Tissue and Organ Culture (PCTOC), 100(2): 183-188.
30. Elsayed, S., Salah, El-H., Msaad, K., Osama, E., Nasser, Al-S., Majed, A., Muhammad, U. T. and Waleed, D. (2021). Combining thermal and RGB imaging indices with multivariate and data-driven modeling to estimate the growth, water status, and yield of

ISBN : 9788119821327

potatoes under different drip irrigation regimes. *Remote Sensing*, 13(9): 1679.

31. Behmann, J., Mahlein, A.-K., Rumpf, T., Römer, C. and Plümer, L. (2014). A review of advanced machine learning methods for the detection of biotic stress in precision crop protection. *Precis. Agric*., 16: 239-260.
32. Lobet, G., Draye, X. and Périlleux, C. (2013). An online database for plant image analysis software tools. *Plant methods*, 9(1): 1-8.
33. Scarth, G. W., Loewy, A. and Shaw, M. (1948). Use of the infrared total absorption method for estimating the time course of photosynthesis and transpiration. *Can. J. Res*., 26: 94-107.
34. Oerke, E. C., Steiner, U., Dehne, H. W. and Lindenthal, M. (2006). Thermal imaging of cucumber leaves affected by downy mildew and environmental conditions. *Journal of Experimental Botany,* 57(9): 2121-2132.
35. Anonymous (2020). https://www.flir.com/discover/rd-science/how-do-thermal-cameras-work/," 16 June 2020. [Accessed 15 June 2021].
36. Yu, L., Guo, Y., Zhu, H., Luo, M., Han, P. and Ji, X. (2020). Low-Cost Microbolometer Type Infrared Detectors. *Micromachines*,11(9): 800.
37. Anonymous (2021). T. P. Instruments, InGaAs Cameras: The Basics, Teledyne Imaging Group, 2021. [Online].Available: https://www.princetoninstruments.com/learn/camerafundamentals/ingaas -sensors-the-basics.
38. Jones, H. G., Stoll, M., Santos, T., de Sousa, C., Chaves, M. M. and Grant, O. M. 2002. Use of infrared thermography for monitoring

ISBN : 9788119821327

stomatal closure in the field: Application to grapevine. *Journal of Experimental Botany*, 53(378): 2249-2260.

39. Leinonen, I. and Jones, H. G. (2004). Combining thermal and visible imagery for estimating canopy temperature and identifying plant stress. *Journal of Experimental Botany*, 55(401): 1423-1431.
40. Mastrodimos, N. Lentzou, D. Templalexis, C. Tsitsigiannis, D.I. and Xanthopoulos, G. (2019). Development of thermography methodology for early diagnosis of fungal infection in table grapes: The case of Aspergillus carbonarius. *Comput. Electron. Agric*., 165: 104972.
41. Al-doski, J., Mansor, S. B., Shafri, H. Z. B. M. and Zulhaidi, H. (2016). Thermal imaging for pests detecting a review. *Int. J. Agric. For. Plant*., 2: 10-30.
42. Yao, Z., He, D. and Lei, Y. (2018). Thermal imaging for early non-destructive detection of wheat stripe rust. In 2018 ASABE Annual International Meeting (p. 1). *American Society of Agricultural and Biological Engineers*.
43. Yang, N., Yuan, M., Wang, P., Zhang, R., Sun, J. and Mao, H. (2019). Tea diseases detection based on fast infrared thermal image processing technology. *Journal of the Science of Food and Agriculture*, 99(7): 3459-3466.
44. Chaerle, L., Hagenbeek, D., De Bruyne, E., Valcke, R. and Van Der Straeten, D. (2004). Thermal and chlorophyll-fluorescence imaging distinguishes plant-pathogen interactions at an early stage. *Plant and Cell Physiology*, 45(7): 887-896.
45. Humplík, J. F., Lazár, D., Husičková, A. and Spíchal, L. (2015). Automated phenotyping of plant shoots using imaging methods

ISBN : 9788119821327

for analysis of plant stress responses review. *Plant methods*, 11(1):1-10.

46. Pang J, Zhang TZ, Bassig BA and Mao C, Green tea consumption and risk of cardiovascular and ischemic related diseases: a meta-analysis. *Int J Cardiol* 202: 967–974 (2016).
47. Humplík, J. F., Lazár, D., Husičková, A. and Spíchal, L. (2015). Automated phenotyping of plant shoots using imaging methods for analysis of plant stress responses review. *Plant methods*, 11(1):1-10.
48. Mahlein, A. K. (2016). Plant disease detection by imaging sensors–parallels and specific demands for precision agriculture and plant phenotyping. *Plant disease*, 100(2): 241-251.
49. Yandun, F., Reina, G., Torres-Torriti, M., Kantor, G. and Cheein, F. A. (2017). A survey of ranging and imaging techniques for precision agriculture phenotyping. *IEEE/ASME Transactions on Mechatronics*, 22(6): 2428-2439.
50. Baranowski, P., Jedryczka, M., Mazurek, W., Babula-Skowronska, D., Siedliska, A. and Kaczmarek, J. (2015). Hyperspectral and thermal imaging of oilseed rape (Brassica napus) response to fungal species of the genus Alternaria. *PloS one*, 10(3): e0122913.
51. Zhu, H., Chu, B., Zhang, C., Liu, F., Jiang, L. and He, Y. (2017). Hyperspectral imaging for presymptomatic detection of tobacco disease with successive projections algorithm and machine learning classifiers. *Sci. Rep*., 7:4125.
52. Knauer, U., Matros, A., Petrovic, T., Zanker, T., Scott, E. S. and Seiffert, U. (2017). Improved classification accuracy of powdery mildew infection levels of wine grapes by spatial-spectral analysis of hyperspectral images. *Plant Methods*, 13(1):47.

ISBN : 9788119821327

53. Ashourloo, D., Mobasheri, M. R. and Huete, A. (2014). Evaluating the effect of different wheat rust disease symptoms on vegetation indices using hyperspectral measurements. *Remote Sens*., 6(6):5107-23
54. Kuska, M., Wahabzada, M., Leucker, M., Dehne, H. W., Kersting, K. et al. (2015). Hyperspectral phenotyping on the microscopic scale: towards automated characterization of plant-pathogen interactions. *Plant Methods*, 11:28.
55. Naga Subramanian, K., Jones, S., Singh, A.K., Sarkar, S., Singh, A. and Ganapathy Subramanian, B. (2019). Plant disease identification using explainable 3D deep learning on hyperspectral images. *Plant Methods*. 21, 15:98.
56. Leucker, M., Mahlein, A.K., Steiner, U. and Oerke, E.C. (2017). Hyperspectral imaging reveals the effect of sugar beet quantitative trait loci on Cercospora leaf spot resistance. Funct. *Plant Biol.,* 44:1-9.
57. Oerke, E. C., Herzog, K. and Toepfer, R. (2016). Hyperspectral phenotyping of the reaction of grapevine genotypes to *Plasmopara viticola*. *J. Exp. Bot*., 67:5529-43.
58. Vukadinovic, D., Polder, G. and Swinkels, G. J. (2016). Automated detection of *Mycosphaerella melonis*-infected cucumber fruits. *IFAC-Papers Online*, 49(16):105-9.
59. Jacquemoud, S and Baret, F. (1990). PROSPECT: a model of leaf optical properties spectra. *Remote Sens. Environ*., 34:75-91.
60. Singh, A., Ganapathy Subramanian, B., Singh, A.K. and Sarkar S. (2016). Machine learning for high-throughput stress phenotyping in plants. *Trends Plant Sci*., 21(2):110-24.

ISBN : 9788119821327

61. Olesen, M. H., Carstensen, J. M. and Boelt, B. (2011). Multispectral imaging as a potential tool for seed health testing of spinach (Spinacia oleracea L.). *Seed Science and Technology*, 39(1): 140-150.
62. Kalkan, H., Beriat, P., Yardimci, Y. and Pearson, T. C. (2011). Detection of contaminated hazelnuts and ground red chili pepper flakes by multispectral imaging. *Computers and Electronics in Agriculture,* 77(1): 28-34.
63. Kuska, M. T., Behmann, J., Großkinsky, D. K., Roitsch, T. and Mahlein, A. K. (2018). Screening of barley resistance against powdery mildew by simultaneous high-throughput enzyme activity signature profiling and multispectral imaging. *Frontiers in Plant Science*, 9: 1074.
64. Fennell, J., Veys, C., Dingle, J., Nwezeobi, J., van Brunschot, S., Colvin, J., and Grieve, B. (2018). A method for real-time classification of insect vectors of mosaic and brown streak disease in cassava plants for future implementation within a low-cost, handheld, in-field multispectral imaging sensor. *Plant methods*,14(1): 1-12.
65. Veys, C., Chatziavgerinos, F., AlSuwaidi, A., Hibbert, J., Hansen, M., Bernotas, G., Smith, M., Yin, H., Rolfe, S. and Grieve, B., 2019. Multispectral imaging for presymptomatic analysis of light leaf spot in oilseed rape. *Plant methods*, 15(1): 1-12.
66. Zhang, W., Zhu, Q., Huang, M., Guo, Y. and Qin, J. (2019). Detection and classification of potato defects using multispectral imaging system based on single shot method. *Food Analytical Methods*, 12(12): 2920-2929.

ISBN : 9788119821327

67. Bendel, N., Kicherer, A., Backhaus, A., Klück, H. C., Seiffert, U., Fischer, M., Voegele, R. T. and Töpfer, R. (2020). Evaluating the suitability of hyper-and multispectral imaging to detect foliar symptoms of the grapevine trunk disease Esca in vineyards. *Plant methods*, 16(1): 1-18.
68. Cui, D., Zhang, Q., Li, M., Zhao, Y. and Hartman, G. L. (2009). Detection of soybean rust using a multispectral image sensor. *Sensing and Instrumentation for Food Quality and Safety*, 3(1):49-56.
69. Patrick, A., Pelham, S., Culbreath, A., Holbrook, C. C., De Godoy, I. J. and Li, C. (2017). High throughput phenotyping of tomato spot wilt disease in peanuts using unmanned aerial systems and multispectral imaging. *IEEE Instrumentation and Measurement Magazine*, 20(3): 4-12.
70. DadrasJavan, F., Samadzadegan, F., Pourazar, S. H. S. and Fazeli, H. (2019). UAV-based multispectral imagery for fast Citrus Greening detection. *Journal of Plant Diseases and Protection*, 126(4): 307-318.
71. Sethy, P. K., Barpanda, N. K., Rath, A. K. and Behera, S. K. (2020). Image Processing Techniques for Diagnosing Rice Plant Disease: A Survey. *Procedia Computer Science*, 167: 516-530.
72. Zhou, R., Kaneko, S. I., Tanaka, F., Kayamori, M. and Shimizu, M. (2014). Disease detection of Cercospora Leaf Spot in sugar beet by robust template matching. *Computers and electronics in agriculture*, 108: 58-70.
73. Ayaz, M., Ammad-Uddin, M., Sharif, Z., Mansour, A. and Aggoune, E. H. M. (2019). Internet-of-Things (IoT)-based smart agriculture: *Toward making the fields talk. IEEE Access*, 7: 129551-129583.

ISBN : 9788119821327

74. Farooq, M. S., Riaz, S., Abid, A., Abid, K. and Naeem, M. A. (2019). A Survey on the Role of IoT in Agriculture for the Implementation of Smart Farming. *IEEE Access*, 7: 156237-156271.
75. Elijah, O., Rahman, T. A., Orikumhi, I., Leow, C. Y. and Hindia, M. N. (2018). An overview of Internet of Things (IoT) and data analytics in agriculture: Benefits and challenges. *IEEE Internet of Things Journal,* 5(5): 3758-3773.

ISBN : 9788119821327

CHAPTER - 13

Image Classification, GIS Integration and Acreage Estimation for Land Management

Dhaval Kirankumar Dwivedi

Faculty of Agriculture and Allied Sciences, Kantaben Kashiram Institute of Agricultural Sciences and Research, Ganpat University, Kherva, Mehsana, Gujarat

Image classification in remote sensing is a crucial technique that involves the categorization and labelling of pixels within satellite or aerial imagery into distinct land cover or land use classes. This process plays a pivotal role in extracting valuable information for various applications, such as agriculture, land management, environmental monitoring, and urban planning. The following are the key components of image classification:

- **Remote Sensing Data Acquisition:** Image classification relies on data collected from satellite or airborne sensors, capturing information across different wavelengths. Common sensor types include optical, multispectral, and hyperspectral sensors.
- **Preprocessing:** Before classification, remote sensing data undergoes preprocessing steps, including radiometric calibration, atmospheric correction, and geometric correction, to enhance the quality and accuracy of the imagery.
- **Training Data Collection:** Image classification requires labeled training data, representing each class of interest. Analysts manually annotate a subset of the image to train machine learning algorithms, teaching them to recognize specific features.
- **Feature Extraction:** Features are characteristics extracted from the image, such as spectral values, texture, and spatial patterns. These features serve as input parameters for classification algorithms.

ISBN : 9788119821327

- **Classification Algorithms:** Various classification algorithms are employed, ranging from traditional methods like Maximum Likelihood and Decision Trees to advanced machine learning techniques such as Support Vector Machines (SVM), Random Forest, and Convolutional Neural Networks (CNNs).
- **Validation and Accuracy Assessment:** The accuracy of the classification is evaluated through validation techniques, comparing the classified results with ground truth data. Common metrics include overall accuracy, producer's accuracy, and user's accuracy.

Applications:

1. **Agriculture:** Image classification aids in monitoring crop types, assessing crop health, and optimizing farming practices. It enables precision agriculture by providing insights into planting patterns, pest infestations, and irrigation needs.
2. **Land Management:** Classification helps in mapping and monitoring land cover changes, assessing deforestation, urban expansion, and ecosystem health. It supports land-use planning and sustainable resource management.
3. **Environmental Monitoring:** Image classification contributes to monitoring environmental parameters, such as water bodies, vegetation cover, and soil composition. It assists in assessing the impact of natural disasters and climate change.
4. **Urban Planning:** The classification of urban areas aids in understanding urban sprawl, infrastructure development, and population density. This information is valuable for city planning and resource allocation.

Challenges:

1. **Data Quality:** Poor data quality, including atmospheric interference and sensor noise, can impact the accuracy of classifications.

ISBN : 9788119821327

2. **Complexity of Land Cover Types:** Distinguishing between classes with similar spectral signatures can be challenging, requiring advanced algorithms and high-resolution imagery.
3. **Scale and Resolution:** The scale and resolution of the imagery influence the ability to identify and classify smaller or intricate features accurately.

Image classification in remote sensing is a powerful tool with diverse applications, providing valuable insights for sustainable agriculture, effective land management, and informed decision-making across various sectors. Advances in technology and machine learning continue to enhance the accuracy and efficiency of image classification methodologies, contributing to the evolving field of remote sensing applications

Classification of images

There are mainly two types of techniques which are used for classifying the images. They are 1) Supervised classification and 2) Unsupervised classification

Supervised classification

In this type of classification, the user selects the pixels which represent specific classes for sample training and then uses the image processing software to utilize the training data as reference for classification of other similar pixels in the image. The training points are selected based on the user's knowledge. The user also decides the bounds for the range of similarity of pixels for grouping. The bounds depend upon the spectral characteristics of the training data and may include pixels with some deviations in brightness or reflection strength. The number of classes in an image is also decided by the user.

The procedure for supervised classification is given below:

Step 1: Selection of training areas

A polygon is drawn on the area of interest and the training samples for various land cover classes are created.

ISBN : 9788119821327

For example: A polygon is drawn oven an urban area consisting of road or a parking lot. Several such polygons are made to represent the urban areas. Once enough training samples have been created, then another class is selected. Similar polygons are drawn over forest area, water body, agricultural land etc. In ArcGIS, the image analysis toolbar should be opened and the training sample manager is added. "Draw polygon" is clicked to add the training samples

If you're using ArcGIS, the following steps are followed:

1) Enable the image analysis toolbar
2) Add the training sample manager
3) Click on "Draw polygon"
4) Draw polygons for each land cover class
5) Merge the polygons with similar pixel information into classes




Fig.1: Draw polygon in the image and define the classes

Step 2: Generate signature file

The training samples for each class are already created. The signature file holds the training sample data and they are saved for using it later. The signature file is created using a "Create a signature file"

ISBN : 9788119821327




Fig. 5.2: Selection of signature file

Step 3: Classification

There are some common methods of supervised classification like maximum likelihood, iso-cluster, class probability, principal components and support vector machine. Previous studies have shown that support vector machine achieves better accuracy compared to other methods. The input file used in classification is the signature file.

Unsupervised classification

In this type of classification, the pixels having common characteristics are grouped by using software analysis of a satellite image without any guidance from the user. The user does not provide any training or sample classes for the software. The software carries out the analysis to determine the related pixels and then groups them into classes. The selection of algorithm and the number of classes are decided by the user but other than that, there is no aid provided by the user. The users can verify the classification by comparing it with the real features and evaluate the accuracy of the classification.

The steps for conducting an unsupervised classification are:

1. Generate clusters
2. Assign classes

Step 1: Generation of clusters

The pixels are clustered into a set number of classes by the software. The number of classes in which the image is to be classified is assigned by the user. The user also selects the appropriate band.

The steps for generating clusters in ArcGIS are given below:

ISBN : 9788119821327

1) The spatial analyst extension is to be activated
(Customize → Extensions → Spatial Analyst)
2) The iso-clusters can be used for unsupervised classification
(Spatial analysis tools → Multivariate → Iso clusters)
Input: The input is the image that is to be classified
Number of classes: The users can specify the number of classes. If a multispectral imagery (Red, green, blue and NIR bands) is being used for unsupervised classification, then the number of classes can be 40.
A cluster is created on clicking OK based on the input parameters.

Step 2: Assigning of classes

The user identifies the land cover classes of the clusters that are formed. The classes can be given various colours. For example: Blue colour can be given to water, green colour for vegetation etc. The classes can then be merged using the reclassify tool.

Landsat 8 Bands and Band combinations

Operational Land Imager (OLI) and Thermal Infrared Sensor (TIRS) are the two main sensors for Landsat 8. The OLI is capable of producing 9 spectral bands at 15 m and 30 m resolution. The 8 bands namely coastal, blue, green, red, NIR, SWIR -1, SWIR – 2 and cirrus have a spectral resolution of 30 metres. The panchromatic band has a finer resolution of 30 m. The TIRS consists of 2 thermal bands with a spatial resolution of 100 metres.

The bands can be arranged in such a way that a unique information can be obtained. The spectral signatures of objects are extracted for various bands. The natural colour combinations are popular in Landsat-8. The band combinations of Landsat-8 can be obtained from Landsat explorer. The bands, description, wavelength and resolution of Landsat 8 are provided in Table 5.1.

ISBN : 9788119821327

Table 5.1: Bands of Lansat-8

Band No.	Description	Wavelength	Resolution
Band 1	Coastal / Aerosol	0.433 to 0453 µm	30 meter
Band 2	Visible blue	0.450 to 0.515 µm	30 meter
Band3	Visible green	0.525 to 0.600 µm	30 meter
Band 4	Visible red	0.630 to 0.680 µm	30 meter
Band 5	Near-infrared	0.845 to 0.885 µm	30 meter
Band 6	Short wavelength infrared	1.56 to 1.66 µm	30 meter
Band 7	Short wavelength infrared	2.10 to 2.30 µm	60 meter
Band 8	Panchromatic	0.50 to 0.68 µm	15 meter
Band 9	Cirrus	1.36 to 1.39 µm	30 meter
Band 10	Long wavelength infrared	10.3 to 11.3 µm	100 meter
Band 11	Long wavelength infrared	11.5 to 12.5 µm	100 meter

Some examples of common band combinations and their examples are as follows:

Natural Colour (4, 3, 2)



Fig. 2: Natural colour composite image

The band combination of red (4), green (3) and blue (2) are used in the natural colour composite image. It resembles the vision of the human eye. The healthy vegetation looks green, unhealthy vegetation looks brows, urban features appear grey and white and the water body appears to be dark blue.

ISBN : 9788119821327

Colour infrared (5, 4, 3)




Fig. 3: Infrared colour image

The near-infrared (NIR) composite image uses near-infrared (5), red (4) and green (3) bands. This band is useful for analysis of vegetation as the chlorophyll reflects NIR light. The areas which appear red have better health of vegetation. The urban areas appear white and the water appears dark in this image.

Short wave infrared (7, 6, 4)



Fig. 5.3: Short wave infrared colour image

The short-wave infrared band combination includes SWIR-2 (7), SWIR-1 (6) and red (4). The vegetation appears in green shade in this composite image. The dark green colour indicates dense vegetation and light green colour indicates sparse vegetation. The urban areas appear blue.

ISBN : 9788119821327

Agriculture (6, 5, 2)




Fig. 4: Agriculture image

The band combination includes SWIR-1 (6), near-infrared (5) and blue (2). This band combination is generally used for the purpose of monitoring of crops. The healthy vegetation looks dark green and the bare earth appears to have magenta hue.

Acreage estimation

Acreage estimation through Remote Sensing (RS) and Geographic Information Systems (GIS) involves a comprehensive process that integrates advanced technologies to accurately determine the extent of agricultural land. The process begins with the acquisition of high-resolution satellite or aerial imagery, capturing detailed information about land cover and crop types. To ensure accuracy, preprocessing steps, including calibration, atmospheric correction, and geometric correction, are applied to the remote sensing data. Image classification algorithms are then employed to categorize land cover and identify different crop types, utilizing either supervised or unsupervised methods. The classified imagery is seamlessly integrated into GIS datasets, which include information such as land parcel boundaries and soil types. Spatial analysis, particularly overlay analysis, is conducted using GIS tools to combine and analyze different layers of information, facilitating accurate acreage estimation.

ISBN : 9788119821327

The critical step of acreage calculation involves overlaying classified land cover maps with cadastral or administrative boundary layers within the GIS environment. GIS tools are employed to compute area statistics for each land cover class, providing precise acreage estimates. Validation and accuracy assessment are crucial components, involving the comparison of estimated acreage with ground truth data obtained through field surveys. This process ensures the reliability and quality of the RS and GIS-derived results. The applications of this integrated approach are extensive, supporting precision agriculture by optimizing resource use, aiding in crop monitoring for better yield predictions, contributing to land use planning, and assisting insurance companies in risk assessment. Despite challenges related to data resolution, cloud cover, and ground truth verification, the integration of RS and GIS remains an indispensable tool in modern agriculture and land management, providing valuable insights for informed decision-making at various levels. Advances in technology continue to refine and enhance the accuracy of acreage estimation methods, reinforcing the significance of this integrated approach in contemporary agricultural practices.

The conventional approach of estimating the area of crop is costly and time consuming. The task of estimating crop is area is difficult to accomplish in hilly regions and several land types. Therefore, in order to overcome the difficulty of estimating area using conventional method, the remote sensing approach is adopted. Remote sensing provides the ability to classify the crops and also to estimate area and yield.

There are various methods of crop area estimation. They are as follows:

Direct estimation or pixel counting

The ground truth data of samples are required in this approach. The image is classified based on the ground truth data. The number of pixels under each crop is identified and this number is multiplied by the

ISBN : 9788119821327

size of the pixel for obtaining the crop area. If a sample segment is selected, the area under crop in that segment is estimated.

Multi-date data analysis




Fig. 5: Weekly/fortnightly composite NDVI images

The variations in phenology are utilized to identify various crops. The products having moderate spatial resolutions and high temporal frequency are used. Data of around 10 days in the growing period of crops are required to classify them. NDVI is useful in crop classification based on the crop cycle. As shown in Fig. 5.6, the NDVI

ISBN : 9788119821327

value for potato is high near the end of December. For mustard, the NDVI value starts decreasing towards the end of February. NDVI keeps increasing for wheat as it approaches February. These temporal variations in the values of NDVI are utilized to classify the crops.




Fig. 6: Temporal NDVI profile of various crops

Ground truth data

For validation of the classification, it is important to obtain data about crops by ground truth. The data included in ground truth for acreage estimation are geographical location, village, district, state, crop, coverage, condition, stage etc. It is also recommended to take pictures of the field. Sample locations are selected across the study region and it is ensured that almost all the diversity is covered. There are many apps available on android for field data collection which can be utilized to store and transmit the data.

Regression estimator

This method combines the information obtained from satellite images with information of sample obtained from ground survey.

ISBN : 9788119821327

$$\hat{Y}_R = \sum_{i=1}^{L} N_h \bar{y}_{h(Reg)}$$

$$\bar{y}_{h(Reg)} = \bar{y}_h + \hat{b}_h (\bar{X}_h - \bar{x}_h)$$

$\bar{y}_h$ = average ground-reported crop area per sample segment of stratum h, that is

$$\bar{y}_h = \frac{1}{n_h} \sum_{j=1}^{n_h} y_{hj}$$

$\hat{b}_h$= regression coefficient of ground-reported area on remote sensing-derived

$\bar{X}_h$= average remote-sensing-based area for all frame units of stratum h

Regression estimator is used my countries like USA, Brazil and China to estimate the crop from the remote sensing data. The area frame is developed using satellite imagery having spatial resolution of 10 m. The agricultural land is stratified and the random segments are interpreted visually using the ground information to estimate the crop statistics based on the area frame. The training sample consists of the visual interpretation of satellite imagery in the algorithm of supervised classification to extract pixel based crop estimates. The coefficient of determination in the linear regression between crop pixels and crop area is obtained.

References

1. QGIS.org, %Y. QGIS 3.28. Geographic Information System User Guide. QGIS Association.
2. Zaki, A., Buchori, I., Sejati, A. W., & Liu, Y. (2022). An object-based image analysis in QGIS for image classification and assessment of

ISBN : 9788119821327

coastal spatial planning. *The Egyptian Journal of Remote Sensing and Space Science*, *25*(2), 349-359.

3. Congedo, L. (2021). Semi-Automatic Classification Plugin: A Python tool for the download and processing of remote sensing images in QGIS. *Journal of Open Source Software*, *6*(64), 3172.
4. Maurya, A. K. (2011, January). Estimation of acreage & crop production through Remote Sensing & GIS. In *Geospatial World Forum, Hyderabad, India*.
5. Kamaraj, M., & Rangarajan, S. (2022). Predicting the future land use and land cover changes for Bhavani basin, Tamil Nadu, India, using QGIS MOLUSCE plugin. *Environmental Science and Pollution Research*, *29*(57), 86337-86348.

ISBN : 9788119821327

CHAPTER - 14

Application of RS, GIS and GPS in Agronomy

Tanveer Ahmad Ahngar[1]*, Ruby Jan[1] and Zahida Rashid[2]

[1]Division of Agronomy, Faculty of Agriculture, Wadura, SKUAST-K, Jammu and Kashmir, India

[2]Dryland Agriculture Research Station, Rangreth, SKUAST-K, Jammu and Kashmir, India.

Corresponding author: tanveeragron@gmail.com

Abstract

This chapter explores the exciting ways in which modern technologies like Geographic Information System (GIS), Global Positioning System (GPS), and Remote Sensing (RS) are revolutionizing farming practices. We take a closer look at how GIS helps farmers make smart decisions about where to plant crops and how to take care of the soil. GPS acts like a farming wizard, guiding tractors and machines with precision, making farming more efficient. Additionally, RS, like a high-tech eye in the sky, helps farmers keep an eye on the health of crops and spot potential issues. Through real-life examples, this chapter showcases how these technologies are making farming not just easier, but also more sustainable. Despite the benefits, we acknowledge that challenges like getting accurate data and convincing everyone to use these tools exist. However, the chapter ends optimistically, discussing upcoming advancements that promise an even brighter future for smart farming through GIS, GPS, and RS applications in agronomy.

Introduction

Agronomy, a branch of agriculture, serves as a foundational activity supporting the economies of numerous nations. Its substantial contributions to employment and the gross domestic product,

ISBN : 9788119821327

particularly in developing countries, make it a crucial aspect of national prosperity. However, the challenge of increasing food production to meet the needs of a growing population, coupled with the finite nature of land resources, poses a significant dilemma for many developing nations. Consequently, improving agronomic practices becomes imperative to ensure prudent land-use planning and efficient resource management in crop cultivation.

With a rising interest in incorporating site-specific information into a spatial and long-term perspective, achieving precision in agronomic practices requires advanced technology capable of calculating spatial and temporal variations in crop growth. Presently, remarkable progress has been made in developing digital systems for applications in soil fertility examination, soil survey, land-use planning, crop production, and yield monitoring. Computer programs, such as Geographic Information System (GIS), (GPS) and Remote Sensing (RS) play a pivotal role in enhancing the speed and efficiency of overall agronomic planning processes.

Remote Sensing and its Applications

Remote sensing encompasses both the art and science of acquiring information about objects or areas in the real world from a distance, eliminating the need for direct physical contact with the subjects under investigation. Employing space technologies alongside ground observations, remote sensing serves as a tool to monitor Earth's resources, offering enhanced precision and accuracy. The underlying principle involves utilizing the electromagnetic spectrum, including visible, infrared, and microwaves, to evaluate the Earth's features. Different responses of targets to these wavelength regions enable the differentiation of vegetation, bare soil, water, and other similar features. Remote sensing finds application in various fields, such as crop growth monitoring, tracking land use patterns and changes in land cover, mapping water resources, assessing water status under field conditions,

ISBN : 9788119821327

monitoring diseases and pest infestations, forecasting harvest dates, estimating yields, enabling precision farming, and supporting weather forecasting, all complemented by ground observations. Essentially, remote sensing techniques play a crucial role in sensing Earth's resources. The data obtained through remote sensing significantly contributes to monitoring Earth's surface features by providing timely, comprehensive, cost-effective, and repetitive information (Justice et al., 2002). Furthermore, remote sensing holds numerous applications in agrometeorology. When combined with crop simulation models, remote sensing inputs become invaluable for crop yield forecasting. Agricultural applications have evolved in satellite remote sensing, initially focusing on land cover classification, especially crop types. However, recent advancements emphasize characterizing plant biophysical properties. Remote sensing proves invaluable in monitoring and analyzing agricultural activities, offering insights into agronomic parameters without destructive crop sampling. The technology's non-invasive nature provides repeated information for precision agriculture, offering a cost-effective alternative over large geographical areas.

In India, satellite remote sensing primarily aids in crop acreage and production estimation. The technology has transformative potential in detecting and characterizing agricultural productivity based on crop and soil biophysical attributes. Remote sensing, coupled with GIS, creates spatiotemporal informative layers applicable in floodplain mapping, hydrological modelling, surface energy flux, urban development, land use changes, crop growth monitoring, and stress detection.

Monitoring vegetation cover is a key focus, where hyperspectral data enhances crop classification accuracy. Various vegetation indices, such as NDVI, VCI, LAI, GYURI, and TCI, contribute to mapping and monitoring drought, assessing vegetation health, and predicting productivity. Additionally, remote sensing aids in assessing nutrient and water status, essential for precision farming, allowing efficient resource use.

ISBN : 9788119821327

Evapo-transpiration estimation is critical for irrigation scheduling, water and energy balance computations, and crop water stress index determination. Remote sensing, especially with hyper-spectral sensors, linked to GIS and GPS technologies, plays a significant role in water management for agricultural systems.

Weed identification and management benefit from remote sensing, offering a non-time-consuming alternative to ground surveying methods. The technology aids in developing weed maps, allowing site-specific herbicide application.

Remote sensing becomes essential in monitoring and quantifying crop stress due to pests and diseases. It provides data for pest management decision support systems and aids in forecasting crop yields based on statistical-empirical relationships between yield and vegetation indices.

Finally, remote sensing is a key component of precision farming, supporting variable rate technology and enabling informed decision-making for reduced cultivation costs, improved control, and enhanced resource use efficiency. Meteorological satellites contribute to weather forecasting, with applications such as drought assessment playing a pivotal role in agriculture.

Remote sensing holds numerous advantages in agronomic research. The evaluation of crop canopies using remote sensing provides valuable insights into various agronomic parameters. Its role is particularly significant in tasks such as crop classification, monitoring, and yield assessment. In agronomic research, the utilization of remote sensing is essential due to the susceptibility of crops to variations in soil conditions, climate, and other physico-chemical changes. Agricultural production systems exhibit strong seasonal patterns aligned with the biological life cycle of crops, which are highly variable in space and time dimensions. Additionally, agricultural productivity can change rapidly due to unfavorable growing conditions, necessitating timely monitoring.

ISBN : 9788119821327

For effective and sustainable agricultural management, it is crucial to analyze factors influencing the agricultural sector on a spatiotemporal basis. Remote sensing, alongside advanced technologies like global positioning systems and geographical information systems, plays a pivotal role in assessing and managing agricultural activities. These technologies have diverse applications in agriculture, including crop acreage estimation, monitoring crop growth, estimating soil moisture and fertility, detecting crop stress, identifying diseases and pest infestations, monitoring drought and flood conditions, estimating yields, and contributing to weather forecasting. The implementation of precision agriculture, facilitated by these technologies, is essential for maintaining agricultural sustainability and fostering economic growth.

Geophysical surveys have enabled the exploration of detailed information beneath the Earth's surface. Remote sensing, with its advantages over traditional techniques, offers enhanced spatial, spectral, radiometric, and temporal data availability. Its capability to acquire real- or near-real-time data, even from remote or inaccessible areas, within a short timeframe makes it an efficient and powerful tool for the assessment, exploration, evaluation, analysis, monitoring, and management of groundwater, leading to long-term societal benefits.

The field of remote sensing in groundwater hydrology is dynamic, evolving rapidly with advancements in sensors, platforms, and data processing algorithms. These developments support the generation of new forms of data, offering novel perspectives of landscapes. Hydrologists and hydraulic engineers, armed with these technological advancements, can better evaluate the Earth's surface and specific features (Running et al. 1994).

Future prospects

Remote sensing proves highly valuable in assessing both abiotic and biotic stresses across various crops, and it is equally effective in identifying and managing crop issues, even on small-scale farm

ISBN : 9788119821327

holdings. To harness the information on crops for economic improvement, there is a necessity to establish state or district-level information systems based on data derived from remote sensing and GIS methodologies.

Governments can leverage remote sensing data to make critical decisions regarding policies and strategies addressing national agricultural issues. A novel and nontraditional application of remote sensing involves the incorporation of nano-chips into plant and seed tissue. These nano-chips enable near-real-time monitoring of crops. Clearly, these innovative approaches, along with others, underscore the continued significance of remote sensing in the future analysis of agricultural sciences.

GIS and its Applications

According to experts, process-based agronomic models primarily analyze temporal variations using point data from specific sites, while GIS facilitates the storage, manipulation, analysis, and visualization of data. The synergy of both spatial and temporal considerations is best addressed by integrating agronomic models with GIS. A GIS, functioning as a thematic mapping system, enables the production of maps based on various themes, such as soils or hydrology. It is a specialized information system that keeps track of events, activities, and things, including their spatial occurrence.

GIS, as part of a suite of technologies enhancing precision in agronomic practices, relies on preliminary basic information relevant to the specific project discipline. Importing information into GIS demands time and attention, as it forms the foundational knowledge of the territory and individual parameters that is challenging to modify later. GIS processes and links information simultaneously, offering a syntactical expression of system changes induced by parameter variations. This dynamic system allows for the rapid adaptation to real conditions and provides answers in near real-time.

ISBN : 9788119821327

Over the past three decades, GIS techniques have been utilized for farm-related assessments at national and regional scales, with widespread applications in agronomic and natural resource management over the last decade. GIS, being a dynamic product, is easy to update, edit, and reproduce maps, providing valuable support in handling voluminous data generated through conventional and spatial formats. Unlike paper maps, GIS can prepare and manage large collections of agronomic and land resource data essential for crop production.

The importance of GIS in agronomy is underscored by its ability to study the nutrient status of individual fields, leading to specific requirements for external nutrient application. In precision agronomic practices, GIS manages the information-intensive environment in crop production by combining site-specific management with computer software modelling for analyzing and interpreting varying inputs and outputs. GIS facilitates within-field variable rate application, addressing spatial variation in crop yields within a field and enhancing the assessment and understanding of variations in field crops.

GIS applications in agronomy have expanded from hydrological applications in the mid-1980s to a wide range of areas, including atmospheric modelling, climate change studies, characterization and zonation, hydrology, water quality assessment, soil science, and spatial yield calculation. Various studies have reported GIS applications in cultivation practices for different crops, including fertility management for tea cultivation, soil quality monitoring, and impact assessments on crop yield and nitrate movement.

In conclusion, GIS serves as an excellent informative tool that enhances the visualization, analysis, and handling of spatial data. Its digital mapping capabilities allow for the periodic review and improvement of soil fertility status, providing updated information on crops, soil, and prevailing climatic conditions. This accurate knowledge

ISBN : 9788119821327

base significantly enhances the management of finite resources and supports accurate land-use planning.

Global Positioning System (GPS):

The Global Positioning System (GPS) is a satellite-based navigation system that allows users to determine their precise location and track movement anywhere on Earth. It consists of a network of satellites orbiting the planet, ground-based control stations, and GPS receivers. The GPS receiver calculates its location by triangulating signals received from multiple satellites, providing accurate position information, including latitude, longitude, and elevation.

Applications of GPS in Agronomy:

1. **Precision Farming:**
   - ***Variable Rate Technology (VRT):*** GPS enables precise application of resources such as seeds, fertilizers, and pesticides at variable rates based on the specific needs of different areas within a field.
   - ***Guidance Systems:*** GPS-guided tractors and machinery ensure accurate and efficient field operations, reducing overlap and optimizing planting, cultivating, and harvesting.
2. **Field Mapping and Surveying:**
   - ***Boundary Mapping:*** GPS helps in accurately mapping field boundaries, aiding in land management and property delineation.
   - ***Topographic Mapping:*** GPS assists in creating detailed topographic maps of fields, providing insights into terrain variations that impact water drainage and overall field management.
3. **Soil Sampling:**
   - ***Grid Sampling:*** GPS is used to create grid patterns for soil sampling, ensuring that samples are collected systematically from specific locations. This helps in understanding soil fertility variations across the field.

ISBN : 9788119821327

4. **Yield Monitoring:**
    - ***Harvest Data Collection:*** GPS technology is employed in yield monitoring systems, recording and geo-tagging crop yield data during harvesting. This data helps farmers analyze yield variability and make informed decisions for future seasons.
5. **Irrigation Management:**
    - ***Precision Irrigation:*** GPS is utilized to precisely map and manage irrigation activities based on the specific needs of different areas within a field, optimizing water use efficiency.
6. **Crop Scouting:**
    - ***Remote Sensing Integration:*** GPS coordinates are integrated with remote sensing technologies to identify and monitor crop health. Farmers can scout fields more efficiently, targeting specific areas for intervention.
7. **Equipment Tracking and Management:**
    - ***Fleet Management:*** GPS helps track the location and movement of farm equipment, facilitating efficient fleet management, reducing downtime, and improving operational logistics.
8. **Record-Keeping:**
    - ***Farm Management Software:*** GPS data is integrated into farm management software, providing a comprehensive record of field activities, input applications, and yield data. This historical information aids decision-making for subsequent growing seasons.
9. **Boundary and Buffer Management:**
    - ***Environmental Compliance:*** GPS is used to establish and manage buffer zones around fields, ensuring compliance with environmental regulations and minimizing the impact of agricultural activities on adjacent areas.

ISBN : 9788119821327

10. **Research and Experimentation:**
    - ***Field Trials:*** GPS technology enables precise location tagging for experimental plots, making it easier to monitor and analyze the results of various agronomic experiments.

The integration of GPS technology in agronomy enhances precision, efficiency, and sustainability in farming practices.

Integration of RS, GIS and GPS

The integration of Remote Sensing (RS), Geographic Information Systems (GIS), and Global Positioning System (GPS), collectively known as 3S, has garnered significant attention in the realm of groundwater hydrology in recent years. Groundwater, a crucial subterranean natural resource, presents a complex array of dimensions. Remote sensing, a widely used technique employing various platforms such as aircraft and satellites, has proven invaluable in enhancing our understanding of subsurface water conditions (Barrett and Kidd, 1987). This encompasses geophysical surveys encompassing gravity, magnetics, and electromagnetics, providing insights into the Earth's subsurface.

Challenges and Resolutions in Implementing RS, GIS, and GPS Technologies in Agronomy

1. **Data Integration:**
    - ***Challenge:*** Integrating data from RS, GIS, and GPS sources can be complex due to differences in data formats, scales, and resolutions.
    - ***Solution:*** Develop standardized protocols and interoperable formats to facilitate seamless integration of data. Use middleware or platforms that support multi-source data integration.

ISBN : 9788119821327

2. **Data Accuracy and Resolution:**
    - *Challenge:* RS and GPS data may have varying levels of accuracy and resolution, affecting the precision of agronomic analyses.
    - *Solution:* Invest in high-resolution RS and GPS equipment. Implement ground control points for accurate georeferencing. Regularly update and calibrate GPS equipment.
3. **Cost of Technology:**
    - *Challenge:* Acquiring and maintaining RS, GIS, and GPS technologies can be expensive for small-scale farmers.
    - *Solution:* Promote collaborative initiatives, subsidize technology costs, and provide training to make advanced technologies more accessible. Encourage sharing of equipment among farmers.
4. **Data Privacy and Security:**
    - *Challenge:* Protecting sensitive agronomic data, such as farm layouts and yield information, from unauthorized access is a concern.
    - *Solution:* Implement secure data storage protocols, encryption, and access controls. Educate users on data privacy practices. Adhere to legal regulations governing data security.
5. **Technical Expertise:**
    - *Challenge:* Many farmers may lack the technical expertise required to effectively use RS, GIS, and GPS technologies.
    - *Solution:* Provide training programs, workshops, and user-friendly interfaces. Collaborate with agricultural extension services to disseminate knowledge on technology usage.
6. **Infrastructure Limitations:**
    - *Challenge:* In remote or underdeveloped areas, limited infrastructure may hinder the implementation of RS, GIS, and GPS technologies.
    - *Solution:* Invest in improving infrastructure, including connectivity and power supply. Explore alternative technologies that operate in low-infrastructure environments.

ISBN : 9788119821327

7. **Data Overload:**
    - *Challenge:* The abundance of data generated by RS, GIS, and GPS can overwhelm users, making it challenging to extract meaningful insights.
    - *Solution:* Implement data filtering algorithms, data reduction techniques, and visualization tools to simplify data interpretation. Provide training on effective data management practices.
8. **Regulatory Compliance:**
    - *Challenge:* Adhering to regulations related to land use, environmental protection, and privacy can be challenging when using RS, GIS, and GPS technologies.
    - *Solution:* Stay informed about relevant regulations, obtain necessary permits, and collaborate with regulatory authorities. Develop transparent practices for data collection and use.
9. **Limited Connectivity in the Field:**
    - *Challenge:* Field locations may lack reliable network connectivity, hindering real-time data transmission and communication.
    - *Solution:* Use offline-capable GPS devices. Implement data synchronization when connectivity is available. Explore satellite-based communication solutions.
10. **Resistance to Change:**
    - *Challenge:* Farmers and agronomists may resist adopting new technologies due to unfamiliarity.
    - *Solution:* Conduct awareness campaigns, showcase success stories, and provide incentives for early adopters. Emphasize the benefits of improved efficiency, cost-effectiveness, and sustainability.

Overcoming these challenges requires a collaborative effort involving government agencies, technology providers, researchers, and farmers to ensure the effective and widespread adoption of RS, GIS, and GPS technologies in agronomy. The rapid evolution of technology

ISBN : 9788119821327

continues to shape the future of agronomy, with RS, GIS, and GPS playing pivotal roles in transforming agricultural practices. Anticipating future trends is crucial for stakeholders in the agronomic field. This section explores emerging trends in RS, GIS, and GPS technologies, offering insights into their potential impact on the future of agronomy.

1. Integration of Artificial Intelligence (AI) and Machine Learning (ML):

- The integration of AI and ML into RS, GIS, and GPS applications will enhance data analysis, providing more accurate predictions and actionable insights for agronomic decision-making.

2. Advancements in Hyperspectral and LiDAR Technologies:

- Continued advancements in hyperspectral and LiDAR sensors will enable more detailed and precise data acquisition, allowing agronomists to gain in-depth insights into crop health, soil conditions, and vegetation characteristics.

3. Enhanced Spatial and Temporal Resolution:

- Future developments are expected to yield sensors with improved spatial and temporal resolutions. This will enable more frequent and detailed monitoring of agricultural landscapes, facilitating timely interventions.

4. Internet of Things (IoT) Integration:

- Integrating IoT devices with RS, GIS, and GPS technologies will enable real-time data collection from sensors deployed in the field. This connectivity will enhance monitoring and management capabilities in agronomy.

5. Autonomous Farming Systems:

- The rise of autonomous vehicles and machinery equipped with GPS technologies will revolutionize farming practices. These systems will enable precise and efficient resource management, reducing labor requirements and enhancing overall productivity.

ISBN : 9788119821327

6. Cloud-Based Solutions for Data Storage and Analysis:

- Cloud computing will become increasingly integral, providing scalable and secure platforms for storing and analyzing large volumes of agronomic data. This shift to cloud-based solutions will enhance collaboration and accessibility.

7. Blockchain for Traceability and Transparency:

- Blockchain technology will find applications in ensuring traceability and transparency in the agricultural supply chain. This will be particularly valuable for quality assurance and compliance with industry standards.

8. Climate-Smart Agriculture Solutions:

- RS, GIS, and GPS technologies will be pivotal in developing climate-smart agriculture solutions. These technologies will assist in adapting agricultural practices to changing climate conditions, optimizing resource utilization, and promoting sustainability. The future trends in RS, GIS, and GPS for agronomy suggest a trajectory towards more sophisticated, interconnected, and data-driven approaches. Embracing these advancements will empower agronomists to make informed decisions, enhance productivity, and contribute to sustainable agricultural practices.

Conclusion:

In conclusion, the integration of Remote Sensing (RS), Geographic Information Systems (GIS), and Global Positioning System (GPS) technologies has significantly transformed the landscape of agronomy, offering multifaceted solutions to complex challenges. The applications of these technologies in agronomy have demonstrated their potential to enhance precision, efficiency, and sustainability in agricultural practices. The applications of RS have proven instrumental in providing timely and accurate information for crop monitoring, disease detection, and yield estimation. The ability to capture real-time

ISBN : 9788119821327

data from various platforms, including satellites and aircraft, empowers agronomists with a comprehensive understanding of crop health, growth patterns, and environmental conditions. GIS plays a pivotal role in spatial analysis and decision-making, offering a powerful tool for mapping, monitoring, and managing agricultural landscapes. The spatial insights derived from GIS applications enable agronomy professionals to optimize land use, plan irrigation systems, and assess the impact of environmental factors on crop performance. The advent of GPS technology has revolutionized precision agriculture, allowing for precise navigation, mapping, and data collection in the field. The accurate geospatial information provided by GPS facilitates variable-rate applications, ensuring targeted resource utilization, minimizing waste, and optimizing overall crop yield. The synergy of RS, GIS, and GPS technologies has also contributed to sustainable agriculture by enabling better natural resource management. From soil fertility evaluation to water resource mapping, the 3S technologies empower agronomists to make informed decisions that align with long-term environmental conservation goals. Despite the numerous benefits, challenges such as data integration, technical expertise, and cost implications need to be addressed for widespread adoption. Continuous advancements in technology, coupled with collaborative efforts among researchers, policymakers, and practitioners, will pave the way for further innovations and improvements in the applications of RS, GIS, and GPS in agronomy.

As we look to the future, the ongoing refinement of these technologies, coupled with a commitment to knowledge sharing and capacity building, promises a continued evolution of precision agriculture. The journey from data acquisition to actionable insights will play a crucial role in shaping a more resilient, sustainable, and productive agronomic landscape. With each technological stride, the applications of RS, GIS, and GPS in agronomy will undoubtedly play a pivotal role in addressing

ISBN : 9788119821327

the dynamic challenges of feeding a growing global population while promoting environmental stewardship.

References

1. Barrett, E. C., and Kidd, C. (1987). The use of SMMR data in support of a VIR/IR satellite rainfall monitoring technique in highly contrasting climatic environments. In *passive microwave observing from environmental satellites, a status report* (pp. 109-123). Washington, DC: NOAA Tech. Rep. NESDIS 35.
2. Justice, C. O., Townshend, J. R. G., Vermote, E. F., Masuoka, E., Wolfe, R. E., Saleous, N., and Morisette, J. T. (2002). An overview of MODIS Land data processing and product status. *Remote sensing of Environment*, *83*(1-2), 3-15.
3. Running, S. W., Justice, C. O., Salomonson, V., Hall, D., Barker, J., Kaufmann, Y. J., and Carneggie, D. (1994). Terrestrial remote sensing science and algorithms planned for EOS/MODIS. *International journal of remote sensing*, *15*(17), 3587-3620.

ISBN : 9788119821327

CHAPTER - 15

Geospatial Technologies for Crops and Soils

Seerat Jan [1], Tanveer A. Ahanger[1*] and Shahid Rasool [2]

[1]Division of Agronomy, Faculty of Agriculture, Wadura, SKUAST-K, Jammu and Kashmir, India

[2]CSIR-Indian Institute of Integrative Medicine (CSIR-IIIM)

Corresponding author: tanveeragron@gmail.com

Abstract

Geospatial technologies have emerged as powerful tools in modern agriculture, playing a transformative role in enhancing precision, efficiency, and sustainability. This book chapter provides an overview of the applications and impact of geospatial technologies in the context of crops and soils. From precision farming to soil mapping and crop monitoring, these technologies offer a suite of tools that enable farmers to make data-driven decisions, ultimately leading to optimized resource management and increased agricultural productivity. The literature delves into the key components of geospatial technologies, including Geographic Information Systems (GIS), Global Positioning Systems (GPS), and remote sensing. These technologies facilitate precise data collection, analysis, and interpretation of spatial information related to crops and soils. Precision farming, a cornerstone application, involves leveraging detailed data about soil characteristics, moisture levels, and topography to make informed decisions on planting, fertilization, irrigation, and harvesting. Soil mapping, another critical aspect, enables farmers to analyze the chemical and physical properties of the soil, aiding in crop selection, fertilizer application, and irrigation scheduling.

ISBN : 9788119821327

Crop monitoring, utilizing remote sensing tools such as drones and satellite imagery, allows real-time tracking of crop health, early detection of stress factors, and informed pest and disease management. The abstract also addresses the challenges associated with geospatial technologies in agriculture, including limited access, high costs, and the need for technical knowledge. Furthermore, it emphasizes the importance of collaborative efforts involving government, the private sector, and academic institutions to overcome these challenges and promote research, development, and knowledge sharing. In conclusion, this book chapter underscores the transformative impact of geospatial technologies on modern agriculture, offering a pathway to sustainable and precision-driven farming practices. The integration of these technologies not only empowers farmers with valuable insights but also contributes to the global goals of ensuring food security and environmental sustainability.

Keywords: Agriculture, Crop monitoring, Geospatial, Geographic Information System, Information Technology, Remote Sensing

Introduction

Geospatial technology is a dynamic field focused on the collection, analysis, and interpretation of spatial data. It encompasses tools such as Geographic Information Systems (GIS), Global Positioning Systems (GPS), and Remote Sensing (RS). In India, geospatial technology has permeated various sectors, including agriculture, telecommunications, oil and gas, environmental management, forestry, public safety, infrastructure, and logistics. The geospatial industry is poised for substantial growth in the coming years, as stakeholders

ISBN : 9788119821327

recognize the value and long-term cost-effectiveness of adopting these technologies. With India's GDP projected to reach $9-10 trillion by 2025, the country is transitioning into a highly industrialized and technologically advanced economy. To support this economic expansion, there is a growing need for a modern information and knowledge regime. Geospatial technologies are expected to play a crucial role in information management in India, with significant social and national implications. States like Andhra Pradesh, Karnataka, Rajasthan, and Tamil Nadu have already incorporated geospatial technology to enhance governance and management. In agriculture, geospatial technology is increasingly vital for improving crop yields, reducing environmental impact, and enhancing sustainability. Various applications of geospatial technology in agriculture include precision farming, crop monitoring, weather forecasting, field mapping, and farm management. Precision farming involves leveraging detailed data on soil type, moisture levels, and topography to make informed decisions about planting, fertilization, irrigation, and harvesting. Crop monitoring utilizes remote sensing tools such as drones and satellite imagery to assess crop health and detect signs of stress or disease. Weather forecasting aids farmers in making informed decisions about crop management, leading to improved yields and reduced environmental impact. Field mapping, analysis, and equipment monitoring contribute to optimizing farm management practices and increasing productivity. Geospatial technology empowers farmers with valuable insights into their fields' physical characteristics, enabling data-driven decisions that promote efficient and sustainable agricultural practices to meet the growing demand for food while preserving natural resources.

ISBN : 9788119821327



Geospatial Technology in Agriculture

Geospatial technology plays a crucial role in agriculture, involving the utilization of tools and methodologies for the collection, analysis, and interpretation of spatial data to enhance farming practices. These technologies encompass Geographic Information Systems (GIS), Global Positioning Systems (GPS), remote sensing, and other related tools. The application of geospatial technology in agriculture has gained significance due to challenges such as climate change, land degradation, water scarcity, and food security. Farmers, grappling with these issues, can leverage geospatial technology to make more informed decisions regarding their crops and fields, ultimately resulting in improved yields, cost reduction, and increased sustainability. Widely adopted in agriculture, geospatial technology contributes to enhancing crop productivity, optimizing resource utilization, and minimizing environmental impacts. Various applications of geospatial technology in agriculture include the following (Esri, 2008).

ISBN : 9788119821327

Precision Agriculture

Precision agriculture represents an agricultural strategy that employs geospatial technology, data analytics, and advanced technologies to maximize crop productivity while mitigating environmental repercussions. This methodology revolves around the utilization of intricate data pertaining to a specific field's attributes, including soil type, moisture levels, and topography. The objective is to make well-informed decisions regarding planting, fertilization, irrigation, and harvesting processes.



Here are some key components of precision agriculture:

1. **Data Collection**: Data Collection in precision agriculture is heavily reliant on gathering and scrutinizing data from diverse sources, including satellite imagery, drones, and sensors. These tools furnish information about the physical attributes of a field, encompassing soil type, moisture levels, temperature, as well as insights into crop health and yield.

ISBN : 9788119821327

2. **Data analysis**: Following the data collection phase, analysis becomes paramount. Employing machine learning algorithms, patterns are discerned, and predictions are made. For instance, data on soil moisture and weather patterns can be harnessed to forecast optimal times for crop irrigation, aiming to maximize yield while minimizing water usage.
3. **Precision application**: Precision agriculture extends to the meticulous application of resources such as fertilizers, pesticides, and water. Specialized equipment is employed to precisely apply these resources based on the data collected and analyzed, ensuring targeted and efficient usage.
4. **Yield monitoring**: is a crucial aspect of precision agriculture. By systematically tracking crop yield and leveraging this data for insights, farmers can enhance future production. This process aids in pinpointing well-performing areas in the field and those requiring improvement, guiding subsequent planting and management decisions.

In essence, precision agriculture holds the potential to elevate crop yields, curtail expenses, and diminish environmental impact by empowering farmers to make informed, data-driven decisions regarding their field management practices.

Crop monitoring

Crop monitoring involves the systematic tracking of crop health and growth through the application of geospatial technology and remote sensing tools, including satellite imagery, drones, and sensors. This practice furnishes farmers with comprehensive insights into their

ISBN : 9788119821327

crops, enabling them to make w ell-informed decisions regarding fertilization, irrigation, and pest management.

Here are some ways crop monitoring can be used in agriculture



Early identification of plant stress: is a key benefit of crop monitoring, enabling farmers to detect issues like water or nutrient deficiencies before they are visibly apparent. This early detection empowers farmers to take corrective measures promptly, preventing any negative impact on crop yield.

Pest and disease management: crop monitoring proves valuable by aiding farmers in the detection and control of these issues. By utilizing remote sensing tools to monitor crop health, farmers can pinpoint areas affected by pests or diseases and implement targeted measures for effective control.

irrigation management: Effective irrigation management is another aspect where crop monitoring plays a crucial role. By offering insights into soil moisture content, crop monitoring assists farmers in

ISBN : 9788119821327


optimizing their irrigation schedules. This helps prevent both over- and under-watering, which can lead to reduced crop yield and the wastage of water resources.

yield forecasting: Additionally, crop monitoring serves as a tool for yield forecasting. By providing information about anticipated crop yields, farmers can make informed decisions regarding marketing and distribution.

In summary, crop monitoring equips farmers with valuable insights into their crops, enabling them to enhance yields, reduce input costs, and minimize environmental impacts by making well-informed decisions throughout the farming process.

Mapping

Mapping serves as a fundamental aspect of geospatial technology within the agricultural context. This practice entails the utilization of Geographic Information System (GIS) tools to gather and analyze data pertaining to the physical attributes of a field, encompassing soil type, topography, and water resources. Subsequently, this information is employed to make well-informed decisions regarding crop management activities, such as planting, fertilization, and irrigation.



ISBN : 9788119821327

Here are some ways mapping is used in agriculture:

Soil Mapping: The process of soil mapping entails the collection and analysis of data concerning the chemical and physical attributes of the soil within a specific field. This data aids farmer in making decisions related to crop selection, determining appropriate fertilizer quantities, and scheduling irrigation.

Topographic Mapping: Topographic mapping involves the creation of a comprehensive map detailing the terrain and features of a field, such as hills, valleys, and watercourses. This information assists farmers in deciding where to plant crops, addressing soil erosion concerns, and planning infrastructure development.

Water Resource Mapping: Water resource mapping encompasses the identification and mapping of water resources like rivers, streams, and groundwater. This information guides decisions on irrigation practices, crop selection, and overall land use.

Precision Farming: Mapping plays a pivotal role in precision agriculture, enabling farmers to make data-driven decisions regarding crop management. Precision farming utilizes detailed field characteristics, such as soil type, moisture levels, and topography, to inform decisions on planting, fertilizing, irrigating, and harvesting crops.

In summary, mapping serves as a crucial tool for farmers, delivering essential information about the physical aspects of a field that influences decisions concerning crop management, land utilization, and infrastructure planning

ISBN : 9788119821327

Weather forecasting

Weather forecasting plays a crucial role in agriculture as a vital application of geospatial technology. Offering precise weather information, this technology empowers farmers to make knowledgeable decisions regarding various aspects of crop management, encompassing activities such as planting, fertilization, irrigation, and pest control.



Here are some ways weather forecasting is used in agriculture

Planting Decisions: Farmers rely on weather forecasts to determine the optimal timing for planting their crops. This involves anticipating the last frost date and identifying periods when temperatures are favorable for the germination of plants.

Fertilization and Irrigation: Weather forecasts aid farmers in scheduling fertilization and irrigation activities. For instance, if rain is anticipated, farmers may opt to postpone irrigation to prevent overwatering.

Pest Control: Weather forecasts assist farmers in forecasting potential pest infestations. Understanding the preferences of pests, such as thriving in warm and humid conditions, allows farmers to implement preventive measures.

ISBN : 9788119821327

Harvest Planning: Weather forecasts are essential for harvest planning. Farmers need to be aware of expected rainfall to avoid harvesting crops in wet conditions, as damp crops are more susceptible to diseases and spoilage. In essence, weather forecasting is pivotal in agriculture, guiding farmers to make judicious decisions about crop management, resulting in enhanced yields and minimized environmental impact.

Farm management

Farm management involves overseeing agricultural operations to attain specific objectives, including heightened productivity, profitability, and sustainability. Geospatial technology plays a crucial role in this process by furnishing farmers with up-to-date information about their fields, crops, and equipment in real-time.



ISBN : 9788119821327

Here are some ways geospatial technology is used in farm management

Field Mapping and Analysis: Utilizing GIS tools, farmers can map their fields and analyze gathered data to identify soil types, slope, and other topographical features. This information aids in planning crop rotation, determining optimal planting patterns, and making decisions regarding nutrient and water management.

Crop Monitoring: Remote sensing tools like drones and satellite imagery enable farmers to monitor crop health and identify signs of stress or disease. This data is instrumental in making decisions about pest control and refining crop management practices.

Equipment Monitoring and Management: Sensors can be employed by farmers to monitor equipment, providing real-time data on performance. This information facilitates timely maintenance scheduling, minimizes downtime, and optimizes equipment usage.

Yield Forecasting and Analysis: By leveraging data from crop monitoring and field mapping, farmers can forecast crop yields and analyze trends over time. This information proves valuable for making informed decisions related to marketing, pricing, and strategic crop rotation.

Collection, analysis, and interpretation of geospatial data for agricultural purposes

Efficiently collecting and evaluating relevant data is imperative for the success of geospatial technology in agriculture. Various strategies have been developed, with a focus on remote sensing, to achieve this goal. The initial step in precision farming involves dividing extensive farmland into management zones, necessitating the use of remote sensing (Grisso, 2009). Each zone has distinct requirements

ISBN : 9788119821327

addressed through the utilization of GIS and GPS. The division into management zones is primarily based on factors such as soil types, pH levels, pest presence, nutrient availability, soil moisture content, fertility needs, weather predictions, crop characteristics, and hybrid responses. The information required for this division can be obtained from existing records, including soil survey maps, historical crop data, and documents reflecting regional cropping practices. Aerial and satellite images are also valuable in this process. An example is the examination of three characteristics using a Daedalus sensor aboard a NASA plane in an image captured on January 30, 2001. The information gleaned from assessing parameters like vegetation density, water distribution, and crop stress enables effective micromanagement of water administration, considering varying soil conditions and vegetation growth. Continuous aerial and satellite images can be generated at different times, providing the farmer with insights into the production of various management zones and detecting growth and yield trends. Various remote sensing techniques enhance the efficiency of this process, with image-based remote sensing data being a prevalent method. The relationship derived from comparing the reflection of red and near-infrared light forms the basis for evaluating productivity/yield. RADARSAT data has also proven beneficial, offering reliable information on soil conditions and crop performance. The data collected through remote sensing is transformed into spatial data through GIS techniques such as kriging. This spatial data enables the identification of potential issues in various management zones, empowering farmers to make informed decisions to address challenges and boost overall farm production. To ensure the relevance of collected point data, proper storage and analysis are crucial, requiring the use of

ISBN : 9788119821327

GIS tools. GIS software aids in creating digital maps that represent geospatial data digitally, either in raster or vector formats. The comparison of results with field notes helps identify trends and connections on the ground, providing insights into soil nutritional content or pest infestations. This information guides site-specific decisions on fertilizers, herbicides, insecticides, irrigation, and other aspects. Storing data systematically is essential for future reference, enhancing the efficiency of subsequent surveys. As demand grows, automated farm machinery becomes more prevalent, relying on GIS and GPS for accurate, safe, and efficient operations, ultimately increasing farm yield (Sohne et al., 1994).

Geospatial Technology's Impact on Agriculture

The adoption of modern information, geospatial, and communication technologies (ICTs) for rural and agricultural development has witnessed significant progress in the Asia-Pacific region over the past decade. Geospatial technology plays a pivotal role, particularly in the surveying and mapping of plantation crops. Notably, countries like Malaysia and Indonesia have undertaken extensive rice mapping initiatives. In Australia, geospatial technologies, including remote sensing and geographic information systems (GIS), are widely embraced, particularly for mapping crops like sugarcane and oil palms. At the micro-level, geospatial techniques are predominantly employed for mapping groundwater resources, analyzing drainage patterns, implementing variable rate fertilizer delivery, and managing pesticides and insecticides. The integration of geospatial technologies in agriculture brings several advantages, including increased yields,

ISBN : 9788119821327

efficient resource management, outcome prediction, and enhanced farming practices (Xiangjian, M. and Gang, L. 2007)

Role in Carbon Sequestration

Carbon sequestration (CS) involves the secure transfer and storage of atmospheric CO2 into long-lived carbon reservoirs, mitigating its release into the atmosphere. The combination of remote sensing (RS) and geographic information systems (GIS) is a powerful tool for advanced spatial and temporal analysis of high-resolution data, contributing significantly to research efforts aimed at managing the global carbon cycle. Integrated RS-GIS serves as a Decision Support System (DSS) tool in CS administration and monitoring, enabling the quantification of climatic and soil conditions across geographical and temporal variations (Niu and Duiker, 2006). Various remote sensing methods, including Synthetic Aperture Radar (SAR), Light Detection and Ranging (LiDAR), and satellite sensors such as Landsat, SPOT, and Ikonos, have been employed to map carbon stocks (Goetz et al., 2009). In an area double-planted with wheat-corn under a Typical tillage regime, the soil CS potential ranged between 0.16-0.43 Pg of C without significantly affecting crop yields (Thomson et al., 2006). The study also suggests that vegetation recovery enhances the CS potential of eroded soils. The development of C-Lock, a system standardizing the estimation of agricultural carbon sequestration credits, utilized a regional-scale GIS as its operating platform. This system incorporates the Century biogeochemical model, facilitating the simulation of carbon, nutrient, and water dynamics in various ecosystems and supporting precision carbon management. The synergistic role of RS and GIS technologies is evident in CS management, with studies highlighting

ISBN : 9788119821327

their effectiveness in estimating biomass, delineating geographic variability, predicting carbon stocks and revenues, and outlining management strategies at local and regional scales for climate change mitigation. The integration of RS and GIS in precision carbon management is expected to become increasingly prominent in the future

Technological Challenges in Agriculture Sector

The agriculture sector confronts various technological challenges that can impede its growth and advancement. Several key challenges include:

1. Limited Access to Technology: Small-scale farmers in developing countries often lack access to modern agricultural technologies due to insufficient infrastructure, resources, and knowledge.
2. High Cost of Technology: Advanced agricultural technologies, including precision farming equipment and remote sensing tools, can be prohibitively expensive for many farmers, limiting their adoption.
3. Lack of Data Sharing: Agriculture data is frequently fragmented and not easily accessible, hindering the development of comprehensive solutions that cater to the diverse needs of farmers.
4. Technical Knowledge Gaps: Many farmers lack the necessary technical knowledge and skills to effectively utilize new technologies in their farming practices, creating a barrier to adoption.
5. Compatibility Issues: Agricultural technologies may not align with existing farming practices or infrastructure, making integration into current systems challenging.

ISBN : 9788119821327

6. Dependence on External Factors: Some agricultural technologies, such as weather forecasting tools, heavily rely on external factors like data from satellites or weather stations, which may not always be available or accurate.
7. Cybersecurity Risks: Agriculture technology systems are susceptible to cybersecurity risks, including data breaches and cyberattacks, which can compromise sensitive information and disrupt farming operations.

Addressing these technological challenges necessitates collaborative efforts involving government, the private sector, and academic institutions. This collaboration can foster research and development, facilitate knowledge sharing, and enhance access to affordable and reliable technology solutions for farmers

Conclusion

Geospatial technology has brought about a revolutionary transformation in the agriculture sector, empowering farmers with valuable data and insights for more informed decision-making in crop management. The applications of geospatial technology, ranging from precision farming to crop monitoring and field mapping, have contributed to increased yields, cost reduction, and improved sustainability in agriculture. Despite these advancements, the sector still grapples with technological challenges, including limited access, high costs, and technical knowledge gaps. Addressing these challenges requires a concerted effort involving collaboration between government entities, the private sector, and academic institutions. This collaborative approach aims to foster research and development, facilitate knowledge sharing, and ensure access to affordable and

ISBN : 9788119821327

reliable technology solutions for farmers. Continued investments in geospatial technology and its integration into agriculture hold the potential for more efficient and sustainable farming practices, securing food production and environmental sustainability for future generations. Geospatial technologies, such as remote sensing, GPS, and GIS, provide farmers with detailed information about site-specific demands on their farms. This information becomes a valuable resource for developing and implementing management strategies that optimize input usage, maximizing production and revenue. By leveraging geospatial technologies, farmers gain access to a repository of information that enables them to make informed decisions, ensuring effective and efficient farm management and enhancing overall productivity. To harness the full potential of these technologies, farmers are encouraged to integrate them with their existing knowledge and expertise (Anjna *et al*., 2023).

References

1. Anjna, G. R. L., Raut. and Hansnath, K. 2023. Role of Geospatial Technology in Agriculture. 3:8.
2. Brisco, B., Brown, R., Hirose, J., McNairn, H. and Staenz, K. 2012. Precision Agriculture and the Role of Remote Sensing: A Review.
3. Esri (2008). GIS for Sustainable Agriculture. GIS Best Practices. New York: ESRI Publications
4. Grisso, B. 2009. Precision Farming: A Comprehensive Approach.
5. Sohne, W., Heinze, O. and Groten, E. 1994. Integrated INS/GPS System for High Precision Navigation Applications. Record-IEEE PLANS, Position Location and Navigation Symposium. **35**(2): 310-313.

ISBN : 9788119821327

6. Xiangjian, M. and Gang, L. 2007. Integrating GIS and GPS to Realise Autonomous Navigation of Farm Machinery. *New Zealand Journal of Research*. **50**(1): 807-812

ISBN : 9788119821327

CHAPTER - 16

Geoinformatics, Nanotechnology and Precision Farming

Insha Jan, Tanveer Ahmad Ahngar*and Arzeena Shabir

Division of Agronomy, Faculty of Agriculture, Wadura, SKUAST-K, Jammu and Kashmir, India

****Corresponding author: tanveeragron@gmail.com***

Abstract

Agriculture is poised for a technology revolution in the twenty-first century, one that could transform farming as a whole and increase production. In this revolutionary story, geoinformatics, nanotechnology, and precision agriculture have come to prominence as key figures guiding the way towards a more efficient and sustainable agricultural future. Farmers and academics are given tremendous tools for the study of geoinformatics—the collection, analysis, and interpretation of geographic data—to comprehend the spatial dynamics of agricultural landscapes. In the meantime, novel approaches to pest management and targeted nutrition delivery are made possible by nanotechnology, which acts at the molecular and atomic levels and opens up hitherto unheard-of possibilities for precise manipulation in the agricultural domain.

Introduction

Geoinformatics, the science of utilizing geographic information through cutting-edge technologies, provides farmers with a strong lens to perceive, analyse, and comprehend the intricacies of their agricultural environments. Farmers can use satellite imagery, Geographic Information Systems (GIS), and spatial data analytics to make well-informed decisions on how best to manage their land, care for their crops, and allocate resources. Thus, geoinformatics serves as

ISBN : 9788119821327

the base, providing a spatial framework for the complex dance of factors influencing contemporary agriculture.

With its ability to modify matter at the molecular level at the nanoscale, nanotechnology has become a game changer in agriculture. This tiny science presents new methods for improving nutrient delivery, reducing pest pressure, and increasing agricultural output. Targeted interventions made possible by nano-enabled technologies enable farmers to tackle issues head-on, lessen their negative effects on the environment, and usher in a new era of precision-driven and sustainable agriculture. Precision farming is an emerging methodology designed to link management actions to site-specific soil and crop conditions, and place inputs of fertilizers, herbicides, and pesticides where they are most needed to maximize farm efficiency and minimize environmental contamination. The core technologies in precision agriculture are GIS, GPS, and remote sensing.

The enormous use of agricultural inputs including machinery, pesticides, water, and others brought about by conventional farming techniques has had a significant impact on the environment, including pollution from greenhouse gas emissions. Research indicates that the two most significant obstacles to the adoption of precision agriculture are those related to education and the economy.

Among the economic challenges, Precision Agriculture and startup expenses have a greater influence than the other problems. Precision Farming avoids soil degradation while increasing agricultural productivity. Precision Farming promotes optimal use of water resources and lessens the need for chemical application in crop cultivation. It is also beneficial in fostering positive attitudes, transforming the socioeconomic standing of farmers, and disseminating contemporary farming techniques to increase production quality, quantity, and lower costs.

Better resource management made possible by geospatial technology increases the likelihood of increased yields and financial

ISBN : 9788119821327

returns from agricultural operations. The agricultural management in this approach is customized to the unique conditions present in each field. Applying fertilizer is a crucial process that needs to be done carefully and effectively in order to get the most return on investment. The goal of geospatial technologies for fertilizer recommendations is to maximize all available nutrient resources in concert for crop management that is site-specific. These kinds of methods are used to minimize adverse environmental effects, raise yield, and lower production costs.

Foundations of precision agriculture

It is believed that the phrase "precision agriculture" was originally used in 1990 to refer to a Montana State University-sponsored workshop that took place in Great Falls, Montana. The phrases "site-specific agriculture" or "site-specific crop management" were in use prior to this. The word "site-specific management" was used in the titles of the first two international conferences on what is now known as "precision agriculture," but the term "precision agriculture" was first used at the third conference in 1996. The phrase "precision agriculture" gained popularity by the mid-1990s to describe what is now known as modern precision agriculture, which originated in the USA in the 1980s. Farmers have used precision agriculture from the beginning of agriculture. Farmers planted crops where the conditions were most optimum by dividing their land into smaller sections that they were familiar with. Just as modern subsistence farmers do, the early farmers carefully tended their land to ensure that they produced enough food for their families—it was a matter of life or death.

ISBN : 9788119821327




Geographically and chronologically, the adoption of precision agriculture technologies has been uneven. According to the economic theory of induced innovation, new technologies that optimize the use of limited productive resources will be created and implemented in those areas. In fact, the quickest uptake of precision agriculture technologies has occurred in areas with relatively cheap labor costs compared to relatively inexpensive land and capital. The uneven adoption rate of precision agriculture in certain areas is linked to regular cycles for updating the costly machinery that houses many of the technology used in precision agriculture. In nations with enough of land and labor, adoption is expected to continue, and it will pick up speed during periods of high commodity prices and low financing rates.

Nanotechnology in Agriculture:

As this section has shown, nanomaterials offer a wide range of possible uses in agriculture to boost crop productivity and improve soil health. Here, we highlight several advancements in the fields of nanopesticides, nanofertilizers, nanobiosensors, and nano-enabled soil contamination remediation.

ISBN : 9788119821327



Nanofertilizers:

Nano fertilizers improve crop yield and quality with higher nutrient use efficiency while reducing the cost of production and thus, contribute towards agricultural sustainability. Introduction of nanotechnology in plant nutrition facilitates the development of slow/controlled release fertilizers which improve the fertilizer use efficiency and reduce the losses of nutrients to the environment making them ecofriendly. Based on the current data, it is clear that because of their high use efficiency, nanofertilizers can minimize the amount of fertilizer needed, hence reducing the environmental effect caused by nutrient losses. This idea comes at a perfect time to protect the environment and assure global food security. However, research into the economic viability of nanofertilizers is still necessary for lucrative and sustainable agriculture.

Nanopesticides:

Commercial agriculture regularly uses pesticides, and the task of creating new, effective, and target-specific pesticides seldom ceases. Although the potentially harmful effects of synthetic pesticides seem to be lessened by biopesticides, their sluggish and environment-

ISBN : 9788119821327

dependent efficacy against pests limits their application. There is a chance that nano pesticides can get beyond these restrictions. In the presence of appropriate NMs, slow degradation and regulated release of active ingredients can provide long-term, efficient pest control. Consequently, nano pesticides can reduce the need for synthetic chemicals and the hazards they pose to the environment while also being crucial for the sustainable and efficient control of a variety of pests. To boost their effectiveness, nano pesticides act differently from traditional pesticides.

Many agricultural uses, including nano pesticides, nano fertilizers, nano biosensors, and environmental cleanup agents, have been made possible by nanotechnology. Nonetheless, a thorough comprehension of the harmful consequences of nanoparticles on plant development and growth is needed. Crop plants experience changes in morphology and physiological processes as a result of the increased nanoparticle concentration. Increased nanoparticle concentrations in the root zone reduce the growth of leaves and biomass production, change the intake of water and nutrients, hinder seed germination, and limit root development. The fate of nanomaterials and their environmental impacts continue to be a major challenge in the fields of agricultural and environmental sciences. Additionally, oxidative outburst caused by nanomaterial toxicity results in chloroplast disorganization, reduced photosynthesis, membrane disruption, cellular damage, and altered gene expression.

Geoinformatics in Agriculture:

Modern agriculture is increasingly dependent on geoinformatics technology, particularly those related to remote sensing, GIS, and GPS. The collection of data on agricultural operations, such as land use, soil condition, weather conditions, etc., that are crucial for site characterization and subsequently site selection for farming, has been transformed by advances in remote sensing. To enable prompt decision-

ISBN : 9788119821327

making, agricultural data must be collected, analysed, and managed using geographic information systems. Layers of spatial data from field surveys, current maps, and remote sensors will be included in the database. The capacity to combine different data sets from these different sources guarantee that the different data sources are compatible as possessed by Geographic Information Systems.

The integration of remote sensing, Geographic Information System and Global Positioning System technologies have taken agriculture to the space age. This has given rise to a new concept variously termed Precision Agriculture (PA), Precision Farming (PF) or Site Specific Crop Management (SCCM). To enable prompt decision-making, agricultural data must be collected, analysed, and managed using geographic information systems. Layers of spatial data from field surveys, current maps, and remote sensors will be included in the database. The capacity to combine different data sets from these different sources and guarantee that the different data sources are compatible is possessed by Geographic Information Systems.



Advancements in Precision Farming, Nanotechnology, and Geoinformatics:

GIS has experienced significant changes in data collecting, data administration, technological platform, application development, and

ISBN : 9788119821327

project execution as a result of the advancement of IT technology. The storage, administration, and analysis of substantial amounts of geographically dispersed data are made possible by GIS applications. These statistics are linked to the corresponding geographic characteristics. Different data types occupying the same geographic location can be managed by a GIS. Despite sharing an identical geographic area, predators, prey, and vegetation can all be mapped as unique elements. We should anticipate a sharp rise in food production in the future given these developments and the upcoming release of new, cutting-edge technology. The face of modern farming will alter.

By concentrating on the minute spatial variations within a field, precision farming maximizes productivity. Variable Rate Application is the concept that sets precision farming apart from conventional agricultural practices. When, when, and in what quantities are needed are the inputs provided. Growers require distinct Variable-Rate Technologies (VRA) at every step and comprehensive, geographical data about their farms in order to use the VRA approach. It is important to highlight that precision farming, in terms of fertilizer application, lowers input consumption while increasing output quantities and quality. It also lessens pollution of the soil, water, and air by using fewer chemical pesticides and fertilizers. Large farms, small farms, and organizations collaborating with several producers can all benefit from precision farming.

ISBN : 9788119821327



In a short amount of time, nanotechnology has dominated all scientific disciplines, including chemistry, biology, and physics. Now, because of the obvious and established advantages of microscaling, it is anticipated to have a significant influence on manufacturing technology. Nanotechnology is currently creating entirely new fields in every sector, including computing, information technology, engineering, medical, agriculture, and food. The terms nanocomputer, nanoengineering, nanoinformatics, nanobiotechnology, nanomedicine, nanoagriculture, and nanofood industries are commonly used to refer to these sectors. While the most innovative and affordable technologies are being launched in the fields of materials science and mechanical sciences, the most dazzling discoveries are being produced in the field of nanomedicine. The amazing ways that nanotechnology is being used in the industrial world lead one to believe that in the not too distant future, its products will be used in every industry imaginable. Nonetheless, we must exercise caution and educate ourselves on the potential health risks to living things as well as environmental and pollution issues that may come from the improper application of nanotechnology. This is significant since sustainability is a factor that is being thought about more and more globally. Therefore, it is possible to ensure that nanotechnology will have a profitable future by combining the sustainability component with nanotechnology.

ISBN : 9788119821327

Conclusion:

The amalgamation of precision agriculture, geoinformatics, and nanotechnology has become a trifecta of innovation in the rapidly changing field of agriculture, altering the fundamentals of conventional agricultural methods. It is clear from the conclusion of this investigation into the revolutionary potential of these technologies that integrating them will be essential to achieving hitherto unheard-of levels of agricultural resilience, sustainability, and efficiency.

A Confluence of Precision and Miniaturization: Nanotechnology's Impact on Agriculture:

The development of nanotechnology's use in agriculture has been nothing short of remarkable. The manipulation of matter at the nanoscale is at the core of this scientific revolution, offering a paradigm leap in our understanding of soil health, nutrient delivery, and crop protection. Precision and focus have been possible in addressing issues that have long beset agriculture thanks to nanotechnology. In addition to increasing efficacy, the use of nanoparticles in crop protection has reduced environmental impact. Pesticide nano-formulations with regulated release show better efficacy and less environmental damage. Similar to this, plants are now able to utilize nutrients to their fullest potential thanks to the precise delivery of nutrients via nanocarriers, which reduces waste and environmental runoff. Agriculture and nanotechnology are a symbiotic couple, with the precision of nanoscale interventions perfectly matching the needs of contemporary farming.

Navigating Agricultural Terrain with Geoinformatics:

Agriculture's complicated terrain has shown to be best navigated with the invaluable assistance of geoinformatics. Geoinformatics has given farmers access to practical insights through the use of satellite images, which offers an aerial view of fields, and Geographic Information Systems (GIS), which facilitates spatial analysis. Farm

ISBN : 9788119821327

management techniques have been completely transformed by the capacity to track crop health, evaluate land use patterns, and make well-informed decisions using geospatial data. Precision agriculture, fostered by geoinformatics, has transformed fields into dynamic, data-driven ecosystems. Farmers having access to geospatial data may increase overall production, reduce environmental hazards, and allocate resources optimally. The combination of geoinformatics with precision agriculture is more than simply a technology partnership; it's a calculated move that gives farmers the potential to produce with never-before-seen levels of environmental awareness, flexibility, and accuracy.

Precision Farming: A Harmony of Data-Informed Effectiveness:

Precision agriculture is at the crossroads of geoinformatics and nanotechnology, bringing data-driven efficiency to a harmonious whole. Precision agriculture is the conductor that unifies the capabilities of nanotechnology and geoinformatics into a smooth and effective agricultural performance. It includes everything from GPS-guided machinery that ensures accurate field coverage to real-time monitoring systems that optimize irrigation. In a time when environmental sustainability and resource conservation are of the utmost importance, the capacity to customize agricultural techniques to the unique requirements of every square meter of land is not just a luxury but also a need. Precision agriculture represents the forefront of a new era in farming, one in which every drop of water, every ounce of fertilizer, and every ray of sunlight is harnessed with surgical precision to maximize agricultural output. Precision agriculture is driven by the insights from geoinformatics and the precision of nanotechnology.

A Vision for the Future:

In conclusion, the combination of geoinformatics, precision agriculture, and nanotechnology represents a vision for the future of agriculture, not only a technical development. It's a future in which

ISBN : 9788119821327

farmers have the means to purposefully and precisely negotiate the complexity of their fields, where all inputs are maximized and all results are sustainable. In this future, innovation and responsibility coexist, with precision agriculture, geoinformatics, and nanotechnology serving as the enablers for a robust and thriving agriculture industry. As we stand at the intersection of these transformative forces, it is our collective responsibility to steer the course toward a future where technology serves as a force for good, enhancing not only the productivity of our fields but also the health of our planet. The trip is far from over, and there is a plethora of opportunities for creativity, teamwork, and beneficial effects in agriculture. Let this trio serve as our compass to lead us to a frontier in agriculture that is more robust, efficient, and sustainable.

References:

1. Bahadur, P. S., Jaiswal, S., Srivastava, R., & Kumar, A. (2021, November). Advanced application of nanotechnology in engineering. In *2021 International Conference on Technological Advancements and Innovations (ICTAI)* (pp. 92-95). IEEE.
   https://agribooks.co/geoinformatics-and-its-branches/
   https://www.linkedin.com/pulse/farming-20-rise-precision-agriculture-digital-world-abhishek-singh
   https://www.sciencedirect.com/science/article/pii/S0981942823007556
2. Meena, B. R., Jatav, H. S., Dudwal, B. L., Kumawat, P., Meena, S. S., Singh, V. K., ... & Sathyanarayana, E. (2022). Fertilizer Recommendations by Using Different Geospatial Technologies in Precision Farming or Nanotechnology. *Ecosyst. Serv*, *14*, 241-257.
3. Nellis, M. D., Price, K. P., & Rundquist, D. (2009). Remote sensing of cropland agriculture. *The SAGE handbook of remote sensing*, *1*, 368-380.

ISBN : 9788119821327

4. Oliver, M. A., Bishop, T. F., & Marchant, B. P. (Eds.). (2013). *Precision agriculture for sustainability and environmental protection* (pp. 1-283). Abingdon: Routledge.
5. Rilwani, M. L., & Gbakeji, J. O. (2009). Geoinformatics in agricultural development: challenges and prospects in Nigeria. *Journal of Social Sciences*, *21*(1), 49-57.
6. Song, W., & Wu, C. (2021). Introduction to advancements of GIS in the new IT era. Annals of GIS, 27(1), 1-4.
7. Swinton, S. M., & Lowenberg-Deboer, J. (2001, June). Global adoption of precision agriculture technologies: Who, when and why. In *Proceedings of the 3rd European conference on precision agriculture* (Vol. 2, pp. 557-562). Citeseer.
8. Tiwari, A., & Jaga, P. K. (2012). Precision farming in India–A review. *Outlook on Agriculture*, *41*(2), 139-143.
9. Udoumoh, U. I., Ikrang, E. G., & Ehiomogue, P. O. (2021). Precision farming and fertilizer recommendation using geographic information system (GIS): A review. *Int. J. Agricult. Earth Sci*, *7*(2), 1-8.
10. Usman, M., Farooq, M., Wakeel, A., Nawaz, A., Cheema, S. A., ur Rehman, H., ... & Sanaullah, M. (2020). Nanotechnology in agriculture: Current status, challenges and future opportunities. *Science of the Total Environment*, *721*, 137778.

ISBN : 9788119821327